

AUTHOR INDEX

- Ali, H.M., Comparative bioavailability of eight brands of ampicillin, 301
- Ali, S.L., see Möller, H., 157
- Alpar, O.H., Bamford, J.B. and Walters, V., The in vivo incorporation and release of hydroxocobalamin by liposomes, 349
- Al-Shora, H., Said, S. and Hammad, A.-L., Sustained release from inert matrixes II. Effect of polyethylene glycols on theophylline release, 77
- Attwood, D., see Wood, J.M., 189
- Augsburger, L.L., see Mehta, A.M., 327
- Baillie, A.J., see Onaga, I.C., 89
- Bamford, J.B., see Alpar, O.H., 349
- Baudhuin, P., see Kante, B., 45
- Bodor, N., Zupan, J. and Selk, S., Improved delivery through biological membranes VII. Dermal delivery of cromoglycic acid (cromolyn) via its prodrugs, 63
- Bryan, A.J., Kaur, R., Robinson, G., Thomas, N.W. and Wilson, C.G., Histological and physiological studies on the intestine of the rat exposed to solutions of Myrj 52 and PEG 2000, 145
- Bundgaard, H. and Johansen, M., Pro-drugs as drug delivery systems XV. Bioreversible derivatization of phenytoin, acetazolamide chlorzoxazone and various other NH-acidic compounds by N-aminomethylation to effect enhanced dissolution rates, 129
- Bundgaard, H. and Larsen, C., Pro-drugs as drug delivery systems XVII. Esters of 4-hydroxybutyric acids as potential pro-drug types for γ -lactones, 169
- Bundgaard, H. and Hansen, J., Studies on the stability of corticosteroids VI. Kinetics of the rearrangement of betamethasone-17-valerate to the 21-valerate ester in aqueous solution, 197
- Bundgaard, H., see Johansen, M., 119
- Collett, J.H., see Wood, J.M., 189
- Couvreur, P., see Kante, B., 45
- D'Arcy, P.F., see McElroy, J.C., 83
- El-Khordagui, L.K., Saleh, A.M. and Khalil, S.A., Diazepam-sodium salicylate solution: dilution with intravenous fluids, in vitro haemolytic activity and protein binding, 111
- Farsam, H., Yahya-Saeed, H.H. and Fawzi, A., Spectrophotometric determination of codeine in pharmaceutical preparations, 343
- Fawzi, A., see Farsam, H., 343
- Franz, J.M., Vonderscher, J.P. and Voges, R., Contribution to the intestinal absorption of ergot peptide alkaloids, 19
- Fujimoto, Y., see Nakano, N.I., 335
- Grant, D.J.W., see Liversidge, G.G., 211
- Gstrein, K., see Nachtmann, F., 55
- Guiot, P., see Kante, B., 45
- Hammad, A.-L., see Al-Shora, H., 77
- Hansen, J., see Bundgaard, H., 197
- Hikal, A.H., Effect of polysorbate 80 on the apparent partition coefficient of drugs and on their intestinal absorption in the rat II. Phenobarbital, 205
- Hirai, S., Yashiki, T., Matsuzawa, T. and Mima, H., Absorption of drugs from the nasal mucosa of rat, 317
- Hölzer, A.W. and Sjögren, J., Friction coefficients of tablet masses, 269
- Huizinga, T., see Schoonen, A.J.M., 29
- Jacobs, G.P., The radiation sterilization of cefoxitin sodium, 279
- Jamali, F., see Mehvar, S.R., 293
- Johansen, M. and Bundgaard, H., Pro-drugs as drug delivery systems XIII. Kinetics of decomposition of N-Mannich bases of salicylamide and assessment of their suitability as possible pro-drugs for amines, 119
- Johansen, M., see Bundgaard, H., 129
- Jung, D. and Øie, S., Decreased in vivo acetylation of sulfisoxazole in the rabbit in the presence of 4-N-acetyl-sulfisoxazole, 179
- Kaneniwa, N., see Watari, N., 307

- Kante, B., Couvreur, P., Lenaerts, V., Guiot, P., Roland, M., Baudhuin, P. and Speiser, P., Tissue distribution of [^3H]actinomycin D absorbed in polybutylcyanoacrylate nanoparticles, 45
- Kaur, R., see Bryan, A.J., 145
- Kellaway, I.W. and Najib, N.M., The effect of hydrophilic polymers on the electrophoretic mobility of suspended particles, 285
- Kent, J.S., Mroszczak, E.J., Roe, R.L. and Runkel, R., Bioavailability and in vitro dissolution of tiopinac from solution and capsule formulations, 245
- Khalil, S.A., see El-Khordagui, L.K., 111
- Kraak, J.C., see Tomlinson, E., 225
- Krasowska, H., Effect of micellar solubilization of the gastrointestinal absorption of indomethacin in the rat, 137
- Larsen, C., see Bundgaard, H., 169
- Lenaerts, V., see Kante, B., 45
- Liversidge, G.G., Grant, D.J.W. and Padfield, J.M., Influence of physicochemical interactions on the properties of suppositories. I. Interactions between the constituents of fatty suppository bases, 211
- Matsuzawa, T., see Hirai, S., 317
- McElnay, J.C., D'Arcy, P.F. and Throne, O., Effect of antacid constituents, kaolin and calcium citrate on phenytoin absorption, 83
- Mehta, A.M. and Augsburger, L.L., A preliminary study of the effect of slug hardness on drug dissolution from hard gelatin capsules filled on an automatic capsule-filling machine, 327
- Mehvar, S.R. and Jamali, F., Dipyrone-ethyl-bisouacetate interaction in man, 293
- Mima, H., see Hirai, S., 317
- Möller, H., Ali, S.L. and Steinbach, D., Pharmaceutical and biological availability of sustained release preparations of potassium chloride, 157
- Moody, R.R. and Taylor, R.B., An HPLC assay for the determination of proguanil hydrochloride in tablets, 177
- Moolenaar, F., see Schoonen, A.J.M., 29
- Mroszczak, E.J., see Kent, J.S., 245
- Nachtmann, F. and Gstrein, K., Comparison of HPLC and some official test methods for different penicillins, 55
- Najib, N.M., see Kellaway, I.W., 285
- Nakano, N.I., Yamaguchi, S., Shimamori, Y. and Fujimoto, Y., Fractionation of a human serum albumin preparation and the salicylate binding characteristics, 335
- Noonan, P.K., see Wester, R.C., 99
- Öie, S., see Jung, D., 179
- Onaga, I.C. and Baillie, A.J., The toxicity of liposomal chloramphenicol for *Tetrahymena pyriformis*, 89
- Padfield, J.M., see Liversidge, G.G., 211
- Poppe, H., see Tomlinson, E., 225
- Reuvers, K.A., see Schoonen, A.J.M., 29
- Robinson, G., see Bryan, A.J., 145
- Roe, R.L., see Kent, J.S., 245
- Roland, M., see Kante, B., 45
- Runkel, R., see Kent, J.S., 245
- Said, S., see Al-Shora, H., 77
- Saleh, A.M., see El-Khordagui, L.K., 111
- Schoonen, A.J.M., Moolenaar, F., Reuvers, K.A. and Huizinga, T., Release of drugs from fatty suppository bases II. A rate-limiting interfacial process, 29
- Selk, S., see Bodor, N., 63
- Shimamori, Y., see Nakano, N.I., 335
- Sjögren, J., see Hölder, A.W., 269
- Speiser, P., see Kante, B., 45
- Spring, M.S., The effects of some process variables on the concentration of minor components in tablet granules, 261
- Steinbach, D., see Möller, H., 157
- Taylor, R.B., see Moody, R.R., 177
- Thomas, N.W., see Bryan, A.J., 145
- Throne, O., see McElnay, J.C., 83
- Tomlinson, E., Poppe, H. and Kraak, J.C., Thermodynamics of functional groups in reversed-phase high performance liquid-solid chromatography, 225
- Voges, R., see Franz, J.M., 19
- Vonderscher, J.P., see Franz, J.M., 19
- Walters, V., see Alpar, O.H., 349
- Watari, N. and Kaneniwa, N., Dissolution of slightly soluble drugs VIII. In vivo drug dissolution in human subjects and rabbits, 307
- Wester, R.C. and Noonan, P.K., Relevance of animal models for percutaneous absorption, 99

- Wilson, C.G., see Bryan, A.J., 145
- Wood, D.A., Biodegradable drug delivery systems, 1
- Wood, J.M., Attwood, D. and Collett, J.H., The swelling properties of poly(2-hydroxyethyl methacrylate) hydrogels polymerized by gamma-irradiation and chemical initiation, 189
- Yahya-Saeb, H.H., see Farsam, H., 343
- Yamaguchi, S., see Nakano, N.I., 335
- Yashiki, T., see Hirai, S., 317
- Zupan, J., see Bodor, N., 63