

AUTHOR INDEX

- Acree, W.E., Jr., Solubility of biphenyl in binary solvent mixtures, 47
- Akaho, E., see Kagayama, A., 247
- Allen, J.C., see Morton, A.K., 39
- Al-Shora, H.I., see Gouda, H.W., 213
- Arnold, E., see Bundgaard, H., 79
- Badian, M., Le Normand, Y., Rupp, W. and Zapf, R., There is no interaction between dipyrone (metamizol) and the anticoagulants, phenprocoumon and ethylbiscoumacetate, in normal caucasian subjects, 9
- Barry, B.W., see Southwell, D., 299
- Barzegar-Jalali, M., Theoretical justification for the midpoint back-extrapolation method of estimating apparent volume of distribution of drugs after a short intravenous infusion, 353
- Blanchard, J., Perry, D.F. and Walson, P.D., Factors influencing the transfer of xanthines across the everted rat jejunum, 259
- Brenner, G.S., see Zografi, G., 99
- Brodin, A., see Nicklasson, M., 149
- Bundgaard, H., Larsen, C. and Thorbeck, P., Prodrugs as drug delivery systems. XXVI. Preparation and enzymatic hydrolysis of various water-soluble amino acid esters of metronidazole, 67
- Bundgaard, H., Larsen, C. and Arnold, E., Prodrugs as drug delivery systems. XXVII. Chemical stability and bioavailability of a water-soluble prodrug of metronidazole for parenteral administration, 79
- Bundgaard, H., see Buur, A., 325
- Buur, A. and Bundgaard, H., Prodrugs as drug delivery systems. XXVIII. Structural factors influencing the rate of hydrolysis of oxazolidinedines—a potential prodrug type, 325
- Carlotti, M.E., see Gasco, M.R., 235
- Carpignano, R., see Gasco, M.R., 235
- Chou, R.C., Ramzan, I.M. and Levy, G., Effect of pregnancy or renal dysfunction on differences in plasma protein binding of phenytoin in peripheral blood and blood exiting the liver of rats, 217
- Chow, A.H.-L., see Grant, D.J.W., 25
- Collier, P.S., The possible range of values for apparent volume of distribution at steady-state when disposition may be characterized by a tri-exponential function, 345
- Corrigan, O.I., Holohan, E.M. and Sabra, K., Amorphous forms of thiazide diuretics prepared by spray-drying, 195
- Daly, P.B., Davis, S.S. and Kennerley, J.W., The effect of anionic surfactants on the release of chlorpheniramine from a polymer matrix tablet, 201
- Davis, S.S., see Wilson, C.G., 1
- Davis, S.S., see Daly, P.B., 201
- de Blaey, C.J., see Fokkens, J.G., 127
- de Haan, F.H.N. and Jansen, A.C.A., The hydrodynamics in a simple transport vessel in two-phase systems of organic solvent-water, 311
- Drabik, B.R., see Miller, S.C., 269
- Edwards, H.E., see Morton, A.K., 39
- El-Fattah, S.A. and Khalil, S.A.H., Variations in dissolution rates of sugar-coated chlorpromazine tablets, 225
- El-Khawas, M., see Khalil, S.A.H., 157
- Fairbrother, J.E., see Grant, D.J.W., 25
- Fokkens, J.G. and de Blaey, C.J., Drug release from non-aqueous suspensions. II. The release of methylxanthines from paraffin suspensions, 127
- Gasco, M.R., Trotta, M., Carlotti, M.E. and Carpignano, R., Contribution of carboxylic acids to the permeability of imipramine, 235
- Gouda, H.W., Moustafa, M.A. and Al-Shora, H.I., Effect of storage on nitrofurantoin solid dosage forms, 213
- Grant, D.J.W., Mehdizadeh, M., Chow, A.H.-L. and Fairbrother, J.E., Non-linear van't Hoff solubility-temperature plots and their pharmaceutical interpretation, 25
- Guy, R.H. and Hadgraft, J., A theoretical description of the effects of volatility and substantivity on percutaneous absorption, 139

- Hadgraft, J., see Guy, R.H., 139
 Hardy, J.G., see Wilson, C.G., 1
 Hirsch, C.A., see Pearlman, R., 53
 Hiura, M., Kawashima, H. and Kaneniwa, N., Hepatic first-pass metabolism of hexobarbital in rabbits and dogs, 185
 Hollenbeck, R.G., see Tabibi, S.E., 169
 Holohan, E.M., see Corrigan, O.I., 195
 Hussain, A., see Kagayama, A., 247
 Jansen, A.C.A., see de Haan, F.H.N., 311
 Kagayama, A., Mustafa, R., Akaho, E., Khawam, N., Truelove, J. and Hussain, A., Mechanism of diffusion of compounds through ethylene vinyl acetate copolymers. I. Kinetics of diffusion of 1-chloro-4-nitrobenzene, 3,4-dimethylphenol and 4-hexylresorcinol, 247
 Kaneniwa, N., see Hiura, M., 185
 Kawashima, H., see Naito, S., 117
 Kawashima, H., see Hiura, M., 185
 Kawashima, Y., Lin, S.Y., Ueda, M. and Takenaka, H., Preparation of directly compressible powders of a physical mixture and a complex of theophylline-phenobarbital using spray-drying, 335
 Kennerley, J.W., see Wilson, C.G., 1
 Kennerley, J.W., see Daly, P.B., 201
 Khalil, S.A.H., Mortada, L.M. and El-Khawas, M., The uptake of ampicillin and amoxycillin by some adsorbents, 157
 Khalil, S.A.H., see El-Fattah, S.A., 225
 Khawam, N., see Kagayama, A., 247
 Kontny, M.J., see Zografi, G., 99
 Larsen, C., see Bundgaard, H., 67
 Larsen, C., see Bundgaard, H., 79
 Le Normand, Y., see Badian, M., 9
 Levy, G., see Chou, R.C., 217
 Lewis, G.A., Determination of dissociation constants of sparingly soluble compounds from solubility data, 207
 Lin, S.Y., see Kawashima, Y., 335
 Li Wan Po, A. and Mroso, P.V., Drug-drug incompatibility in the solid state: kinetic interpretation, modelling and prediction, 287
 Mehdizadeh, M., see Grant, D.J.W., 25
 Meltzer, N.M., see Repta, A.J., 277
 Mereish, K.A., see Repta, A.J., 277
 Miller, S.C. and Drabik, B.R., Rheological properties of poloxamer vehicles, 269
 Mizuho, M., see Naito, S., 117
 Mortada, I.M., see Khalil, S.A.H., 157
 Morton, A.K., Edwards, H.E. and Allen, J.C., Some physicochemical and biological characteristics of heparin fractions prepared by gel chromatography, 39
 Moustafa, M.A., see Gouda, H.W., 213
 Mroso, P.V., see Li Wan Po, A., 287
 Mustafa, R., see Kagayama, A., 247
 Naito, S., Tanaka, S., Mizuno, M. and Kawashima, H., Concentrations of human interferons α and β in rabbit body fluids, 117
 Nicklasson, M. and Brodin, A., The relationship between intrinsic dissolution rates and solubilities in the water-ethanol binary solvent system, 149
 Parr, G.D., see Wilson, C.G., 1
 Pearlman, R., Rutherford, B.S., Pozsgai, K.M. and Hirsch, C.A., The isolation and identification of some degradation products of flurandrenolide in Cordran cream, 53
 Periviita, L., see Salminen, L., 17
 Perry, D.F., see Blanchard, J., 259
 Pozsgai, K.M., see Pearlman, R., 53
 Ramzan, I.M., see Chou, R.C., 217
 Rees, J.A., see Wilson, C.G., 1
 Repta, A.J., Sternson, L.A., Mereish, K.A. and Meltzer, N.M., Complexation in the retention of benzoic acids on a bonded xanthine HPLC stationary phase, 277
 Ritchie, K. and Winfield, A.J., An investigation of phenobarbitone-sucrose glass systems, 89
 Rupp, W., see Badian, M., 9
 Rutherford, B.S., see Pearlman, R., 53
 Sabra, K., see Corrigan, O.I., 195
 Salminen, L., Urtti, A. and Periviita, L., Effect of ocular pigmentation on pilocarpine pharmacology in the rabbit eye. I. Drug distribution and metabolism, 17
 Southwell, D., Barry, B.W. and Woodford, R., Variations in permeability of human skin within and between specimens, 299
 Sternson, L.A., see Repta, A.J., 277
 Tabibi, S.E. and Hollenbeck, R.G., Interaction of water vapor and compressible sugar, 169
 Takenaka, H., see Kawashima, Y., 335
 Tanaka, S., see Naito, S., 117
 Taylor, M.J., see Wilson, C.G., 1
 Thorbeck, P., see Bundgaard, H., 67
 Trotta, M., see Gasco, M.R., 235
 Truelove, J., see Kagayama, A., 247

Ueda, M., see Kawashima, Y., 335

Urtti, A., see Salminen, L., 17

Walson, P.D., see Blanchard, J., 259

Wilson, C.G., Parr, G.D., Kennerley, J.W.,
Taylor, M.J., Davis, S.S., Hardy, J.G. and
Rees, J.A., Pharmacokinetics and in vivo scin-
tigraphic monitoring of a sustained release
acetylsalicylic acid formulation, 1

Winfield, A.J., see Ritchie, K., 89

Woodford, R., see Southweil, D., 299

Yang, A.Y.S., see Zografi, G., 99

Zapf, R., see Badian, M., 9

Zografi, G., Kontny, M.J., Yang, A.Y.S. and
Brenner, G.S., Surface area and water vapor
sorption of microcrystalline cellulose, 99