

Erratum

Erratum to “Enhancing effect of polyoxyethylene alkyl ethers on the skin permeation of ibuprofen” [Int. J. Pharm. 209 (2000) 109–119] ☆

Eun-Seok Park, Si-Young Chang, Mikyoung Hahn, Sang-Cheol Chi *

College of Pharmacy, Sungkyunkwan University, Suwon 440-746, South Korea

The publisher regrets that in the above article Fig. 4 was inadvertently omitted, also the legends to figures 4,5,6 and 7 were published incorrectly. The figures and their correct legends are reproduced overleaf.

☆ PII of original article: S0378-5173(00)00559-7.

* Corresponding author. Tel.: +82-31-2907709; fax: +82-31-2907729.

E-mail address: scchi@yurim.skku.ac.kr (S.-C. Chi).




Fig. 4. A histogram showing the dependence of ibuprofen flux (J_s) on EO chain length with a constant alkyl chain length of 12.




Fig. 5. A histogram showing the dependence of ibuprofen flux (J_s) on EO chain length with a constant alkyl chain length of 16.




Fig. 6. A histogram showing the dependence of ibuprofen flux (J_s) on EO chain length with a constant alkyl chain length of 18.




Fig. 7. A histogram showing the dependence of ibuprofen flux (J_s) on EO chain length with a constant alkyl chain length of 1:1 mixture of C16 and C18.