

Erratum

Erratum to “The effect of ethanol on the simultaneous transport and metabolism of methyl *p*-hydroxybenzoate in excised skin of Yucatan micropig”  
[Int. J. Pharm. 236 (2002) 35–42]<sup>☆</sup>

S.Y. Oh, M. Fujii <sup>\*</sup>, Y. Takeda, K. Yoda, N. Utoguchi, M. Matsumoto, Y. Watanabe

*Showa Pharmaceutical University, 3-3165 Higashi-Tamagawa-Gakuen, Machida, 194-8543 Tokyo, Japan*

The publisher regrets that Fig. 1 in the above article was published incorrectly. Please find the corrected figure produced below.




Fig. 1. Cumulative amounts of HBA, HBE and HBM permeated through full-thickness skin of YMP after application of 10 mM HBM without (a) and with 10% ethanol (b). HBM ( $\circ$ ), HBA ( $\bullet$ ), HBE ( $\blacktriangle$ ), HBM<sub>total</sub> ( $\square$ ). Each point represents the mean  $\pm$  S.E. of at least six experiments.

<sup>☆</sup> PII of original article: S0378-5173(02)00005-4.

<sup>\*</sup> Corresponding author. Tel.: +81-42-7211556; fax: +81-42-7233585.

E-mail address: [fujii@ac.shoyaku.ac.jp](mailto:fujii@ac.shoyaku.ac.jp) (M. Fujii).