





International Journal of Pharmaceutics 252 (2003) 281

www.elsevier.com/locate/ijpharm

## **Erratum**

Erratum to "Shape imposed by secondary structure of a polypeptide affects its free diffusion through liquid-filled pores" [Int. J. Pharm. 244 (2002) 1–8]<sup>☆</sup>

Nazila Salamat-Miller, Montakarn Chittchang, Ashim K. Mitra, Thomas P. Johnston\*

Division of Pharmaceutical Sciences, School of Pharmacy, University of Missouri-Kansas City, Katz Pharmacy Building, Room 211 A, 5005 Rockhill Road, Kansas City, MO 64110-2499, USA

The publisher regrets that there were three mistakes in the above article when it was published.

On page 3 left column line 8 mw/(Mn) should have been written as  $(M_{\rm w}/M_{\rm n})$  and in line 41 t=092%:8% should have been written as t=0 (92%:8%), also on page 6 left column line 18  $f_{\rm exp}=t\cdot T/D_{\rm aq}$  should have been written as  $f_{\rm exp}=k\cdot T/D_{\rm aq}$ .

<sup>&</sup>lt;sup>☆</sup> PII of original article S0378-5173(02)00320-4.

<sup>\*</sup> Corresponding author. Tel.: +1-816-235-1624; fax: +1-816-235-5190. *E-mail address:* johnstont@umkc.edu (T.P. Johnston).