



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

International Journal of Pharmaceutics 278 (2004) 475

international  
journal of  
pharmaceutics

[www.elsevier.com/locate/ijpharm](http://www.elsevier.com/locate/ijpharm)

## Corrigendum

### Corrigendum to “Solubility of drugs in aqueous solutions Part 2: binary nonideal mixed solvent” [Int. J. Pharm. 260 (2003) 283–291]☆

E. Ruckenstein\*, I. Shulgin

*Department of Chemical Engineering, State University of New York at Buffalo, Furnas Hall, Amherst, NY 14260, USA*

Available online 14 May 2004

---

The authors regret that in the above published paper there was a misprinting of equation (B.2) in Appendix B. Please find the correct equation below.

$$\begin{aligned} I = & \frac{L_{13}^2(\ln V - \ln(x_1^{b,1-3} + x_3^{b,1-3}L_{13}))}{(V_3^0 - L_{13}V_1^0)^2} + \frac{L_{31}(-\ln V + \ln(x_3^{b,1-3} + x_1^{b,1-3}L_{31}))}{(V_1^0 - L_{31}V_3^0)^2} \\ & + \frac{V_3^0(L_{13}L_{31}V_3^0 - L_{13}V_1^0 + L_{13}L_{31}V_1^0 - L_{31}V_3^0)}{(V_3^0 - L_{13}V_1^0)(L_{31}V_3^0 - V_1^0)(V_3^0 - V_1^0)V} + \frac{(-1 + L_{31})\ln((1 - L_{31})V)}{(V_3^0L_{31} - V_1^0)(V_3^0 - V_1^0)} \\ & + \frac{(x_3^{b,1-3} + x_1^{b,1-3}L_{31})\ln(x_3^{b,1-3} + x_1^{b,1-3}L_{31})}{(L_{31}V_3^0 - V_1^0)V} \end{aligned} \quad (\text{B.2})$$

All numerical calculations in the above paper have been carried out with the correct equation.

---

☆ doi of original article 10.1016/S0378-5173(03)00273-4.

\* Corresponding author: Tel.: +1-716-645-2911x2214; fax: +1-716-645-3822.

E-mail addresses: feaeliru@acsu.buffalo.edu (E. Ruckenstein), ishulgin@eng.buffalo.edu (I. Shulgin).