

Erratum

Erratum to “Tuning the MR properties of blood-stable
pH-responsive paramagnetic liposomes”
[Int. J. Pharm. 274 (2004) 75–83]☆

Knut-Egil Løkling^{a,*}, Roald Skurtveit^b, Knut Dyrstad^b,
Jo Klaveness^a, Sigrid L. Fossheim^b

^a Department of Medical Chemistry, School of Pharmacy, University of Oslo, P.O. Box 1155 Blindern, N-0318 Oslo, Norway

^b Amersham Health AS, P.O. Box 4220 Nydalen, N-0401 Oslo, Norway

Available online 13 May 2004

The publisher regrets that in the above published article there was a typing error in Table 3. Please find the correct table below.

Table 3

Curve fit parameters for various liposomal GdDTPA-BMA formulations in MES-buffered human whole blood (20 min incubation, 37 °C)

Liposome composition	r_1^{\min} (mM ⁻¹ s ⁻¹)	$r_1^{\min,E}$ (mM ⁻¹ s ⁻¹)	pH ₂₀	R^2	$r_1^{\max,E} - r_1^{\min}$ (mM ⁻¹ s ⁻¹)
DMPE/DPSG					
10 mol% DPSG	1.05	1.52	6.30	0.99	0.47
30 mol% DPSG	1.26	1.53	6.04	0.99	0.27
DPPE/DPSG					
10 mol% DPSG	0.62	2.28	6.95	0.99	1.66
20 mol% DPSG ^a	0.43 ± 0.03	2.03 ± 0.06	6.63 ± 0.08	0.99	1.60 ± 0.03
30 mol% DPSG	0.57	2.38	6.52	0.99	1.81
DSPE/DPSG					
10 mol% DPSG	0.26	1.21	6.92	0.99	0.95
30 mol% DPSG	0.56	2.24	6.65	0.99	1.68

^a Arithmetic mean of duplicates ± S.E.M.

☆ doi of original article 10.1016/j.ijpharm.2003.12.027.

* Corresponding author. Tel.: +47-23-18-5605; fax: +47-23-18-6014.

E-mail address: knut-egil.lokling@amersham.com (K.-E. Løkling).