

Corrigendum

Corrigendum to “Role of the mucous/glycocalyx layers in insulin permeation across the rat ileal membrane”
[Int. J. Pharm. 297 (2005) 98–109]

Yoshinobu Aoki^{a,b}, Mariko Morishita^{a,*}, Kozo Takayama^a

^a Department of Pharmaceutics, Hoshi University, Ebara 2-4-41, Shinagawa, Tokyo, 142 8501, Japan

^b Research Center, Kyorin Pharmaceutical Co. Ltd., Nogi 1848, Nogi-machi, Shimotsugo-gun, Tochigi 329-0114, Japan

Available online 19 September 2005

The authors regret that there was an error in Table 2 in the above-published article.

In Table 2, apparent permeability of FD-4 for PBS- and hyaluronidase-treated groups, 6.144 ± 0.354 and 6.004 ± 0.605 , should be corrected to 61.44 ± 3.54 and 60.04 ± 6.05 , respectively.

Table 2

Apparent permeability coefficient of insulin, FD-4 and antipyrine during transport across the rat ileum with or without hyaluronidase pretreatment

Solute	Apparent permeability, P_{app} (10^{-8} cm/s)		Ratio
	PBS	Hyaluronidase	
Insulin	0.358 ± 0.051	$1.429 \pm 0.072^*$	4.0
FD-4	61.44 ± 3.54	60.04 ± 6.05	1.0
Antipyrine	264.8 ± 27.1	272.0 ± 15.8	0.9

Data: mean \pm S.E. ($n = 4-7$).

* $p < 0.05$ against control. There were no significant differences in the groups of FD-4 and antipyrine.

DOI of the original article: [10.1016/j.ijpharm.2005.03.004](https://doi.org/10.1016/j.ijpharm.2005.03.004).

* Corresponding author. Tel.: +81 3 5498 5782;

fax: +81 3 5498 5782.

E-mail address: morisita@hoshi.ac.jp (M. Morishita).