




Contents lists available at [ScienceDirect](#)

International Journal of Pharmaceutics

journal homepage: www.elsevier.com/locate/ijpharm



Corrigendum

Corrigendum to “Drug delivery to the nail following topical application” [Int. J. Pharm. 236 (2002) 1–26]

Sudaxshina Murdan

Department of Pharmaceutics, The School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX, UK

The author regrets that in the above published article there was an error on Page 2, Section 2, line 12.

The sentence is reproduced correctly below.

The nail plate is a thin (0.25–0.6 mm), hard, yet slightly elastic, translucent, convex structure and is made up of approximately 80–90 layers of dead, keratinised, flattened cells which are tightly bound to one another via numerous intercellular links, membrane-coating granules and desmosomes.

The author would like to apologise for any inconvenience this error may have caused readers of this article. The reference for the number of layers of cells in the nail plate is.

Achten, G., Andre, J., Laporte, M., 1991. Nails in light and electron microscopy. *Semin. Dermatol.* 10, 54–64.

DOI of original article: [10.1016/S0378-5173\(01\)00989-9](https://doi.org/10.1016/S0378-5173(01)00989-9).

E-mail address: sudax.murdan@ulsop.ac.uk.