



Pharmaceutical Nanotechnology

Inhibition of bacterial growth and intramniotic infection in a guinea pig model of chorioamnionitis using PAMAM dendrimers

Bing Wang^{a,c}, Raghavendra S. Navath^{b,c}, Anupa R. Menjoge^{b,c}, Bindu Balakrishnan^a, Robert Bellair^b, Hui Dai^{a,c}, Roberto Romero^c, Sujatha Kannan^{a,c,*}, Rangaramanujam M. Kannan^{b,c,**}

^a Department of Pediatrics (Critical Care Medicine), Children's Hospital of Michigan, Wayne State University, Detroit, MI 48201, United States

^b Department of Chemical Engineering and Material Science, Wayne State University, Detroit, MI 48202, United States

^c Perinatology Research Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NICHD, NIH, DHHS, Detroit, MI 48201, United States

ARTICLE INFO

Article history:

Received 24 February 2010

Received in revised form 30 April 2010

Accepted 15 May 2010

Available online 24 May 2010

Keywords:

PAMAM dendrimer

Antimicrobial activity

Gram negative bacteria

Cytotoxicity

Cell membrane

ABSTRACT

Dendrimers have emerged as topical microbicides to treat vaginal infections. This study explores the *in vitro*, *in vivo* antimicrobial activity of PAMAM dendrimers, and the associated mechanism. Interestingly, topical cervical application of 500 µg of generation-4 neutral dendrimer (G₄-PAMAM-OH) showed potential to treat the *Escherichia coli* induced ascending uterine infection in guinea pig model of chorioamnionitis. Amniotic fluid collected from different gestational sacs of infected guinea pigs posttreatment showed absence of *E. coli* growth in the cultures plated with it. The cytokine level [tumor necrosis factor (TNFα) and interleukin (IL-6 and IL-1β)] in placenta of the G₄-PAMAM-OH treated animals were comparable to those in healthy animals while these were notably high in infected animals. Since, antibacterial activity of amine-terminated PAMAM dendrimers is known, the activity of hydroxyl and carboxylic acid terminated PAMAM dendrimers was compared with it. Though the G₄-PAMAM-NH₂ shows superior antibacterial activity, it was found to be cytotoxic to human cervical epithelial cell line above 10 µg/mL, while the G₄-PAMAM-OH was non-cytotoxic up to 1 mg/mL concentration. Cell integrity, outer (OM) and inner (IM) membrane permeabilization assays showed that G₄-PAMAM-OH dendrimer efficiently changed the OM permeability, while G₄-PAMAM-NH₂ and G_{3,5}-PAMAM-COOH damaged both OM and IM causing the bacterial lysis. The possible antibacterial mechanism are G₄-PAMAM-NH₂ acts as polycation binding to the polyanionic lipopolysaccharide in *E. coli*, the G₄-PAMAM-OH forms hydrogen bonds with the hydrophilic O-antigens in *E. coli* membrane and the G_{3,5}-PAMAM-COOH acts as a polyanion, chelating the divalent ions in outer cell membrane of *E. coli*. This is the first study which shows that G₄-PAMAM-OH dendrimer acts as an antibacterial agent.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Dendrimers are emerging as the new topical antimicrobial agents (Svenson and Tomalia, 2005). Recent developments show that the polylysine dendrimers are effective against the simplex herpes virus (Bourne et al., 2000; Svenson and Tomalia, 2005). The polylysine (SPL7013) dendrimer is currently undergoing Phase I and Phase II human clinical trials to test its efficacy against genital herpes and HIV, safety, tolerability, retention and duration of

* Corresponding author at: Department of Pediatrics (Critical Care Medicine), Children's Hospital of Michigan, Wayne State University, Detroit, MI 48201, United States.

** Corresponding author at: Department of Chemical Engineering and Material Science, Wayne State University, Detroit, MI 48202, United States.
Tel.: +1 313 577 3879; fax: +1 313 577 3810.

E-mail addresses: skannan@med.wayne.edu (S. Kannan), rkannan.wsu@gmail.com, rkannan@eng.wayne.edu (R.M. Kannan).

activity (Halford, 2005; Mumper et al., 2009). Different approaches have been used to enhance the antimicrobial activities employing dendrimers and the most conventional being carriers for the antimicrobial agents, as encapsulating or complexing agents for antibacterial agents, by competitively inhibiting the binding of bacteria to the host cells (Cloninger, 2002; Hou et al., 2009) and by exerting the microbial activity by themselves. Quinolone drugs encapsulated in PAMAM dendrimers were active and useful for topical delivery of microbicides (Cheng et al., 2007). Triazine-based antibiotics encapsulated in the dendrimer beads provided enhanced inhibition of microbial growth when compared to the non-dendrimerized triazine antibiotics (Lebreton et al., 2003). The silver complexes and nanocomposites with PAMAM dendrimer resulted in increased antibacterial activity towards the *S. aureus*, *P. aeruginosa* and *Escherichia coli* due to the improved contact of microbes with the organized silver domains (Balogh et al., 2001).

Literature reveals that the dendrimer core, the surface charge and functionality, 3D structure and the size of the dendrimer

are key factors that affect the antibacterial activity. The PAMAM (polyamidoamine) dendrimer with amine terminations exhibited antibacterial activity against gram negative *E. coli*, *P. aeruginosa* and gram positive *S. aureus* (Calabretta et al., 2007; Lopez et al., 2009). Glycodendrimers are known as antimicrobial agents (Cloninger, 2002). Ortega et al. (2008) reported that the amine and ammonium terminated cationic carbosilane dendrimers were more potent towards the gram positive bacteria than the gram negative bacteria. Amine-terminated dendrimers with poly(propyleneoxide) amine core and methylacrylate and ethylenediamine core were found to demonstrate antibacterial and antifungal activities comparatively higher or equipotent to the antibacterial and antifungal agents used conventionally (Tulu et al., 2009). The quaternary ammonium terminated poly(propyleneimine) (PPI) dendrimer were found to be antibacterial against gram negative and gram positive bacteria (Chen et al., 2000; Chen and Cooper, 2002). The mechanism of antibacterial activity of the polycationic PPI dendrimers was attributed to the displacement of the divalent cations from the outer membrane of the gram negative bacteria causing cell wall disruption and lysis. Though the amine-terminated PPI dendrimers show excellent transfection and antibacterial activity, the major limitation to their use is the cytotoxicity and hence surface modification was sought to reduce cytotoxicity (Dutta et al., 2008; Tziveleka et al., 2007; Yang et al., 2008). Considerable effort has been directed in the past towards the surface group modification of the amine-terminated PAMAM dendrimers for reduce the cytotoxicity (Jevprasesphant et al., 2003; Kim et al., 2008; Kolhatkar et al., 2007; Yang et al., 2008). Size of the dendrimer affects its ability to penetrate the bacterial cell and therefore its antibacterial activity. For example, G₃-PAMAM dendrimers were more effective antibacterial agents than G₅-PAMAM dendrimers (Lopez et al., 2009). Similarly, lower generation carbosilane dendrimers were found to be more effective than higher generation dendrimers (Ortega et al., 2008). Typically, dendrimers have been investigated as carriers for drugs and imaging agents, but these recent findings present the perspective of dendrimers as new antimicrobial agents (Svenson and Tomalia, 2005).

Efforts continue to identify the newer and better antimicrobial agents. Antibacterial activity of amine-terminated PAMAM dendrimers is well recognized (Calabretta et al., 2007; Lopez et al., 2009), to the best of our knowledge, the assessment of the antibacterial activity of carboxylic and hydroxyl terminated PAMAM dendrimers has not been previously reported. In the present study, we have evaluated mechanisms by which hydroxyl and carboxylic acid terminated PAMAM dendrimers affect the cell wall of *E. coli*. Interestingly, we found that the hydroxyl terminated PAMAM dendrimer exhibited antibacterial activity *in vitro*. We evaluated the ability of the hydroxyl terminated PAMAM dendrimer to treat the *E. coli* induced uterine infection *in vivo* in pregnant guinea pig. Since, the guinea pig model of chorioamnionitis is well established to demonstrate the injury caused to the fetus resulting from *E. coli* infections (Patrick et al., 2004), we used this model for evaluating the *in vivo* activity of G₄-PAMAM-OH.

Intrauterine infection is usually caused by microorganisms ascending from vaginal and affecting the fetus and amniotic fluid leading to chorioamnionitis, cerebral palsy, increased efficiency of HIV seroconversion, miscarriage, and spontaneous preterm birth (Chaim et al., 1997; Romero et al., 2003; Ugwumadu, 2007). Chorioamnionitis is known to cause fetal brain injury (Patrick et al., 2004) due to the generation of proinflammatory cytokines (Dickinson et al., 2009; Harnett et al., 2007). Antibacterial and antifungal agents are applied to vagina and cervix to treat intrauterine infections in the pregnant women (Chaim et al., 1997; Ugwumadu, 2007). In the present study we show that *E. coli* infection in pregnant guinea pig can be treated by topical vaginal and cervical application of G₄-PAMAM-OH dendrimer. This is the first report

using the guinea pig model of chorioamnionitis to induce *E. coli* infections and show the effective inhibition of bacterial growth by treatment with G₄-PAMAM-OH. Our results show that the cytokine levels in placenta of the G₄-PAMAM-OH treated animals were comparable to those in healthy animals and significantly less than infected animals. Although PAMAM dendrimers are the most extensively studied dendrimers the antimicrobial activity of unmodified G₄-PAMAM-OH and G_{3.5}-PAMAM-COOH has not been reported previously. Though G₄-PAMAM-NH₂ dendrimer shows strong antibacterial activity it shows high cytotoxicity to human cervical cell line and the antibacterial activity of G₄-PAMAM-OH dendrimer is notable since it is non-cytotoxic at higher concentrations. Our findings bring out that G₄-PAMAM-OH has a potential as antibacterial agent.

2. Experimental

2.1. Materials

The PAMAM dendrimers (generation 4, with end groups OH, NH₂ and generation 3.5 COOH 14.93% w/w in methanol) were purchased from Dendritech. 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT), 2-nitrophenyl-β-D-galactopyranoside (ONPG), osmium tetroxide N-phenyl-1-naphthylamine (NPN), glutaraldehyde and hexamethyldisilazane were purchased from Invitrogen. Nutrient broth and nutrient agar were purchased from BD Biosciences. Mouse TNFα, IL-6 and IL-1β ELISA kits were purchased from R&D Systems.

2.2. Preparation of bacteria

E. coli (ATCC 11775) isolated from human urine is the bacterial strain used in this study. Single colony on nutrient agar was used to inoculate 5 mL of nutrient broth at 37 °C overnight. A small volume (100 μL) of this growth was used to inoculate 20 mL of nutrient broth media at 37 °C for 6 h. The bacteria were resuspended at 10⁶ colony forming units (CFU)/mL for the experiments.

2.3. Bacterial growth inhibition assays

The inhibitory concentration (IC₅₀) of dendrimers was determined using the broth microdilution method (Lopez et al., 2009; Wiegand et al., 2008). Briefly, serial dilutions of dendrimers (0.76 μg/mL to 200 mg/mL) were prepared in PBS and combined 1:1 (v/v) with bacteria at 10⁶ CFU/mL in a 96-well polypropylene plate. After incubation at 37 °C for 18 h, the absorbance was measured at 650 nm using a microplate reader to assess the cell growth. The positive control wells contained PBS and nutrient broth medium inoculated with bacteria (5 × 10⁵ CFU/mL), and the negative control wells contained PBS and nutrient broth medium without bacteria. The IC₅₀ value was determined as the concentration of the dendrimers which inhibits 50% of microbial growth after 18–24 h incubation (Lopez et al., 2009; Wiegand et al., 2008). The % survival of the bacteria was determined on the basis of the positive control which was considered as 100%.

2.4. Morphology of the bacteria

Scanning electron microscopy (SEM) analyses were performed to investigate the morphological changes of the *E. coli* following 8 h treatment with dendrimers G₄-PAMAM-OH, G_{3.5}-PAMAM-COOH and G₄-PAMAM-NH₂ at concentrations arrived at from bacterial growth inhibition assays. *E. coli* were fixed with 2.5% glutaraldehyde for 2 h. The samples were washed thrice with deionised water, and postfixed with 1% osmium tetroxide for 2 h, followed by washing thrice with deionised water. The samples were dehydrated

with a series of ethanol solutions (50%, 70%, 90% and 100%) followed by 10 min in 100% hexamethyldisilazane (HMDS), and were air dried. The samples were coated with gold and observed under SEM (HITACHI S-2400 Scanning Electron Microscope) at 20 kV.

2.5. Cell integrity

The bacterial cell membrane integrity was examined by determination of the release of nuclear acids (RNA/DNA) material absorbing at 260 nm (Je and Kim, 2006b; Ibrahim et al., 1991). *E. coli* were harvested, washed and resuspended in 0.5% NaCl solution with an absorbance of 0.7 at 420 nm. 150 μ L of G₄-PAMAM-OH, G_{3.5}-PAMAM-COOH and G₄-PAMAM-NH₂ solutions at concentrations arrived at from bacterial growth inhibition assays were mixed with 150 μ L of bacterial suspension in a 96-well plate, control was carried out with 0.5% NaCl alone, and the release over time of nuclear materials was monitored with a UV spectrometer by measuring the absorbance at 260 nm.

2.6. Outer membrane (OM) permeabilization assay

The OM permeabilization of *E. coli* by dendrimers was evaluated using the hydrophobic NPN (1-N-phenyl-1-naphthylamine) fluorescent probe (Helander and Mattila-Sandholm, 2000; Je and Kim, 2006a). *E. coli* were harvested, washed, and resuspended in 0.5% NaCl solution with an absorbance of 1.0 at 420 nm. The concentration of dendrimers was the same as described in previous section. 100 μ L of *E. coli* and 50 μ L of dendrimers were mixed with 50 μ L of 40 μ M NPN in black fluorotiter plate, control was carried out with 0.5% NaCl alone. An increase in fluorescence due to partitioning of NPN into the OM was recorded as a function of time until no further increase in intensity was observed. Excitation and emission wavelengths were set at 350 and 420 nm, respectively.

2.7. Inner membrane (IM) permeabilization assay

IM permeabilization was determined by measuring the release of cytoplasmic β -galactosidase activity from *E. coli* into the culture medium using ONPG as the substrate (Je and Kim, 2006b). *E. coli* were harvested, washed and resuspended in 0.5% NaCl solution an absorbance of 1.2 at 420 nm, then resuspended in half volume of 0.5% NaCl solution. The concentration of dendrimers was the same as described in previous section. 100 μ L of *E. coli* and 100 μ L of dendrimers were mixed with 10 μ L of ONPG (30 mM) in a 96-well plate, control was carried out with 0.5% NaCl alone. The production of *O*-nitrophenol over time was determined by monitoring the change in absorbance at 420 nm using a spectrophotometer.

2.8. Cell culture

Human cervical epithelial End1/E6E7 cells were obtained from ATCC, maintained in DMEM/F-12K medium supplemented with 20% fetal bovine serum (FBS). Mouse microglial BV-2 cells were obtained from Children's Hospital of Michigan Cell Culture Facility and maintained in DMEM supplemented with 5% FBS. These cells were cultured at 37 °C with 5% CO₂ and the media was replaced at 2 days interval.

2.9. Evaluation of normal cell cytotoxicity

End1/E6E7 and BV-2 (passage 19) cells were seeded into a 96-well plate at 1.5×10^4 /well, and 5×10^3 /well, respectively. After 24 h, cells were exposed to various concentrations of dendrimers (10 ng/mL to 1 mg/mL) in serum free medium for 24 h. Controls were carried out with medium alone. Cytotoxic effect was deter-

mined using MTT assay. The proportion of viable cells in the treated group was compared to that of the control.

2.10. Flow cytometry analysis

E. coli (10^6 CFU) and BV-2 (10^6) cells were treated with 10 μ g/mL of FITC-labeled G₄-PAMAM-OH dendrimer. The FITC-labeled G₄-PAMAM-OH dendrimer was prepared as previously reported (Kolhe et al., 2006). The *E. coli* and cells were washed 3 times with PBS, resuspended in 1% formaldehyde, and analyzed using a flow cytometer (FACS caliber, Becton Dickinson) by counting 20,000 and 10,000 events, respectively. The mean fluorescence intensity of cells was calculated using the histogram plot.

2.11. Confocal laser scanning microscopy

Microglial cells (10^6) were treated for 2 h with 10 μ g/mL of G₄-PAMAM-OH-FITC. The cells were washed with PBS 3 times and fixed with 4% para-formaldehyde for 20 min. Images were captured using a confocal microscope (Zeiss LSM 310) using a magnification of 400 \times . The excitation and emission wavelengths were 488 nm and 518 nm respectively, for FITC.

2.12. Evaluation the antimicrobial activity in guinea pig model of chorioamnionitis

All the animal experimental procedures were approved by the institutional animal care and use committee of Wayne State University. Intracervical bacterial inoculation was performed as previously reported (Patrick et al., 2004). Briefly, pregnant Dunkin–Hartley strain guinea pigs (Charles River) at 52 days of gestation were anesthetized with 1.5% isoflurane using the mask. An endoscope was used to visualize the cervix. Guinea pigs were inoculated intra-cervically with 150 CFU *E. coli* ($n = 11$) to induce infection. Dendrimer G₄-PAMAM-OH 500 μ g was injected into the cervix 5 min after *E. coli* inoculation in the treatment group 3 ($n = 4$). The *E. coli* inoculated guinea pigs without treatment (group 2) were used as positive control ($n = 4$). The guinea pigs without any treatment (group 1) and inoculation were used as negative controls ($n = 3$). Forty-eight hours after intervention, guinea pigs were euthanized with pentobarbital sodium (120 mg/kg) and mid-line laparotomy was performed to expose uterus. Amniotic fluid was collected from each gestational sac and 50 μ L was plated on nutrient agar to determine the presence of microbiologic chorioamnionitis.

2.13. Cytokine quantification in placenta

The placental tissue (0.3 g) was homogenized in 1 mL RIPA lysis buffer. The homogenate was kept on ice for 30 min, centrifuged at $10,000 \times g$ for 25 min at 4 °C and the protein concentration of supernatant was determined. Cytokines; tumor necrosis factor (TNF α), interleukin (IL-6 and IL-1 β) concentrations were measured in the total protein fraction using ELISA kits (Ethier-Chiasson et al., 2008).

2.14. Statistical analysis

Data are presented as mean \pm SD. Specific comparisons between control and individual experiment were analyzed by ANOVA test with p -value less than 0.05 considered as statistically significant.




Fig. 1. Bacterial growth inhibition assays. *E. coli* was treated with the indicated concentration of G₄-PAMAM-NH₂ (A) and (B), G₄-PAMAM-OH (C) and (D), G_{3.5}-PAMAM-COOH (E) and (F) dendrimers for 18 h. The initial concentration used for bacterial seeding was 5×10^5 CFU/mL. Three samples were in each group. Bacterial growth was measured by turbidity as the optical density at 650 nm using a microplate reader. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. positive control.

3. Results

3.1. Antimicrobial assay

We used the antibacterial assay procedure reported previously (Lopez et al., 2009) to assess the antimicrobial activity of G₄-PAMAM-OH and G_{3.5}-PAMAM-COOH dendrimers towards the gram negative bacteria *E. coli* and compared it with the activity of G₄-PAMAM-NH₂. We used *E. coli* in this study since it is known to cause the chorioamnionitis in pregnancy, in an established guinea pig model based on *E. coli* infection (Patrick et al., 2004). We have used *E. coli* for *in vitro* and *in vivo* evaluations to demonstrate the antibacterial activity of PAMAM dendrimers. In the present study we have measured the IC₅₀ values of PAMAM dendrimers using a modified broth microdilution assay in a 96-well plate format. The optical density of the suspension of bacteria in different dendrimer solutions was measured at 650 nm. The IC₅₀ value of the dendrimer was then obtained from the plot of % survival of bacteria vs. the concentrations of the dendrimer and the plot of optical densities vs. the concentrations of the dendrimer. G₄-PAMAM-OH, G_{3.5}-PAMAM-COOH and G₄-PAMAM-NH₂ dendrimers inhibited the growth of *E. coli* in a concentration-dependent manner as seen from 18 h treatment (Fig. 1). The strong antimicrobial activity of G₄-PAMAM-NH₂ is consistent with that reported previously (Calabretta et al., 2007). It is interesting to note that G₄-PAMAM-OH markedly inhibited the growth of *E. coli* from 3.13 mg/mL to 25.0 mg/mL concentration. G_{3.5}-PAMAM-COOH also inhibited the growth of *E. coli* but at relatively higher concentrations 6.25–100 mg/mL. For our set of conditions we observed the IC₅₀ values for G₄-PAMAM-OH, G_{3.5}-PAMAM-COOH and G₄-PAMAM-

NH₂ as 5.4 mg/mL, 22.0 mg/mL and 3.8 μg/mL respectively. Since G₄-PAMAM-NH₂ dendrimer exhibits high cytotoxicity, the G₄-PAMAM-OH was considered for *in vivo* evaluations in guinea pigs.

For the amine-terminated PAMAM dendrimers, it is proposed that the amino groups form nanoscale holes in supported lipid bilayers of bacterial membrane causing its rupture and cell lysis (Calabretta et al., 2007; Hong et al., 2006; Mecke et al., 2005; Milovic et al., 2005). The quaternary ammonium dendrimers adsorb onto negatively charged bacterial cell surfaces, diffuse through the cell wall, bind to cytoplasmic membrane, disrupt and disintegrate the cytoplasmic membrane, release of electrolytes such as potassium ions and phosphate from the cell and release nucleic materials such as DNA and RNA, all contributing to the death of the bacterial cell (Chen et al., 2000). These reports suggest that dendrimers mediate their antimicrobial activity by disrupting the bacterial outer and inner membrane. The antibiotic ampicillin is known to penetrate the outer membrane of gram negative bacteria and inhibits the bacterial cell wall synthesis. The antibacterial activity of dendrimers is limited to its effect on bacterial membrane permeabilities. However, there has not been any experimental data to support this suggestion and we performed the outer and inner membrane permeabilization assays to gain insights into the bactericidal mechanism of action of the PAMAM dendrimers. This is discussed in the subsequent sections in the manuscript.

3.2. SEM analysis of the *E. coli*

SEM studies were performed to examine if dendrimers damage the cell wall of *E. coli*. The concentration representing the IC₅₀ values of dendrimers was used. The SEM images collected




Fig. 2. SEM images of *E. coli*. (A) Untreated *E. coli*, (B) 8 h treatment of $G_{3.5}$ -PAMAM-COOH, (C) 8 h treatment of G_4 -PAMAM-OH and (D) 8 h treatment of G_4 -PAMAM-NH₂. Magnification 20,000 \times . Scale bars indicate 5 μ m. The treatment with dendrimers shows the damage to the bacterial cell wall.

after 8 h of treatment with dendrimers showed a marked difference in morphology (Fig. 2). The untreated *E. coli* (Fig. 2A) shows distinct rod shape with the intact cell wall. The treatment of *E. coli* with $G_{3.5}$ -PAMAM-COOH (Fig. 2B), G_4 -PAMAM-OH (Fig. 2C) and G_4 -PAMAM-NH₂ (Fig. 2D) shows shrinking, significant disruption of cell walls, and erosion of cell membrane. The major morphological changes observed by SEM support the antibacterial activity obtained in previous section. These studies indicated that major changes in the bacterial cell wall were induced by the dendrimers at the IC₅₀ concentrations, indicative of their antibacterial potential. In order to understand the underlying mechanism for the possible damage to the bacterial cell wall, the cell integrity, outer and inner membrane permeabilization studies were conducted.

3.3. Cell integrity

The cytoplasmic bacterial cell membrane undoubtedly is the target for many antimicrobial agents. The release of intracellular components is a good indicator of the compromise in cytoplasmic membrane integrity of gram negative bacteria resulting from the microbicidal activity of the tested compounds. In the present study, *E. coli* suspensions were treated with G_4 -PAMAM-OH, $G_{3.5}$ -PAMAM-COOH and G_4 -PAMAM-NH₂ dendrimers at the IC₅₀ concentrations and the absorbance of the suspensions was recorded at 260 nm. The absorbance significantly increased up to 20 min for each dendrimer at the tested concentration and thereafter the absorbance was almost unchanged (Fig. 3). The increase in the absorbance at 260 nm is due to the leaching of the small ions such as potassium and phosphate, but largely it's attributed to the

large nucleotides; DNA and the RNA released from the compromised bacterial cell. Similar observation was reported previously (Chen and Cooper, 2002) for QAC-functionalized dendrimers which exhibited fast killing kinetics till 10 min for *E. coli*. The increased absorbance is an indicator that at tested concentrations the three dendrimers were capable of altering the cytoplasmic membrane (outer cell membrane and cell wall) integrity for the gram negative bacteria.

3.4. Outer membrane (OM) permeabilization

The outer membrane in the gram negative bacteria acts as a strong barrier to foreign materials such as drugs. When the bacterial outer membrane is intact, it is impervious to the hydrophobic probe NPN. However once the outer membrane is compromised the NPN partitions in the perturbed outer membrane. Hence NPN is used to determine the changes induced by the biocides in the outer cell membrane. Typically, the greater the damage induced on the membrane, higher is the fluorescence reading, meaning greater is the NPN partitioning in membrane (Je and Kim, 2006b). The *E. coli* was treated with solutions of dendrimers as described in previous section. The G_4 -PAMAM-NH₂ dendrimer with amine groups on surface showed the highest fluorescence indicative of the rapid and bulk changes induced in the outer membrane of *E. coli* (Fig. 4). The divalent ions; calcium and magnesium play a key role in the stabilization of the bacterial membrane and are electrostatically bound to the anionic lipopolysaccharides (Chen and Cooper, 2002). The displacement of these divalent cations from the phospholipid membrane structure results in the structural instability, causing the disruption and eventually leading to cell death.




Fig. 3. Release of intracellular components of *E. coli* suspensions treated with (A) $G_{3.5}$ -PAMAM-COOH, (B) G_4 -PAMAM-OH and (C) G_4 -PAMAM-NH₂. Four samples were evaluated in each group. The increase in the absorbance is an indicator of the compromised cell integrity resulting in leaching of the nuclear components which are absorbed at 260 nm.




Fig. 4. Uptake of NPN by *E. coli* suspensions treated with (A) $G_{3.5}$ -PAMAM-COOH, (B) G_4 -PAMAM-OH and (C) G_4 -PAMAM-NH₂. Four samples were in each group.

At physiological pH, all of the primary amines and half of the tertiary amines in the dendrimer core of G_4 -PAMAM-NH₂ are protonated (Cakara and Borkovec, 2007; Cakara et al., 2003). The G_4 -PAMAM-NH₂ dendrimer is a polycation and it possibly replaces the divalent ions binding to the polyanions present in the cell membrane. This could lead to the insoluble polyanion-polycation complex and efflux of the calcium ions both contributing to major structural changes in outer cell membrane resulting in bacterial cell death.

It is interesting to note that the hydroxyl and carboxylic acid terminated dendrimers also showed an increase in the fluorescent intensity suggesting the NPN partitioning in treated *E. coli* (Fig. 4). These results clearly indicate that both G_4 -PAMAM-OH and $G_{3.5}$ -PAMAM-COOH alter the outer membrane structure of the *E. coli*. The underlying mechanism by which these dendrimers induce changes in the OM needs to be investigated. Several approaches can be considered to explain how the hydroxyl terminated dendrimer might act on the OM of bacteria. The microscopic protonation mechanism showed that at pH 7.4 and 7.5 the tertiary amines in the core of PAMAM dendrimers are protonated (Cakara and Borkovec, 2007; Cakara et al., 2003). The number of tertiary amines in the core is abundant when compared to the number of surface groups. Hence at pH 7.4, the core tertiary amines of G_4 -PAMAM-OH dendrimer are protonated and the surface hydroxyls are expected to bear negative charge and behave as anions. Anions are known to be bactericidal (Ellison et al., 1988; Khan et al., 2000). This could be one of the possible mechanisms by which G_4 -PAMAM-OH dendrimer affects the *E. coli* cell wall permeability. Further, the outer membrane of gram negative bacteria is extremely hydrophilic (Benz, 1988). The permeability of gram negative bacterial cell is altered by interaction of its O-antigens (outer polysaccharide) with hydroxyl groups and surface bound water (Jucker et al., 1997, 1998). The lipopolysaccharide of *E. coli* forms with hydroxyls and amines in the chitosan forming a complex which contributes to bactericidal effect of chitosan (Nabereznykh et al., 2008). The lipopolysaccharide from *E. coli* forms ionic, electrostatic and hydrogen bonds with chitosan and the hydroxyl groups of chitosan act as proton donors contributing towards these interactions. Further, the antibacterial activity due to hydroxyl groups is well established (Aslama et al., 2009; Myers, 1928). The literature clearly reveals the ability of the outer membrane of gram negative bacteria to form hydrogen bonds with hydroxyl groups. Our SEM images and OM permeability assay measurements together show the changes brought out in the bacterial cell membrane due to the activity of G_4 -PAMAM-OH dendrimer. Based on experimental and the inferential evidence, one can interpret that G_4 -PAMAM-OH dendrimer forms hydrogen bonds with the outer membrane of *E. coli*, another possible mechanism by which it could bring the permeability changes. The in-depth understanding on the mechanism by which G_4 -PAMAM-OH works is required and it is beyond the scope of this paper.

At physiological pH, the carboxylic acid terminated dendrimer acts as polyanion and chelates the positive calcium and magnesium ions in the outer membrane altering its structure and permeability. The phosphate group in the phosphatidyl choline (lipid) layers of bacteria has free carboxyl groups and phosphate groups bearing a negative charge which bind to cations. The outer membrane of gram negative bacteria is stabilized by the ionic interaction and binding of the positive ions (calcium and magnesium) to the polar anionic residues on the lipid head groups. This complex is reversible. Since this complex is reversible, the carboxylic groups from dendrimer replace the carboxylic groups from the lipids and bind to positive calcium ions destabilizing the membrane. This could be a possible mechanism by which the $G_{3.5}$ -PAMAM-COOH alters the outer membrane of the *E. coli*. There are reports which support this assumption. The ethylenediamine tetraacetic acid is antibac-

terial as it binds to the membrane stabilizing cations (Ellison et al., 1988). Incorporation of palmitic acid residues in lysozyme rendered it bactericidal to *E. coli* as hydrophobic interactions occur between the two in LPS zone (Ibrahim et al., 1991). The anionic phosphatidyl shows antibacterial activity towards *E. coli* by chelating calcium ions in outer membrane (Khan et al., 2000). Further, anionic peptide MDpep5 acts as antimicrobial by binding to the bacteria through hydrophobic interactions causing disorders in bilayer structure. The membrane disruption is not complete for anionic MDpep5 when compared to the cationic (AMPs) antimicrobial peptide (Tang et al., 2008). Based on the literature, when we review the results of present study, it appears that at physiological pH the carboxylate ions on the G_{3.5}-PAMAM-COOH act as polyanions interacting with the calcium ions in the outer membrane of gram negative bacteria affecting its permeability. As compared to the anionic G_{3.5}-PAMAM-COOH, the cationic G₄-PAMAM-NH₂ causes more damage to outer membrane structure, which is again consistent with the observations reported by Tang et al. (2008). From our studies we found that PAMAM dendrimers with amine, hydroxyl and carboxyl terminal group, could each show a different path for interacting with the bacterial outer membrane.

3.5. IM permeabilization

Destabilization of the outer membrane in bacteria is essential for establishing contact with the inner membrane (Benz, 1988; Je and Kim, 2006b). The inner layer of bacteria consists of phosphatidyl glycerol and cardiolipin (Je and Kim, 2006a). The cytoplasmic β -galactosidase is released as a consequence of alteration/compromise in inner membrane and it reacts with ONPG to yield ortho-nitrophenol which is absorbed at 420 nm. The concentrations of three dendrimers were the same as IC₅₀ values. There was a lag of 20 min followed by which a slow release of cytoplasmic β -galactosidase was observed for all the dendrimers tested. The release of cytoplasmic β -galactosidase was found to be time dependant (Fig. 5). Amongst the dendrimers evaluated, the G₄-PAMAM-NH₂ showed a marked increase in OD values indicative its capability to induce major permeability changes in inner membrane. The G_{3.5}-PAMAM-COOH dendrimer treated bacteria showed increase in OD with respect to time, again indicating the changes brought in the structure of inner wall. The OD values were low for G₄-PAMAM-OH dendrimer as compared to the other dendrimers but they increased slowly over time as seen from Fig. 5. These results indicate that the G₄-PAMAM-OH dendrimer seems to induce profound permeability changes in outer membrane as compared to the inner membrane. While the G_{3.5}-PAMAM-COOH dendrimer appears to release the cell intracellular nucleotides more than the other dendrimers, the cationic G₄-PAMAM-NH₂ dendrimer appears to disrupt the IM and OM to a significantly higher extent compared to the other dendrimers.

3.6. Cytotoxicity assay

The cytotoxicity of PAMAM dendrimers was evaluated against human cervical epithelial (End1/E6E7) and immune cells; mouse microglial cells (BV-2). MTT assay showed that G₄-PAMAM-OH and G_{3.5}-PAMAM-COOH dendrimers were non-cytotoxic to End1/E6E7 cells and BV-2 cells in 24 h treatment at concentrations 10 ng/mL to 1 mg/mL (Fig. 6). The G₄-PAMAM-NH₂ showed high cytotoxicity above 10 μ g/mL concentration to human cervical epithelial End1/E6E7 cells. Also the G₄-PAMAM-NH₂ exhibited cytotoxicity at 1 mg/mL concentration to microglial cells. On the basis of the MTT assay, G₄-PAMAM-OH did not exhibit cytotoxicity up to 1 mg/mL concentration, while the G₄-PAMAM-NH₂ was found to be cytotoxic at higher concentrations. *E. coli* infections in the vagina can lead to preterm birth. The experimental data shows that G₄-




Fig. 5. Release of cytoplasmic β -galactosidase of *E. coli* treated with (A) G_{3.5}-PAMAM-COOH, (B) G₄-PAMAM-OH and (C) G₄-PAMAM-NH₂. Four samples were in each group.

PAMAM-OH dendrimer is non-cytotoxic to the human cervical cell line and also exhibits antibacterial activity towards *E. coli*, hence it was chosen as antibacterial agent to prevent preterm birth in pregnant guinea pigs. In our *in vivo* experiments we applied a total of 500 μ g of G₄-PAMAM-OH to the cervix of *E. coli* infected pregnant guinea pigs and at this concentration the dendrimer showed efficacy.




Fig. 6. Cytotoxicity assay. (A) Human cervical epithelial End1/E6E7 cells and (B) mouse microglial cells were treated with the G₄-PAMAM-OH, G_{3.5}-PAMAM-COOH and G₄-PAMAM-NH₂ dendrimers at concentrations indicated for IC₅₀ values. Three samples were in each group. Cell viability was assessed by MTT method. The proportion of viable cells in the treated group was compared to that of negative control.




Fig. 7. Flow cytometry of the cell entry dynamics of (A) G₄-PAMAM-OH-FITC in *E. coli* and (B) BV-2 microglial cell line. The log of FITC absorption intensity (FL1-H on x-axis) is plotted against the number of cells (counts on y-axis). The maximum uptake of G₄-PAMAM-OH-FITC in *E. coli* occurs at 3 h. The rapid cellular uptake of G₄-PAMAM-OH-FITC within 15 min in microglial cells is evident. The transport of conjugate into microglial cell increased with increasing time. Confocal microscopy images (400×) showed that G₄-PAMAM-OH-FITC appeared to be mainly localized in the cytoplasm of BV-2 cells while the nucleus appeared to be relatively free of the presence of any fluorescence at this time point (C).

3.7. Cell entry dynamics of dendrimer in *E. coli* and microglial cells

The size of the dendrimer affects its antibacterial activity and G₅-PAMAM-NH₂ was less bactericidal than G₃-PAMAM-NH₂ which could penetrate the bacterial cell because of relatively smaller size (Lopez et al., 2009). In the present study, we have evaluated the generation-4 PAMAM dendrimers and hence the ability of G₄-PAMAM-OH to enter the cells was determined using a FITC tagged G₄-PAMAM-OH-FITC conjugate. The *E. coli* did not show significant increase in fluorescence intensity after treatment with G₄-PAMAM-OH-FITC for 2 h. However, a significant increase in fluorescence intensity was observed after 3 h (Fig. 7A) confirming a slow entry of G₄-PAMAM-OH-FITC conjugate into the *E. coli*. This demonstrates that G₄-PAMAM-OH enters the *E. coli* effectively while the exact location of G₄-PAMAM-OH in the bacteria remains unclear and needs to be further elucidated using other methods. These results support the previous studies where the G₄-PAMAM-OH shows changes in the permeability of the bacterial membrane. The microglial cells showed a significant fluorescence intensity increase within 15 min, as evidenced by a two-order of magnitude increase in the fluorescence intensity. This was followed by a moderate increase in fluorescence over the next 4 h (Fig. 7B) indicating that G₄-PAMAM-OH-FITC conjugate entered the microglial cells. In case of BV-2 cells the cellular entry was visualized by using confocal microscopy showing the localization of G₄-PAMAM-OH-FITC in cytoplasm (Fig. 7C). The uptake of the dendrimer by *E. coli* was slower than that in microglial cells. The difference in the structure

of cell membrane may result in different cell entry dynamics of dendrimer in *E. coli* and microglial cells.

3.8. Antimicrobial activity in guinea pig model of chorioamnionitis

The *in vitro* studies brought out antibacterial potential of G₄-PAMAM-OH dendrimer as seen from the antibacterial assay, OM and IM permeabilization assays and bulk changes in morphology seen from SEM analysis. These interesting results coupled with its non-cytotoxicity to human cervical epithelial cells encouraged the evaluation of G₄-PAMAM-OH as an antibacterial agent *in vivo* using the guinea pig model of chorioamnionitis. Though this model is established for creating infection and assessing injury to the fetus, it has not been previously used to demonstrate the effective treatment. This is the first study to our knowledge, which shows the treatment of the pregnant guinea pig using the model of chorioamnionitis. The ascending *E. coli* infection causes chorioamnionitis which is associated with development of cerebral palsy, a motor disorder in children due to stimulation of proinflammatory cytokines causing white matter damage and fetal brain injury (Patrick et al., 2004). We optimized the dose of the *E. coli* inoculation in the guinea pigs ($n=17$) in the pilot experiments for the strain (ATCC 11775). We observed that 1000 CFU of *E. coli* effectively induced the infection causing extreme sickness in mother, leading to abortion of dead fetuses within 48–72 h. The lower CFU of *E. coli* were subsequently inoculated to identify the optimum dose which lead to infection and yet the guinea

Table 1The inhibition of *E. coli* growth after treatment with G₄-PAMAM-OH dendrimer in guinea pig model of chorioamnionitis.

Inoculation with <i>E. coli</i>			Treatment: with G ₄ -OH after <i>E. coli</i> inoculation		
Guinea Pig	Amniotic fluid from different gestational sacs		Guinea Pig	Amniotic fluid from different gestational sacs	
	Bacterial growth	%		Bacterial growth	%
Mother 1	Fetuses (4/5)	80.0	Mother 1	Fetuses (0/5)	0
Mother 2	Fetuses (5/6)	83.3	Mother 2	Fetuses (0/6)	0
Mother 3	Fetuses (1/4)	25.0	Mother 3	Fetuses (0/5)	0
Mother 4	Fetuses (2/5)	40.0	Mother 4	Fetuses (0/3)	0
	Average	57.1		Average	0

pigs did not abort up to 48 h. Based on this evaluation a dose of 150 CFU of *E. coli* was found to effectively induce the infection in the pregnant guinea pigs without leading to abortion of fetuses.

In the present study, chorioamnionitis was induced after intracervical inoculation with *E. coli* in 8 guinea pigs of which $n=4$ were considered as positive control (group 2) and $n=4$ were used for treatment with G₄-PAMAM-OH (group 3). None of the amniotic fluid samples plated from the negative control group 1 ($n=3$) showed evidence of microbiologic chorioamnionitis. Of the pregnant guinea pigs (group 2, $n=4$) that were inoculated with 150 CFU of *E. coli*, 57.1% of the amniotic fluid samples for different fetus were positive with bacterial growth (indicative of induced infection) as seen from the culture inoculated with it (see Table 1). Prenatal exposure to maternal infection alters cytokine expression in the placenta (Urakuboa et al., 2001). Abundance of cytokines in placental tissues is an indicator of activation of inflammatory response in gestational membranes with term and preterm parturition (Keelan et al., 1999). The cytokine IL-6 is known to peak after 48 h of infection (Dickinson et al., 2009) and hence in present study animals were sacrificed after 48 h to determine the cytokine level in positive and treated animals. When we compared the expression of cytokine levels in negative control vs. the positive controls, the cytokines, especially TNF α and IL-6, increased significantly in the placenta of positive controls after 48 h of inoculation with *E. coli* (Fig. 8). These results demonstrated that chorioamnionitis was successfully induced after 48 h of cervical inoculation with 150 CFU of *E. coli*.

The G₄-PAMAM-OH dendrimer was applied topically at a dose of (625 μ g/kg) on the cervical endometrium of guinea pigs (group 3, $n=4$) in form of aqueous solution in saline after *E. coli* inoculation. The total amount applied was 500 μ g dissolved in saline based on the average weight of the guinea pigs (800 g/animal). The amniotic fluid samples for different fetus were collected after 48 h and were plated on the culture plates and evaluated for the bacterial growth. All these samples did not show any bacterial growth (0%) on the culture plates (Table 1). The study shows that the treatment with G₄-PAMAM-OH dendrimer completely eliminated the bacterial growth and prevented bacteria ascension into uterine cavity and amniotic fluid i.e. from 57.1 (positive control) to 0% (treatment group) bacterial growth. Earlier, the *in vitro* data showed antibacterial nature of G₄-PAMAM-OH at higher concentration and the *in vivo* results show that amniotic fluid samples for different fetus in treatment group 3 were found to be negative. The comparison of cytokine expression in placenta of the treatment group, negative and positive control groups shows that the cytokine levels (TNF α and IL-6) in treatment group are comparable to the negative control while they are overly expressed in positive controls (Fig. 8). These results indicate the potential of G₄-PAMAM-OH to effectively kill gram negative bacteria *E. coli* in cervix of guinea pig and prevent chorioamnionitis. This is a significant finding since the chorioamnionitis is known to cause fetal brain injury (Patrick et al., 2004) which could possibly be averted by treatment with G₄-




Fig. 8. The placental tissue (0.3 g) was homogenized in 1 ml RIPA lysis buffer. The homogenate was kept on ice for 30 min and the protein concentration of supernatant was determined. Cytokines concentrations were measured in the total protein fraction using ELISA. (A) TNF α measurements in normal, *E. coli* infected and G₄-PAMAM-OH treated guinea pigs (B) IL-6 measurements in normal, *E. coli* infected and G₄-PAMAM-OH treated guinea pigs (C) IL-1 β measurements in normal, *E. coli* infected and G₄-PAMAM-OH treated guinea pigs * $P<0.05$, *** $P<0.001$ vs. Normal control. ▲ $P<0.01$, ▲▲ $P<0.001$ vs. *E. coli* group.

PAMAM-OH as indicated from these findings. It is interesting to note that the G₄-PAMAM dendrimers show very negligible transport across the fetal membranes and can therefore be possibly used for selective treatment of mother without crossing into the fetus when administered by intravaginal route (Menjoge et al., 2010).

4. Conclusion

The bactericidal activity of hydroxyl and carboxylic acid terminated PAMAM dendrimer was evaluated against gram negative *E. coli* and compared with amine-terminated PAMAM dendrimers. The antimicrobial assay, SEM analysis, cell integrity, inner and outer membrane permeability assays showed that G₄-PAMAM-OH and G_{3.5}-PAMAM-COOH dendrimers affect the cell wall of *E.*

coli, and were antibacterial at the concentrations evaluated. The major finding was the bactericidal effect of G₄-PAMAM-OH dendrimer and its ability to treat *E. coli* infections *in vivo* in pregnant guinea pigs. Topical cervical application of 500 µg of G₄-PAMAM-OH treated the *E. coli* infections induced in guinea pig model of chorioamnionitis. The amniotic fluid collected from different fetus in the infected guinea pigs, posttreatment showed absence of *E. coli* growth in the cultures plated with it. The cytokines levels were higher in the positive controls confirming presence of infection after inoculation with *E. coli*. The cytokine expression (TNFα and IL-6) in the treatment group was comparable to that in negative control showing the efficacy of G₄-PAMAM-OH to treat the *E. coli* infections. The G₄-PAMAM-NH₂ dendrimer is known to be potent antibacterial agent, however, it was found to be highly cytotoxic to above 10 µg/mL to human cervical epithelial (End1/E6E7) cells and immune cells (BV-2) while the G₄-PAMAM-OH dendrimer was non-cytotoxic up to 1 mg/mL concentrations to both cell lines. Each dendrimer appears to affect the bacterial cell wall in a different way. The possible mechanisms involve the G₄-PAMAM-NH₂ acting as polycation binding to the polyanionic lipopolysaccharide, the G₄-PAMAM-OH binding via hydrogen bonds to the hydrophilic O-antigens and the G_{3,5}-PAMAM-COOH acting as a polyanion chelating the divalent ions in outer cell membrane. The outer and inner membrane permeabilization assay shows that G₄-PAMAM-OH brings major structural changes to the outer membrane whereas G₄-PAMAM-NH₂ brings major changes to both outer and inner membrane.

Acknowledgments

This study was supported by the Intramural Research Program of the National Institute of Child Health and Human Development, NIH, DHHS, and the Pediatric Critical Care Scientist Development Program NICHD-K08.

References

- Aslama, S.N.S., Kokubun, T., Halla, D.R., 2009. Antibacterial and antifungal activity of cicerfuran and related 2-arylbenzofurans and stilbenes. *Microbiol. Res.* 164, 191–195.
- Balogh, L., Swanson, D.R., Tomalia, D.A., Hagnauer, G.L., McManus, A.T., 2001. Dendrimer–silver complexes and nanocomposites as antimicrobial agents. *Nano Lett.* 1, 18–21.
- Benz, R., 1988. Structure and function of porins from gram negative bacteria. *Microbiol. Lett.* 42, 359–393.
- Bourne, N., Stanberry, L.R., Kern, E.R., Holan, G., Matthews, B., Bernstein, D.I., 2000. Dendrimers, a new class of candidate topical microbicides with activity against herpes simplex virus infection. *Antimicrob. Agents Chemotherapy* 44, 2471–2474.
- Cakara, D., Borkovec, 2007. Microscopic M, protonation mechanism of branched polyamines: poly(amidoamine) versus poly(propyleneimine) dendrimers. *Croat. Chem. Acta* 80, 421–428.
- Cakara, D., Kleimann, J., Borkovec, M., 2003. Protonation equilibria of poly(amidoamine) dendrimers from macroscopic titrations. *Macromolecules* 36, 4201–4207.
- Calabretta, M.K., Kumar, A., McDermott, A.M., Cai, C., 2007. Antibacterial activities of poly(amidoamine) dendrimers terminated with amino and poly(ethylene glycol) groups. *Biomacromolecules* 8, 1807–1811.
- Chaim, W., Mazor, M., Leiberman, J.R., 1997. The relationship between bacterial vaginosis and preterm birth. A review. *Arch. Gynecol. Obstet.* 259, 51–58.
- Chen, C.Z., Beck-Tan, N.C., Dhurjati, P., van Dyk, T.K., LaRossa, R.A., Cooper, S.L., 2000. Quaternary ammonium functionalized poly(propylene imine) dendrimers as effective antimicrobials: structure–activity studies. *Biomacromolecules* 1, 473–480.
- Chen, C.Z., Cooper, S.L., 2002. Interactions between dendrimer biocides and bacterial membranes. *Biomaterials* 23, 3359–3368.
- Cheng, Y., Qu, H., Ma, M., Xu, Z., Xu, P., Fang, Y., Xu, T., 2007. Polyamidoamine (PAMAM) dendrimers as biocompatible carriers of quinolone antimicrobials: an *in vitro* study. *Eur. J. Med. Chem.* 42, 1032–1038.
- Cloninger, M.J., 2002. Biological applications of dendrimers. *Curr. Opin. Chem. Biol.* 6, 742–748.
- Dickinson, M.A., Harnett, E.L., Venditti, C.C., Smith, G.N., 2009. Transient lipopolysaccharide-induced cytokine responses in the maternal serum and amniotic fluid of the guinea pig. *Am. J. Obstet. Gynecol.* 200, pp. 534 e531–536.
- Dutta, T., Garg, M., Jain, N.K., 2008. Poly(propyleneimine) dendrimer and dendrosome mediated genetic immunization against hepatitis B. *Vaccine* 26, 3389–3394.
- Ellison 3rd, R.T., Giehl, T.J., LaForce, F.M., 1988. Damage of the outer membrane of enteric gram-negative bacteria by lactoferrin and transferrin. *Infect. Immun.* 56, 2774–2781.
- Ethier-Chiasson, M., Forest, J.-C., Giguere, Y., Masse, A., Marseille-Tremblay, C., Levy, E., Lafond, J., 2008. Modulation of placental protein expression of OLR1: implication in pregnancy-related disorders or pathologies. *Reproduction* 136, 491–502.
- Halford, B., 2005. Dendrimers branch out. *C. EN* 83, 30–36.
- Harnett, E.L., Dickinson, M.A., Smith, G.N., 2007. Dose-dependent lipopolysaccharide-induced fetal brain injury in the guinea pig. *Am. J. Obstet. Gynecol.* 197, pp. 179 e171–177.
- Helander, I.M., Mattila-Sandholm, T., 2000. Fluorometric assessment of gram-negative bacterial permeabilization. *J. Appl. Microbiol.* 88, 213–219.
- Hong, S., Leroueil, P.R., Janus, E.K., Peters, J.L., Kober, M.M., Islam, M.T., Orr, B.G., Baker Jr., J.R., Banaszak Holl, M.M., 2006. Interaction of polycationic polymers with supported lipid bilayers and cells: nanoscale hole formation and enhanced membrane permeability. *Bioconjug. Chem.* 17, 728–734.
- Hou, S., Zhou, C., Liu, Z., Young, A.W., Shi, Z., Ren, D., Kallenbach, N.R., 2009. Antimicrobial dendrimer active against *Escherichia coli* biofilms. *Bioorg. Med. Chem. Lett.* 19, 5478–5481.
- Ibrahim, H.R., Kato, A., Kobayashi, K., 1991. Antimicrobial effects of lysozyme against gram-negative bacteria due to covalent binding of palmitic acid. *J. Agric. Food Chem.* 39, 2077–2082.
- Je, J.Y., Kim, S.K., 2006a. Antimicrobial action of novel chitin derivative. *Biochim. Biophys. Acta* 1760, 104–109.
- Je, J.Y., Kim, S.K., 2006b. Chitosan derivatives killed bacteria by disrupting the outer and inner membrane. *J. Agric. Food Chem.* 54, 6629–6633.
- Jevprasesphant, R., Penny, J., Jalal, R., Attwood, D., McKeown, N.B., D'Emanuele, A., 2003. The influence of surface modification on the cytotoxicity of PAMAM dendrimers. *Int. J. Pharm.* 252, 263–266.
- Jucker, B.A., Harms, H., Hug, S.J., Zehnder, A.J.B., 1997. Adsorption of bacterial surface polysaccharides on mineral oxides is mediated by hydrogen bonds. *Colloids Surf. B: Biointerfaces* 9, 331–343.
- Jucker, B., Zehnder, A.B.D., Harms, H., 1998. Quantification of polymer interactions in bacterial adhesion. *Environ. Sci. Technol.* 32, 2909–2915.
- Keelan, J.A., Marvin, K.W., Sato, T.A., Coleman, M., McCowan, L.M., Mitchell, M.D., 1999. Cytokine abundance in placental tissues: evidence of inflammatory activation in gestational membranes with term and preterm parturition. *Am. J. Obstet. Gynecol.* 181, 1530–1536.
- Khan, M.A.S., Nakamura, S., Ogawa, M., Akita, E., Azakami, H., Kato, A., 2000. Bactericidal action of egg yolk phosphatidylcholine against *Escherichia coli* under thermal stress. *J. Agric. Food Chem.* 48, 1503–1506.
- Kim, Y., Klutz, A.M., Jacobson, K.A., 2008. Systematic investigation of polyamidoamine dendrimers surface-modified with poly(ethylene glycol) for drug delivery applications: synthesis, characterization, and evaluation of cytotoxicity. *Bioconjug. Chem.* 19, 1660–1672.
- Kolhatkar, R.B., Kitchens, K.M., Swaan, P.W., Ghandehari, H., 2007. Surface acetylation of polyamidoamine (PAMAM) dendrimers decreases cytotoxicity while maintaining membrane permeability. *Bioconjug. Chem.* 18, 2054–2060.
- Kolhe, P., Khandare, J., Pillai, O., Kannan, S., Lieh-Lai, M., Kannan, R.M., 2006. Preparation, cellular transport, and activity of polyamidoamine-based dendritic nanodevices with a high drug payload. *Biomaterials* 27, 660–669.
- Lebreton, S., Newcombe, N., Bradley, M., 2003. Antibacterial single-bead screening. *Tetrahedron* 59, 10213–10222.
- Lopez, A.I., Reins, R.Y., McDermott, A.M., Trautner, B.W., Cai, C., 2009. Antibacterial activity and cytotoxicity of PEGylated poly(amidoamine) dendrimers. *Mol. Biosyst.* 5, 1148–1156.
- Mecke, A., Majoros, I.J., Patri, A.K., Baker Jr., J.R., Holl, M.M., Orr, B.G., 2005. Lipid bilayer disruption by polycationic polymers: the roles of size and chemical functional group. *Langmuir* 21, 10348–10354.
- Menjoge, A.R., Navath, R.S., Asad, A., Kannan, S., Kim, C.J., Romero, R., Kannan, R.M., 2010. Transport and biodistribution of dendrimers across human fetal membranes: implications for intravaginal administration of dendrimer–drug conjugates. *Biomaterials* 31, 5007–5021.
- Milovic, N.M., Wang, J., Lewis, K., Klibanov, A.M., 2005. Immobilized N-alkylated polyethylenimine avidly kills bacteria by rupturing cell membranes with no resistance developed. *Biotechnol. Bioeng.* 90, 715–722.
- Mumper, R.J., Bell, M.A., Worthen, D.R., Cone, R.A., Lewis, G.R., Paull, J.R., Moench, T.R., 2009. Formulating a sulfonated antiviral dendrimer in a vaginal microbicidal gel having dual mechanisms of action. *Drug Develop. Ind. Pharm.* 35, 515–524.
- Myers, R.P., 1928. The effect of hydroxyl ion concentration on the thermal death rate of *Bacterium coli*. *J. Bacteriol.* 15, 341–356.
- Naberezhnykh, G.A., Gorbach, V.I., Likhatskaya, G.N., Davidova, V.N., Solov'eva, T.F., 2008. Interaction of chitosans and their N-acylated derivatives with lipopolysaccharide of gram-negative bacteria. *Biochemistry (Mosc.)* 73, 432–441.
- Ortega, P.C.-P.J.L., Munoz-Fernandez, M.A., Soliveri, J., Gomez, R., Mata, F., Jdela, 2008. Amine and ammonium functionalization of chloromethylsilane-ended dendrimers. *Antimicrobial activity studies*. *J. Biomol. Chem.* 6, 3264–3269.
- Patrick, L.A., Gaudet, L.M., Farley, A.E., Rossiter, J.P., Tomalty, L.L., Smith, G.N., 2004. Development of a guinea pig model of chorioamnionitis and fetal brain injury. *Am. J. Obstet. Gynecol.* 191, 1205–1211.
- Romero, R., Chaiworapongsa, T., Espinoza, J., 2003. Micronutrients and intrauterine infection, preterm birth and the fetal inflammatory response syndrome. *J. Nutr.* 166S–167S.

- Svenson, S., Tomalia, D.A., 2005. Dendrimers in biomedical applications—reflections on the field. *Advanced drug delivery reviews* 57, 2106–2129.
- Tang, Y.L., Shi, Y.H., Zhao, W., Hao, G., Le, G.W., 2008. Insertion mode of a novel anionic antimicrobial peptide MDpep5 (Val-Glu-Ser-Trp-Val) from Chinese traditional edible larvae of housefly and its effect on surface potential of bacterial membrane. *J. Pharm. Biomed. Anal.* 48, 1187–1194.
- Tulu, M., Aghatabay, N.M., Senel, M., Dizman, C., Parali, T., Dulger, B., 2009. Synthesis, characterization and antimicrobial activity of water soluble dendritic macromolecules. *Eur. J. Med. Chem.* 44, 1093–1099.
- Tziveleka, L.A., Psarra, A.M., Tsiourvas, D., Paleos, C.M., 2007. Synthesis and characterization of guanidinylated poly(propylene imine) dendrimers as gene transfection agents. *J. Control. Release* 117, 137–146.
- Ugwumadu, A., 2007. Role of antibiotic therapy for bacterial vaginosis and intermediate flora in pregnancy. *Best Pract. Res.* 21, 391–402.
- Urakuboa, A., Jarskog, L.F., Lieberman, J.A., Gilmore, J.H., 2001. Prenatal exposure to maternal infection alters cytokine expression in the placenta, amniotic fluid, and fetal brain. *Schizophr. Res.* 47, 27–36.
- Wiegand, I., Hilpert, K., Hancock, R.E., 2008. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat. Protoc.* 3, 163–175.
- Yang, H., Lopina, S.T., DiPersio, L.P., Schmidt, S.P., 2008. Stealth dendrimers for drug delivery: correlation between PEGylation, cytocompatibility, and drug payload. *J. Mater. Sci. Mater. Med.* 19, 1991–1997.