



Erratum

Erratum to “Evaluation of gastrointestinal motility directly from human pharmacokinetic data” [Int. J. Pharm. 419 (2011) 43–51]

John A. Roush *

GlaxoSmithKline, Five Moore Drive, Research Triangle Park, NC 27709, United States

The publisher regrets that in the above named article, the references were not numbered. Please see the list of numbers and references below.

- [1] Dokoumetzidis, A., Kalantzi, L., Fotaki, N., 2007. Predictive models for oral drug absorption: from *in silico* methods to integrated dynamic models. *Expert Opin. Metab. Toxicol.* 3, 491–505.
- [2] Goodacre, B.C., Murray, R.J., 1981. A mathematical model of drug absorption. *J. Clin. Hosp. Pharm.* 6, 117–133.
- [3] Suttle, B.A., Pollack, G.M., Brouwer, K.L.R., 1992. Use of a pharmacokinetic model incorporating discontinuous gastrointestinal absorption to examine the occurrence of double peaks in oral concentration-time profiles. *Pharm. Res.* 9, 350–356.
- [4] Yu, L.X., Lipka, E., Crison, J.R., Amidon, G.L., 1996. Transport approaches to the biopharmaceutical design of oral drug delivery systems: prediction of intestinal absorption. *Adv. Drug Deliv. Rev.* 19, 359–376.
- [5] Sawamoto, T., Haruta, S., Kurosaki, Y., 1997. Prediction of plasma concentration profiles of orally administered drugs in rats on the basis of gastrointestinal transit kinetics and absorbability. *J. Pharm. Pharmacol.* 49, 450–457.
- [6] Kimura, T., Higaki, K., 2002. Gastrointestinal transit and drug absorption. *Biol. Pharm. Bull.* 25, 149–164.
- [7] Dressman, J.B., Fleisher, D., Amidon, G.L., 1984. Physicochemical model for dosedependent drug absorption. *J. Pharm. Sci.* 73, 1274–1279.
- [8] Dressman, J.B., Fleisher, D., 1986. Mixing-tank model for predicting dissolution rate control of oral absorption. *J. Pharm. Sci.* 75, 109–116.
- [9] Hintz, R.J., Johnson, K.C., 1989. The effect of particle size distribution on dissolution rate and oral absorption. *Int. J. Pharm.* 51, 9–17.
- [10] Oberle, R.L., Amidon, G.L., 1987. The influence of variable gastric emptying and intestinal transit rates on plasma level curve of cimetidine: an explanation for the double peak phenomenon. *J. Pharmacokinet. Biopharm.* 15, 529–544.
- [11] Luner, P., Amidon, G.L., 1993. Description and simulation of a multiple mixing tank model to predict the effect of bile sequestrants on bile salt excretion. *J. Pharm. Sci.* 82, 311–318.
- [12] Cutler, D.J., 1978. Numerical deconvolution by least squares: use of prescribed input functions. *J. Pharmacokinet. Biopharm.* 6, 227–241.
- [13] Cutler, D.J., 1978. Numerical deconvolution by least squares: use of polynomials to represent the input function. *J. Pharmacokinet. Biopharm.* 6, 243–263.
- [14] Johnson, D.A., Amidon, G.L., 1988. Determination of intrinsic membrane transport parameters from perfused intestine experiments: a boundary layer approach to estimating the aqueous and unbiased membrane permeabilities. *J. Theor. Biol.* 131, 93–106.
- [15] Li, J., Cutler, D.J., 1998. Stability of finite difference deconvolution. I: theoretical analysis. *Biopharm. Drug Dispos.* 19, 547–554.

DOI of original article: [10.1016/j.ijpharm.2011.07.011](https://doi.org/10.1016/j.ijpharm.2011.07.011).

* Tel.: +1 919 483 8205; fax: +1 919 483 0443.

E-mail address: john.a.roush@gsk.com