



## Corrigendum

# Corrigendum to “Prestin binding peptides as ligands for targeted polymersome mediated drug delivery to outer hair cells in the inner ear” [Int. J. Pharm. 424 (2012) 121–127]

Elena V. Surovtseva<sup>a,\*</sup>, Alexander H. Johnston<sup>b</sup>, Weikai Zhang<sup>a</sup>, Ya Zhang<sup>a</sup>, Adele Kim<sup>a</sup>,  
Michio Murakoshi<sup>c</sup>, Hiroshi Wada<sup>c</sup>, Tracey A. Newman<sup>d</sup>, Jing Zou<sup>a</sup>, Ilmari Pyykkö<sup>a</sup>

<sup>a</sup> Department of Otolaryngology, University of Tampere, Tampere 33014, Finland

<sup>b</sup> Centre for Biological Sciences, Life Sciences Building 85, Highfield Campus, University of Southampton, SO17 1BJ, United Kingdom

<sup>c</sup> Department of Bioengineering and Robotics, Tohoku University, 6-6-01 Aoba-yama, Sendai 980-8579, Japan

<sup>d</sup> Clinical Neurosciences, Life Sciences Building 85, Highfield Campus, University of Southampton, SO17 1BJ, United Kingdom

The authors regret that there were errors in the original article text. It is correctly represented below.

The peptide sequence for A<sub>666</sub> should be reported as NH<sub>2</sub>-Leu-Glu-Pro-Arg-Trp-Gly-Phe-Gly-Trp-Trp-Leu-Lys-COOH. The cystine terminated versions of A<sub>665</sub> and A<sub>666</sub> for attachment to the PEG-*b*-PCL polymer should be reported as NH<sub>2</sub>-Cys-Leu-Ser-Thr-His-Thr-Thr-Glu-Ser-Arg-Ser-Met-Val-COOH and NH<sub>2</sub>-Cys-Leu-Glu-Pro-Arg-Trp-Gly-Phe-Gly-Trp-Trp-Leu-Lys-COOH, respectively. Iodoacetate-PEG6K-PCL19K (15 mg) was dissolved into DMF (1 mL) and 1 mg of a cysteine-terminated peptide was added.

The authors would like to apologise for any inconvenience caused.

DOI of original article: [10.1016/j.ijpharm.2011.12.042](https://doi.org/10.1016/j.ijpharm.2011.12.042).

\* Corresponding author.

E-mail addresses: [Elena.Surovtseva@uta.fi](mailto:Elena.Surovtseva@uta.fi), [esurovtseva@gmail.com](mailto:esurovtseva@gmail.com) (E.V. Surovtseva).