



Corrigendum

Corrigendum to “Improvement of survival in C6 rat glioma model by a sustained drug release from localized PLGA microspheres in a thermoreversible hydrogel” [International Journal of Pharmaceutics 427 (2012) 299–304]

Tetsuya Ozeki^{a,*}, Daiki Kaneko^b, Kosuke Hashizawa^b, Yoshihiro Imai^b, Tatsuaki Tagami^a, Hiroaki Okada^b

^a Drug Delivery and Nano Pharmaceutics, Graduate School of Pharmaceutical Sciences, Nagoya City University, 3-1 Tanabe-dori, Mizuho-ku, Nagoya, Aichi 467-8603, Japan

^b Department of Pharmaceutics and Drug Delivery, School of Pharmacy, Tokyo University of Pharmacy and Life Sciences, 1432-1, Horinouchi, Hachioji, Tokyo 192-0392, Japan

The authors regret the errors in original version of Figs. 2A and 5. We have reprinted Figs. 2A and 5 below. The authors would like to apologise for any inconvenience caused.

(A) CPT/PLGA




Fig. 2. SEM images of drug-incorporated PLGA microspheres. (A) CPT/PLGA microspheres.




Fig. 5. Whole body weight changes after the treatment with different formulations. Each point represents mean $\pm$ SE ($n = 5-7$). * $p < 0.05$ and *** $p < 0.005$, statistically significant compared with control rats.

DOI of original article: <http://dx.doi.org/10.1016/j.ijpharm.2012.02.012>.

* Corresponding author. Tel.: +81 52 836 3463; fax: +81 52 836 3463.

E-mail address: ozekit@phar.nagoya-cu.ac.jp (T. Ozeki).