

2-Soliton-Solution of the Novikov-Veselov Equation

J. Nickel^{1,2} and H. W. Schürmann¹

Received November 9, 2005; Accepted February 22, 2006

Published Online: May 16, 2006

Based on a superposition method recently proposed to obtain 1-solitary wave solutions of the KdV-Burgers equation (Yuanxi and Jiashi, 2005, *International Journal of Theoretical Physics* **44**, 293–301), we show that this method can also be used to find a 2-solitary wave solution of the Novikov-Veselov equation. Thus, it seems that the method of Yuanxi and Jiashi in general is not restricted to constructing 1-solitary wave solutions of nonlinear wave and evolution equations (NLWEs).

KEY WORDS: linear superposition; solitary wave solution; Novikov-Veselov equation.

1. INTRODUCTION

We analyze the Novikov-Veselov equation (NV equation) given by Hu (1994) which is another $(2+1)$ -dimensional analog of the KdV equation besides the well-known Kadomtsev-Petviashvili equation (Cheng, 1990). It has relevance in nonlinear physics (in particular in inverse scattering theory) (Athorne and Nimmo, 1991; Cheng, 1990; Hu, 1994; Hu and Willox, 1996; Konopelchenko and Moro, 1996; Tagami, 1989) and mathematics (cf. e.g. (Ferapontov, 1999; Taimanov, 1995)). To obtain a 2-soliton solution of the NV equation Hu has developed a nonlinear superposition formula (Hu, 1994, Equation (5)). In the following we show that a 2-solitary wave solution of the NV equation can even be obtained by a linear superposition method given by Yuanxi and Jiashi (2005) that was proposed to find 1-solitary wave solutions of nonlinear wave and evolution equations.

2. 2-SOLITON SOLUTION OF THE NV EQUATION OBTAINED BY LINEAR SUPERPOSITION

We consider the following equations

$$U_t + U_{xxx} + 3(U^2)_x = 0, \quad (1)$$

¹ Department of Physics, University of Osnabrück, D-49069 Osnabrück, Germany.

² To whom correspondence should be addressed at Department of Physics, University of Osnabrück, D-49069 Osnabrück, Germany; e-mail: jnickel@uos.de.

$$U_t + U_{yyy} + 3(U^2)_y = 0, \quad (2)$$

$$2U_t + U_{xxx} + U_{yyy} + 3(U \partial_y^{-1} U_x)_x + 3(U \partial_x^{-1} U_y)_y = 0. \quad (3)$$

Obviously, the KdV Eqs. (1), (2) and the NV Eq. (3) are related: The linear terms of Eq. (3) are equal to the superposition of those of Eqs. (1), (2) and the nonlinear terms of Eq. (3) are equal to the superposition of those of Eqs. (1) and (2) if traveling waves are considered with $\partial_x^{-1} = \partial_y^{-1}$. Following the ideas of Yuanxi and Jiashi (2005) we construct the solutions of Eq. (3) by linear superposition of those to Eqs. (1) and (2).

According to a method described by Schürmann and Serov (2004) we can evaluate the following 1-solitary wave solutions of Eqs. (1) and (2)

$$U(x, t) = \frac{c}{2k} \operatorname{sech}^2 \left[\frac{1}{2} \sqrt{\frac{c}{k^3}} (kx - ct - y_0) \right], \quad (4)$$

$$U(y, t) = \frac{c}{2k} \operatorname{sech}^2 \left[\frac{1}{2} \sqrt{\frac{c}{k^3}} (ky - ct - x_0) \right], \quad (5)$$

where x_0 and y_0 are arbitrary constants. Combining these solutions and choosing $k = 1$, so that $\partial_x^{-1} = \partial_y^{-1}$,

$$U(x, y, t) = \frac{c}{2} \operatorname{sech}^2 \left[\frac{1}{2} \sqrt{c} (x + y - ct) \right] \quad (6)$$

is a 1-solitary wave solution of Eq. (3).

We tentatively write a 2-solitary wave solution according to

$$U(x, y, t) = a \operatorname{sech}^2 \left[\frac{1}{2} \sqrt{c} (k_1 x + p_1 y - ct) \right] + b \operatorname{sech}^2 \left[\frac{1}{2} \sqrt{c} (k_2 x + p_2 y - ct) \right],$$

$$z_i = \frac{1}{2} \sqrt{c} (k_i x + p_i y - ct), \quad i \in \{1, 2\}, \quad (7)$$

with $c > 0$ and a, b, k_1, k_2, p_1, p_2 to be determined. Inserting Eq. (7) into Eq. (3) leads to

$$\begin{aligned} & \frac{2a\sqrt{c}\operatorname{sech}^2 z_1 \tanh z_1}{4k_1 k_2 p_1 p_2} (-2ck_1 k_2 p_1 p_2 (k_1^3 + p_1^3 - 2) \\ & - 6k_2 p_2 (2a - ck_1 p_1) (k_1^3 + p_1^3) \operatorname{sech}^2 z_1) \\ & - \frac{2a\sqrt{c}\operatorname{sech}^2 z_1 \tanh z_1}{4k_1 k_2 p_1 p_2} (6b(k_2 p_1 + k_1 p_2)(k_1^2 k_2 + p_1^2 p_2) \operatorname{sech}^2 z_2) \end{aligned}$$

$$\begin{aligned}
& + \frac{2b\sqrt{c}\operatorname{sech}^2 z_2 \tanh z_2}{4k_1 k_2 p_1 p_2} (-2ck_1 k_2 p_1 p_2 (k_2^3 + p_2^3 - 2) \\
& - 6k_1 p_1 (2b - ck_2 p_2) (k_2^3 + p_2^3) \operatorname{sech}^2 z_2) \\
& - \frac{2b\sqrt{c}\operatorname{sech}^2 z_2 \tanh z_2}{4k_1 k_2 p_1 p_2} (6a(k_2 p_1 + k_1 p_2) (k_1 k_2^2 + p_1 p_2^2) \operatorname{sech}^2 z_1) = 0. \quad (8)
\end{aligned}$$

Assuming $\frac{2a\sqrt{c}\operatorname{sech}^2 z_1 \tanh z_1}{4k_1 k_2 p_1 p_2} \neq 0$ and $\frac{2b\sqrt{c}\operatorname{sech}^2 z_2 \tanh z_2}{4k_1 k_2 p_1 p_2} \neq 0$ and setting the coefficients of $\operatorname{sech}^2 z_i$ equal to zero we obtain

$$a = \lambda_1 \frac{cp_2(p_2^3 - 2)^{1/3}}{2(p_2^3 - 1)^{2/3}}, \quad b = -\frac{1}{2}\lambda_2 cp_2(p_2^3 - 2)^{1/3}, \quad k_1 = \lambda_3 \left(2 - \frac{p_2^3}{p_2^3 - 1}\right)^{1/3}, \quad (9)$$

$$k_2 = -\lambda_4(p_2^3 - 2)^{1/3}, \quad p_1 = \lambda_5 \frac{p_2}{(p_2^3 - 1)^{1/3}} \quad \text{with}$$

	λ_1	λ_2	λ_3	λ_4	λ_5
I	1	1	1	1	1
II	1	$-(-1)^{1/3}$	$(-1)^{2/3}$	$-(-1)^{1/3}$	$-(-1)^{1/3}$
III	1	$(-1)^{2/3}$	$-(-1)^{1/3}$	$(-1)^{2/3}$	$(-1)^{2/3}$
IV	$-(-1)^{1/3}$	1	$(-1)^{2/3}$	1	$(-1)^{2/3}$
V	$-(-1)^{1/3}$	$-(-1)^{1/3}$	$-(-1)^{1/3}$	$-(-1)^{1/3}$	1
VI	$-(-1)^{1/3}$	$(-1)^{2/3}$	1	$(-1)^{2/3}$	$-(-1)^{1/3}$
VII	$(-1)^{2/3}$	1	$-(-1)^{1/3}$	1	$-(-1)^{1/3}$
VIII	$(-1)^{2/3}$	$-(-1)^{1/3}$	1	$-(-1)^{1/3}$	$(-1)^{2/3}$
IX	$(-1)^{2/3}$	$(-1)^{2/3}$	$(-1)^{2/3}$	$(-1)^{2/3}$	1

New solutions are given if $p_2 \in \mathbb{R}$ can be chosen so that k_i and p_1 are real, a and b may be complex. Subject to certain conditions some of these solutions are even physical (real and bounded) solutions. As an example the solution

$$\begin{aligned}
U(x, y, t) &= \frac{cp_2(p_2^3 - 2)^{1/3}}{2(p_2^3 - 1)^{2/3}} \operatorname{sech}^2 \left(\frac{\sqrt{c}(p_2 y + (p_2^3 - 2)^{1/3}x - c(p_2^3 - 1)^{1/3}t)}{2(p_2^3 - 1)^{1/3}} \right) \\
&- \frac{cp_2(p_2^3 - 2)^{1/3}}{2} \operatorname{sech}^2 \left(\frac{\sqrt{c}}{2}(-p_2 y + (p_2^3 - 2)^{1/3}x + ct) \right), \quad (10)
\end{aligned}$$

$$p_2 > \sqrt[3]{2} \quad (11)$$




Fig. 1. 2-soliton solution (cf. Eq. (7), I) for $t = -10, 0, 10, 20$ ($c = 1, p_2 = 1.5$).

(according to Eqs. (7), (9) and line I of the above table) is shown in Fig. 1. We have verified this solution by putting it into the original Eq. (3) utilizing Mathematica.

3. SUMMARY AND CONCLUDING REMARKS

By analyzing the structure of the NV equation we have shown that by using a superposition method proposed to construct 1-solitary wave solutions of NLWEES 2-solitary wave solutions can be obtained. We suppose that the technique of Yuanxi and Jiashi (2005) may lead to multi-solitary wave solutions of certain NLWEES if the NLWEE in question can be considered as a superposition of NLWEES that have the same type of solitary wave solution (e.g., sech^2); we leave this to future study.

ACKNOWLEDGMENTS

The work was supported by the German Science Foundation (DFG) (Graduate College 695 “Nonlinearities of optical materials”).

REFERENCES

- Athorne, C. and Nimmo, J. J. C (1991). *Inverse Problems* **7**, 809–826.
Cheng, Y. (1990). *Journal of Mathematical Physics* **32**, 157–162.

- Ferapontov, E. V. (1999). *Differential Geometry and its Applications* **11**, 117–128.
- Hu, X.-B. (1994). *Journal of Physics A: Mathematical and General* **27**, 1331–1338.
- Hu, X.-B. and Willox, R. (1996). *Journal of Physics A: Mathematical and General* **29**, 4589–4592.
- Konopelchenko, B. and Moro, A. (2004). *Journal of Physics A: Mathematical and General* **37**, L105–L111.
- Schürmann, H. W. and Serov, V. S. (2004). *Proceeding of Progress in Electromagnetics Research Symposium* March 28–31 (Pisa), 651–654.
- Tagami, Y. (1989). arXiv: dg-ga/9511005 v5 20 Nov 1995.
- Taimanov, I. A. (1995). *Physics Letters A* **141**, 116–120.
- Yuanxi, X. and Jiashi, T. (2005). *International Journal of Theoretical Physics* **44**, 293–301.