



Analysis of histone modifications from tryptic peptides of deuterioacetylated isoforms

Elisabeth Hersman¹, Dwella M. Nelson^{1,2}, Wendell P. Griffith³, Christine Jelinek, Robert J. Cotter^{*}

Pharmacology and Molecular Sciences, Johns Hopkins University School of Medicine, Baltimore, MD 21205, United States

ARTICLE INFO

Article history:

Received 14 February 2011

Received in revised form 19 April 2011

Accepted 22 April 2011

Available online 22 June 2011

Keywords:

Histones
Acetylation
Methylation
Orbitrap
Proteomics

ABSTRACT

The *in vitro* deuterioacetylation of histones obtained from biological sources has been used previously in bottom-up mass spectrometry analyses to quantitate the percent of endogenous acetylation of specific lysine sites and/or peptides. In this report, derivatization of unmodified lysine residues on histones is used in combination with high performance mass spectrometry, including combined HPLC MS/MS, to distinguish and quantitate endogenously acetylated isoforms occurring within the same tryptic peptide sequence and to extend this derivatization strategy to other post-translational modifications, specifically methylation, dimethylation and trimethylation. The *in vitro* deuterioacetylation of monomethylated lysine residues is observed, though dimethylated or trimethylated residues are not derivatised. Comparison of the relative intensities ascribed to the deuterioacetylated and monomethylated species with the deuterioacetylated but unmethylated analog, provides an opportunity to estimate the percent of methylation at that site. In addition to the observed fragmentation patterns, the very high mass accuracy available on the Orbitrap mass spectrometer can be used to confirm the structural isoforms, and in particular to distinguish between trimethylated and acetylated species.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

In the nucleus, DNA is packaged as nucleosomes consisting of 146-basepairs of double-stranded DNA wrapped around the core histones H2A, H2B, H3 and H4 [1]. Each histone has two general domains, an inner core region and an exposed N-terminal “tail”, which can be post-translationally modified with acetylation, methylation, phosphorylation or other modifications [2]. Specific post-translational modifications (PTMs) and combinations of PTMs are proposed to function as a *histone code* that directs gene silencing, transcription, replication, cellular memory and gene recombination [3–5]. Elucidating histone post-translational modifications has thus become essential to understanding epigenetics.

One approach to characterizing histone PTMs is to use “bottom-up” mass spectrometric analysis [6], which begins with proteolytic digestion of the histone into peptide fragments. The molecular weight and mass spectral fragmentation pattern of each of these peptides are then used to determine its amino acid sequence and

any modifications. The most commonly used protease is trypsin, which cleaves the amide peptide bond on the carboxyl side of the basic amino acid residues lysine and arginine except before proline [7]. Proteolysis by trypsin typically yields peptides ranging 200–4000 Da, which are readily detected by mass spectrometry. However, histones have an abundance of lysine and arginine residues, so that tryptic digestion yields many small peptides that are not easily observed and comprise too few and repetitive sequences that make it difficult to establish the location of the modification. Moreover, the principal sites for modification by acetylation, methylation, ubiquitylation, etc., are the lysine residues. When modified, these sites generally become “missed cleavages” [8], producing different peptide segments for the modified and unmodified forms that are then difficult to compare quantitatively. Smaller fragments are also less likely to encompass multiple modification sites, and therefore do not provide quantification for combinatorial isoforms.

The “top-down” or “middle-down” strategies used by Kelleher et al. [9], along with derivatization of lysine residues described by Garcia et al. [8,10], address some of these problems by using intact or longer sequences, and by direct fragmentation of these via *electron capture dissociation* (ECD) on a Fourier transform mass spectrometer (FTMS) [11–13]. It is assumed that the number and/or position of basic residues for the intact proteins will have minimal effect, compared with smaller structures, on their chemistry and therefore on the opportunities for relative quantitation of

* Corresponding author. Tel.: +1 410 955 3022; fax: +1 410 955 3420.

E-mail address: rcotter@jhmi.edu (R.J. Cotter).

¹ These authors contributed equally to this study.

² Current address: Division of Hematology, Johns Hopkins University, Baltimore, MD 21205, United States.

³ Current address: Department of Chemistry, University of Toledo, Toledo, OH 43606-3360, United States.

the different isoforms [14,15]. Furthermore, all possible structural/positional combinations will be intact and Kelleher et al., using hydrophilic interaction liquid chromatography (HILIC) in combination with FTMS have been able to identify and distinguish more than 150 differentially modified forms of histone H3.2 in HeLa cells using a top-down approach [9].

Alternatively, several groups have developed lysine-specific derivatization techniques that modify histone samples prior to digestion and mass spectrometric analysis [16–19]. Smith et al. used a mixture of deuterated (d_4) acetic acid and deuterated (d_6) acetic anhydride as the derivatization reagent, converting all endogenously unmodified lysines present in the sample to deuterioacetylated species, with a mass shift of +45 Da [16,17]. These label-modified lysines are then chemically and chromatographically equivalent to their *in vivo* acetylated counterparts, but distinguishable by a three Dalton mass difference (+45 Da vs. +42 Da, respectively). All lysine residues are then equivalently “blocked” from cleavage by trypsin, cleavage occurs at arginine residues only, and the fragments generated are now larger in size, providing in some cases the ability to observe multiple modifications. From the mass spectra, three Dalton mass shifts are used to determine and quantitate endogenously acetylated (d_0) and *in vitro* deuterioacetylated (d_3) peptides. For peptides containing multiple lysines, this approach can quantitate the numbers of lysines acetylated, distinguishing and quantitating their positional isomeric forms. When additional modifications, such as methylation, are also present in the same peptide, this approach can be used to compare quantitatively those isoforms differing in acetylation but having a common methylation, or other modification, site. For these isoforms, the deuterioacetyl label eliminates several “biases” common in quantitation methods, since enzymatic digestion, chromatographic separation, and ionization efficiency (in the mass spectrometer) are effectively normalized. In our laboratory we have used deuterioacetylation with MALDI tandem time-of-flight mass spectrometry previously to determine the percent acetylation at K56 on histone H3_{54–63} [FQK₅₆STELLIR] in yeast deficient in sirtuins Hst3 and Hst4p [20] and identified the major isoforms spanning residues K5, K8, K12, and K16 in yeast histone H4 [21].

An alternative approach uses propionic anhydride to form propionyl analogs that differ in mass from the acetylated peptides by 14 Da and obviate the need for isotopic corrections to the relative abundances [22]. This approach has been used as well in a two step derivatization process, carried out before and after tryptic digestion, in which the second derivatization carries a stable isotope d_5 analog to enable quantitative comparisons between two samples [23]. In our study, which focuses on the degree of acetylation at specific sites in a single sample, the use of a stable isotope analog was thought to provide the most similar chromatographic behavior, ionization efficiency and quantitative accuracy.

While the top-down approach would appear to be the most direct, intact histone isoforms are relatively resistant to reverse-phase chromatographic separation [9], and the targeting of individual isoforms for data-dependant MS/MS analysis becomes exceedingly difficult as multiple precursor ion species with similar retention times can result in missing the lower abundance forms. Methods to improve the top-down approach include the use of hydrophilic interaction chromatography (HILIC) to better separate histone mixtures [24] and “middle-down” mass spectrometric analysis using endoprotease Glu-C digestion to enable focusing on the so-called “tail regions” where the bulk of the modifications are found [25,26].

In this context, we have continued to develop a bottom-up approach that exploits and extends the advantages of prior lysine deuterioacetylation using high performance mass spectrometry and the application of the approach to other modifications. Specifically, both MALDI tandem time-of-flight (TOF) and high resolution

nanospray/Orbitrap mass spectrometry are used here to characterize the positional microheterogeneity of the histone H4 tail region acetylation isoforms in HeLa cells and to observe the specific changes to the H4 tail region for cells treated with trichostatin A (TSA) and nicotinamide (NIA), two broad spectrum inhibitors of histone deacetylases (HDACs). In addition, deuterioacetylation enables the observation of methylated (lysine and arginine) species in the context of larger peptides that may also be acetylated. In several examples reported here it is shown that monomethylated lysines are derivatized by deuterioacetylation, while dimethylated and trimethylated lysines are not. While the ability to distinguish isobars with the same nominal mass is accomplished relatively quantitatively for methylated as well as acetylated species using this approach, the additional advantages of high mass resolution and accuracy to distinguish acetyl and trimethyl modifications are also shown.

2. Materials and methods

2.1. Preparation of yeast histones

Yeast cells containing the appropriate plasmids were grown as reported previously [20]. Briefly, a 25 mL culture of yeast cells were inoculated in YPD medium and grown for 9–10 h at 30 °C with shaking. This was transferred to 1 L YPD and grown overnight to an OD₆₀₀ of 1. Cells were centrifuged at 500 × g for 5 min and the resultant pellets were washed in 100–200 mL of sterile water and pooled. The pooled pellet was re-spun, collected and resuspended in 20 mL of DTT/Tris buffer pH 9.4 followed by incubation with shaking for 15 min at 30 °C and centrifugation. The cell pellet was resuspended in 20 mL sorbitol/HEPES buffer pH 7.4 and lysed by adding 0.5 mg zymolase and incubating at 30 °C with gentle shaking (160 rpm) until cell wall digestion was complete. Spheroplasts were lysed in a buffer containing 0.5% NP-40. Histones were extracted with 0.25 N hydrochloric acid, precipitated at –20 °C with 8 volumes of cold acetone and centrifuged at 30K × g, 4 °C for 30 min. The resultant histone-enriched pellet was resuspended in Tris-HCl buffer pH 8.0 and resolved in a 15% SDS polyacrylamide gel.

2.2. HeLa cell culture and histone purification

HeLa cells were cultured in Dulbecco's modified eagle's medium (DMEM) containing 10% fetal bovine sera (GIBCO; Invitrogen, Carlsbad, CA). Cells were plated and grown to 90% confluence. One sample was treated with deacetylase inhibitors: 1 μM trichostatin A (TSA, Sigma-Aldrich, St. Louis, MO) and 10 mM nicotinamide (NIA, Sigma-Aldrich, St. Louis, MO) for 3 h while the other sample was supplemented with an equal volume of solvent alone. TSA and NIA were dissolved in DMSO and deionized water, respectively. Cells were then harvested, pelleted and fractionated to isolate nuclei. Histones were acid extracted as described in the method by Wang et al. [23] that provides relatively purified histones, and resolved by SDS-PAGE (4–12% NuPAGE™ gels; Invitrogen, Carlsbad, CA) with purified chicken core histones (Millipore; Billerica, MA) run as markers, and the gel was stained with Coomassie Brilliant Blue (CBB).

2.3. Deutero (d_3)-acetylation, chemical derivatization and digestion

Using reagents provided in the Trypsin Profiler IGD Kit for in-gel digestions (Sigma-Aldrich, St. Louis, MO), CBB-stained HeLa H4 gel bands were excised, cut into 1 mm pieces and destained for 1 h with a 200 mM ammonium bicarbonate solution containing 40% acetonitrile. The destained protein bands were resuspended in a solution

containing 50 μL deuterated (d_4) acetic acid and 10 μL deuterated (d_6) acetic anhydride (deuterated reagents; Sigma–Aldrich, St. Louis, MO). To ensure the derivatization reaction went to completion, gel bands were incubated in the deuterated reagents for 5 h at room temperature. The treated gel slices were then rinsed with distilled water, titrated with 200 mM ammonium bicarbonate (Sigma–Aldrich, St. Louis, MO) to pH 8, rinsed with water again to remove remaining buffer salts, and dehydrated/dried in a Speed Vac. Proteins were digested in-gel with 0.4 μg trypsin (1 mg/mL in 1 mM HCl) in 40 mM ammonium bicarbonate containing 9% acetonitrile, incubated overnight at 37 °C [27]. Resultant peptides were extracted from the gel and lyophilized to dryness before mass spectrometric analysis. All samples were reconstituted as stock solutions in 5 μL water with 0.1% trifluoroacetic acid (TFA; Pierce, Rockford, IL).

2.4. MALDI-TOF mass spectrometry

Matrix solution was prepared as 10 mg/mL α -cyano-4-hydroxycinnamic acid (Sigma Aldrich, St. Louis, MO) dissolved in a 1:1 solution of acetonitrile:deionized water containing 0.1% TFA. Undissolved matrix particles were removed via centrifugation. Using the dried droplet application technique, resultant matrix solution (1 μL) and the resuspended histone H3 and H4 peptide mixtures (1 μL) were spotted on the stainless steel MALDI target [28]. To ensure co-crystallization, target plates were spotted at room temperature using the sandwich method: deposit 1 μL of matrix, 1 μL of digest and 1 μL of matrix, then redissolve in 50% AcN containing 0.1% TFA.

Deuteroacetylated HeLa histone H4 peptides were analyzed using a Shimadzu AXIMA-TOF² tandem time-of-flight mass spectrometer (Manchester, UK) equipped with a 337 nm pulsed nitrogen laser, high energy collision chamber and curved-field reflectron [29,30]. The acceleration voltage was set at 20 kV. A four point, external calibration was applied per sample spot using a peptide mixture containing bradykinin (m/z 757.40), angiotensin II (m/z 1046.54), P₁₄R (m/z 1533.86), ACTH fragment 18–39 (m/z 2465.20), and insulin B chain (m/z 3495.65). All peptides used in the calibration mixture were from Sigma Aldrich (St. Louis, MO). Each mass spectrum acquired consisted of an average of 500 profiles of 10 shots accumulated per profile.

2.5. Sample preparation for HPLC-ESI-MS/MS analysis

Four microliters of stock deuteroacetylated peptide mixture were diluted to 2.5 fold (to a final volume of 10 μL) and placed in an Agilent 1200 autosampler (Agilent, Santa Clara, CA); 5 μL of each 10 μL sample were loaded by the autosampler through a trapping column onto a fused silica PicoFrit (New Objective, Woburn, MA) capillary column, 75 μm inner diameter (i.d.) \times 120 mm long column packed in-house with 5 μm , 300 Å BioBasic C₁₈ (Thermo Electron, Bremen, Germany) stationary phase, at a flow rate of 300 $\mu\text{L}/\text{min}$. Peptides were separated on-line via reversed phase nano high-performance liquid chromatography using the Eksigent Nano 2D high-performance liquid chromatography (HPLC) pumping system (Eksigent, Dublin, CA). The Eksigent Nano 2D HPLC system was controlled by XCalibur software, Version 2.0 (Thermo Electron, San Jose, CA). Separations were performed at mobile phase flow rate of 300 nL/min on the binary pump system using 0.1% formic acid in deionized water (mobile phase A) and 90% (v/v) acetonitrile with 0.1% (v/v) formic acid (mobile phase B) using a linear gradient of 20–50% A over 60 min. The outlet flow of the nano-HPLC interfaced directly with the inlet of an LTQ-Orbitrap-XL (Thermo Electron, San Jose, CA) allowing for introduction of the analyte into the mass spectrometer.

2.6. Tandem mass spectrometric analysis (LTQ-Orbitrap XL)

The LTQ-OrbitrapXL mass spectrometer was operated in *data dependent* mode. MS precursor scan spectra (m/z 300–2000) were acquired in the Orbitrap with mass resolution of 60,000; the six most intense ions from each MS scan were automatically targeted for fragmentation (MS/MS) in the ion trap. Using nitrogen as the collision gas, collision induced dissociation (CID) mediated peptide fragmentation in the linear ion trap. The LTQ-OrbitrapXL was controlled by XCalibur software. For each experiment, the source voltage was set at 2.4 V, the capillary voltage at 48 V, and the capillary temperature at 200 °C. Sheath and auxiliary gases were not necessary because nanoflow parameters were being used. The tube lens voltage was kept at 105 V, and the ion gauge pressure was 1.5×10^{-5} Torr. The automatic gain control (AGC) was used at the manufacturers default settings for MS collection in the orbitrap (200,000) and MS/MS collection in the ion trap (10,000). Normalized collision energy was established at 35% for MS/MS. The default charge state was set at two. The isolation window for precursor ion selection was fixed at 2 Da. After an initial MS/MS fragmentation event, ions were excluded from additional fragmentation rounds for 30 s using dynamic exclusion. The ion selection threshold, the minimum signal required to trigger tandem mass spectrometry, was set to 500. The activation Q was set at 0.25.

2.7. Quantitation of peptide isoforms

Determining the relative ratio between acetyl (d_0) and deuteroacetyl (d_3) ion pairs gives the relative abundance of the *in vivo* acetylated and unmodified peptide forms. The relative acetylation of each peptide was determined by an in-house processing method that compares the peak height of acetylated peptides to the total for all observed forms of the peptide as measured from the charge-state reduced chromatogram of the peptides' elution peak (generated using Xtract software package within XCalibur). Charge state reduction was employed such that the peak intensities from all charge states of each species were summed over the elution profile. Determining the relative ratio between acetyl (d_0) and deuteroacetyl (d_3) ion pairs gives the relative abundance between *in vivo* acetylated and unmodified peptide forms. To determine the relative abundance of positional isomers, where all isomeric forms fragment from the same precursor mass in the same MS/MS scan, we compared the peak intensities from selected b and y ion pairs distinguishing these isoforms [31]. Peak intensities from b and y ion pairs were summed together and this value was then divided by the total produced between both species.

3. Results and discussion

3.1. Acetyl quantitation

Fig. 1 shows the MALDI time-of-flight mass spectrum of the tryptic digest of deuteroacetylated histone H3 from the *hst3-H184A* yeast mutant, a deletion mutant of the Sirtuin (HDAC) Hst3. Several of the tryptic peptides expected or identified previously are shown [20]. The peptide FQKSTELLIR encompasses the K56 residue, an acetylation site just inside the histone core region. In the expanded mass spectrum (Fig. 1, inset) the acetylated FQK_{Ac}STELLIR form is observed at m/z 1276.94 along with its molecular ion distribution resulting from naturally occurring isotopes. Without derivatization, the unacetylated analog could not be observed, as K₅₆ would be a tryptic cleavage site. With derivatization, the unacetylated analog is also observed as the deuteroacetylated peptide FQK_{dAc}STELLIR three mass units higher at m/z 1279.95. Quanti-




Fig. 1. MALDI time-of-flight mass spectrum of the tryptic digest of purified and deuterioacetylated histone H3 from yeast cells. Masses corresponding to the peptides expected from several of the known post-translational modifications are annotated. Inset: expansion of the H3₅₄₋₆₃ peptide showing the distribution of naturally acetylated and deuterioacetylated species.

tative assessment of the degree of acetylation at this site is then based upon the fact that both species are chemically equivalent and that the ionization efficiencies are also equivalent. In this case 43% acetylation is determined after correction for the contribution of the 3rd isotope peak in the isotopic distribution of the naturally acetylated species to the peak intensity of the monoisotopic peak of the deuterioacetylated species. To determine the distribution in acetylation, the isotopic distributions of the endogenous and deuterioacetylated H4 peptides were simulated based on calculations using MS-isotope (Protein Prospector Software), available online at <http://prospector.ucsf.edu/>. For this mass and elemental composition the 3rd isotopic peak in the naturally acetylated peptide will have an area of 8.60% of that of its monoisotopic mass [see Supplemental Figure S1].

It should be noted that without derivatization the absence of a particular acetylated peptide would not preclude its presence in the sample, as this could be the result of poor ionization efficiency or other factors which generally limit full sequence coverage. However, the absence of acetylated peptides in the presence of their deuterioacetylated analogs can be definitive. Conversely, we were able to show previously that the yeast *hst3hst4* double deletion mutant did indeed result in nearly 100% acetylation at this site [20], as little or no deuterioacetylated analogs were observed in this mutant, while they were observed in the single *hst3* or *hst4* mutants.

3.2. Acetylated H4 tail peptide GK₅GK₈GLGK₁₂GGAK₁₆R

Fig. 2 is the MALDI-TOF mass spectrum of the tryptic digest of deuterioacetylated histone H4 from HeLa cells, again showing the most prominent expected peptides. Of particular interest is the histone H4 tail peptide GKGGKGLGKGGAKR encompassing residues 4–17. It includes four lysine residues, and in their 2003 report Smith et al. [16] had previously shown how conversion of all the unmodified lysine residues to their d₃ deuterioacetylated derivatives could

be used to quantitatively assess the distribution of tail peptides containing between 0 and 4 acetylated lysines.

In our laboratory, this same approach was used with both MALDI-TOF and ESI Orbitrap mass spectrometry to monitor effects of the HDAC inhibitors trichostatin A and nicotinamide on the extent of acetylation and on the specific distribution of the positional isomeric forms. The MALDI mass spectrum in Fig. 3(a) was used to determine the number of acetylated lysine residues for the peptide GKGGKGLGKGGAKR obtained from HeLa cells treated with TSA/NIA. Following derivatization all lysine residues are either acetylated (+42 Da) or deuterioacetylated (+45 Da), so that the lowest mass d₀ at m/z 1439.03 in this isotopic distribution corresponds to GKGGKGLGKGGAKR that is fully acetylated. A single acetylation corresponds to the d₉ peak, containing one acetylated lysine and three deuterioacetylated lysines. The unacetylated peptide is the d₁₂ peak at m/z 1451.09. All of the peptides are chemically equivalent and the distribution is obtained from the peak heights of the d₀, d₃, d₆, d₉ and d₁₂ species.

Then, when calculating the actual abundance of each differently modified peptide, the contribution of the 3rd isotope peak of the lighter isoforms to the intensity of the monoisotopic peak of the isoform heavier by 3 Da must be subtracted. For example, the corrected intensity of the d₃ GK₅GK₈GLGK₁₂GGAK₁₆R peptide species is calculated by subtracting 10.75% of the d₀ monoisotopic peak from the measured d₃ peak intensity [see Supplemental Figure S2]. Corrected distributions for the unacetylated to tetra-acetylated species obtained in the MALDI-TOF mass spectra are shown in Table 1 for both the endogenous and deacetylase inhibited HeLa cell samples.

Fig. 3(b) shows the analogous distribution for the histone tail peptide from untreated HeLa cells. Peptides with four acetylated lysines are not observed, and in fact three acetylated lysines are of very low abundance. Most of the peptides are unacetylated or mono-acetylated. Comparison with Fig. 3(a) shows the effect of the HDAC inhibitors in promoting a high degree of acetylation. From Table 1, approximately 57% of isomers present in the endogenous




Fig. 2. MALDI time-of-flight mass spectrum of the tryptic digest of purified and deuterioacetylated histone H4 from HeLa cells. Masses corresponding to the peptides expected from several of the known post-translational modifications are annotated.

HeLa H4 tail peptide had no acetylated lysine, i.e., the most abundant isoform of this peptide was unmodified. 31% of the peptide had one acetylated lysine, and 9% had two acetylated lysine residues. Less than 4% of the $\text{GK}_5\text{GGK}_8\text{GLGK}_{12}\text{GGAK}_{16}\text{R}$ isoforms contained three or four acetylated lysines.

There are sixteen potential positional isomeric forms of the acetylated/deuterioacetylated peptide $\text{GK}_5\text{GGK}_8\text{GLGK}_{12}\text{GGAK}_{16}\text{R}$, including four mono-, six di- and four tri-acetylated species. Quantitation of positional isomers having the same precursor mass could not be determined using MALDI tandem analysis. Time-of-flight instruments generally have somewhat limited mass selection capability, so one cannot select isotopically pure precursors. Nonetheless, our preliminary work with this instrument indicated that the major monoacetylated species was acetylated at K16, while the major diacetylated species was acetylated on K8 and K16 [21]. This of course motivated our use of the LTQ/Orbitrap mass spectrometer to provide detailed analysis of the positional isomers.

3.3. Quantitation of H4 tail peptide isoforms using ESI orbitrap MS/MS analysis

Tryptic digests of the deuterioacetylated histones were analyzed using on-line HPLC MS/MS. Quantitation based upon a comparison of relative peak heights is easily accomplished when the comparison is made between chemically and chromatographically equivalent species, specifically species differing only in isotopic composition. Thus, all 16 of the acetylated/deuterioacetylated iso-

forms of the tail peptide $\text{GK}_5\text{GGK}_8\text{GLGK}_{12}\text{GGAK}_{16}\text{R}$ eluted from the column at the same time (Fig. 4). While some small chromatographic differences were detected for peptides with different numbers of *in vivo* acetylations, by contrast positional isoforms, with the same number of naturally occurring acetylations (and same molecular weights), had identical narrow retention times. Fig. 5 is the ESI Orbitrap mass spectrum of the doubly charged molecular ions of the tail peptides from the (a) untreated and (b) TSA/NIA treated HeLa samples. The extracted (isotopically corrected) relative abundances of the unacetylated to tetra-acetylated species are shown in Table 1, and are similar to those obtained by MALDI.

Doubly charged molecular ions of the unacetylated to tetra-acetylated tail peptides $\text{GK}_5\text{GGK}_8\text{GLGK}_{12}\text{GGAK}_{16}\text{R}$ were then each targeted for MS/MS fragmentation to quantitate the distributions of their positional isomers. Tandem mass spectra collected from




Fig. 3. Expansion of the MALDI mass spectral region for the deuterioacetylated H4_{4-17} peptide GKGGKGLGKGGAKR showing the distribution of the non-acetylated (d_{12}) to tetra-acetylated (d_0) species for (a) histone derived from TSA-treated HeLa cells and (b) histone from untreated HeLa cells.

Table 1

Relative abundances of each differentially modified peptide population for the model GKGGKGLGKGGAKR peptide. Comparison of isotopically corrected data derived from both the LC-ESI MS and MALDI-TOF.

	No deacetylase inhibitor		With deacetylase inhibitors Trichostatin and Nicotinamide	
	MALDI-TOF	ESI Orbitrap	MALDI-TOF	ESI Orbitrap
No acetylation	57%	63%	10%	8%
1 acetylation	31%	29%	19%	14%
2 acetylations	9%	7%	21%	20%
3 acetylations	3%	1%	22%	26%
4 acetylations	0%	0.3%	28%	32%




Fig. 4. Chromatographic traces for the acetylated/deuteroacetylated isoforms of the H4_{4–17} peptide GKGGKGLGKGGAKR obtained by monitoring the doubly charged molecular ions on an LTQ/Orbitrap mass spectrometer. Separation is by C18 column interfaced directly to a nanospray ionization source.


each precursor mass were summed using the processing feature of Qual Browser (Xcaliber, Thermo Electron, San Jose, CA) and the average of the peak intensities from the fragment ions was used to determine the percentage of each fragment ion in the spectra. Calculated fragment ion masses for each were generated using Protein Prospector MS-Product. The standard deviation for the calculated percentage was determined by comparing the ratios of b-series and y-series ion intensities corresponding to cleavage at the same bond (e.g., y5 and b9, or y7 and b5) on the endogenous and deacetylase treated peptides, and was determined to be less than 5%.

Fig. 6 shows the MS/MS spectrum of the doubly charged molecular ion *m/z* 719.90, which corresponds to the *d*₀ (fully acetylated)

Table 2
Fragment ions of monoacetylated GKGGKGLGKGGAKR peptide from histone H4 from HeLa cells: fractional abundance of y and b ion pairs.

Fragment ions	acK5	acK8	acK12	acK16
y5 and b9	0.256	0.744		
y7 and b5	0.127	0.873		
y11 and b3	0.073	0.927		
Untreated HeLa cells	7%	5%	13%	74%
y5 and b9	0.303	0.697		
y7 and b5	0.146	0.854		
y11 and b3	0.063	0.937		
TSA-treated HeLa cells	6%	8%	16%	70%

species from the TSA/NIA-treated sample. There is only one positional isomer with this molecular weight giving rise to a single set of b-series and y-series ions. From these we selected three b and y fragment ion “pairs” that describe the fragmentation between the four lysine residues:



In choosing these ions we selected those that appear with reasonable intensity in all of the subsequent spectra of mixed isoforms.

Fig. 7 then shows expansions of the mass regions about these six fragment ion masses for the mono-acetylated isomers, of which there are four. Each of these b and y ions appears at two masses and can be used to describe the relative abundances of the isomeric forms. For example, *b*₃ shows the ratio between ions acetylated at K₅ and those acetylated at any of the other three lysines. The *y*₁₁ fragment does the same, though the order of mass is reversed. These two ratios were averaged to produce the entry in Table 2 of 7.3% acetylation at K₅ and 93% acetylation on all other lysines. The *b*₅ and *y*₇ ratios then determine the relative abundances of acetylation at K₅ or K₈ versus K₁₂ or K₁₆. And the *b*₉ and *y*₅ ratios determine the percentage acetylated at K₁₆. Combining this data (Table 2), it is possible to determine that the composition for the




Fig. 5. Nanospray LTQ/Orbitrap mass spectra of the doubly charged molecular ion regions for the acetylated/deuteroacetylated isoforms of peptide GKGGKGLGKGGAKR integrated across the chromatographic retention times shown in Fig. 4. (a) untreated HeLa cells and (b) TSA-treated HeLa cells.




Fig. 6. Full scan nanospray LTQ/Orbitrap MS/MS spectrum for the doubly charged molecular ion of the fully acetylated (d_0) isoform of the peptide GKGGKGLGKGGAKR highlighting the b-series (b_3 , b_5 and b_9) and y-series (y_5 , y_7 and y_{11}) ions used to quantitate the positional isomers.

mono-acetylated isomers from untreated cells is 7% K_5 , 5% K_8 , 13% K_{12} and 74% K_{16} .

This computational assessment agrees with earlier MALDI-TOF results showing $K_{16}Ac$ is the major mono-acetylated isoform [21]. Our findings are also consistent with previous biological studies. Histone H4 K_{16} acetylation is a reversible modification implicated

in the widespread process of chromatin condensation by recruiting HATs to acetylate K_{12} , K_8 , and K_5 . In mammals, K_{16} acetylation also correlates with K_{20} trimethylation [32]. Using an antibody approach to study site specific lysine acetylation in yeast histone H4, Suka et al. [33] found that lysine acetylation in the H4 N-terminal peptide_{4–17} occurs sequentially from the C-terminus to




Fig. 7. Expanded mass spectral regions of the b-series (b_3 , b_5 and b_9) and y-series (y_5 , y_7 and y_{11}) ions used to quantitate the positional isomers of the monoacetylated (d_9) GKGGKGLGKGGAKR peptide.

Table 3
Fractional abundances of fragment ions of di-acetylated GKGGKGLGKGGAKR peptide from histone H4 from HeLa cells.

Fragment ions	acK5 acK8	acK5 acK12	acK5 acK16	acK8 acK12	acK8 acK16	acK12 acK16
y7 and b5	0.056	0.541				0.402
y11 and b3	0.255			0.745		
Untreated HeLa cells	6%	20%		34%		40%
y7 and b5	0.032		0.373			0.595
y11 and b3	0.168			0.832		
TSA-treated HeLa cells	3%		14%		24%	59%

the N-terminus: K16 acetylation precedes K12Ac, which precedes K8Ac, leading finally to K5Ac. We found similar results for H4 tail peptides from TSA/NIA treated HeLa cells: 6% of the monoacetylated peptide was acetylated at the K5 residue, 8% at the K8 residue, 16% at K12, and 70% at K16 (Table 2).

There are six potential isomeric forms of the diacetylated H4 tail peptide_{4–17}. It is not possible to distinguish all the diacetylated isomeric species from unique fragment masses. However, we determined that about 6% of untreated diacetylated peptide_{4–17} was acetylated at the K5 and K8 residues, whereas 20% were mixtures of acK₅acK₁₂ and acK₅acK₁₆, 34% comprised mixtures of acK₈acK₁₂ and acK₈acK₁₆, and 40% were acK₁₂acK₁₆ (Table 3). When the deacetylase inhibitor-treated histone H4 sample was evaluated, the most dominant diacetylated isomer was the acK₁₂acK₁₆ peptide, which constituted 59% of all diacetylated species. Three percent of the treated, diacetylated peptide_{4–17} was acK₅acK₈, 14% was the combination of acK₅acK₁₂ and acK₅acK₁₆, and 24% was the combination of acK₈acK₁₂ and acK₈acK₁₆. Results for the tri-acetylated peptide species are summarized in Table 4. In untreated samples, 23% of the triacetylated peptide_{4–17} was acK₅acK₈acK₁₂, 10% was acK₅acK₈acK₁₆, 28% was acK₅acK₁₂acK₁₆, and acK₈acK₁₂acK₁₆ comprised 39%. For the TSA/NIA samples, 6% of all triacetylated peptide_{4–17} species were acK₅acK₈acK₁₂, 11% of the total was acK₅acK₈acK₁₆, 20% was acK₅acK₁₂acK₁₆ and 62% was acK₈acK₁₂acK₁₆.

Combining these distributions with those in Table 1, we can determine the relative abundance for all positional isomers. Thus Table 5 displays the relative abundance of all 16 acetylated isoforms for the tail peptide. Without deacetylase inhibition, the most abundant peptide isoform is unacetylated (63%). The second most abundant isoform had a single acetylation at residue K16 (22%). In the presence of deacetylase inhibitors, as would be predicted, the most abundant peptide is fully acetylated (32%). Compared to untreated controls, no single modification profile predominates; however, isoforms that include modifications at K16 are generally more abundant.

3.4. Methylation of the K₇₉ site in histone H3

It is more difficult to design an analogous strategy that uses isotopically labeled methylation to elucidate methylation sites for two

Table 4
Fractional abundances of fragment ions of tri-acetylated GKGGKGLGKGGAKR peptide from histone H4 from HeLa cells.

Fragment ions	acK5	acK5	acK5	acK8
	acK8	acK8	acK12	acK12
	acK12	acK16	acK16	acK16
b9 and y5	0.231	0.768		
y7 and b5	0.327	0.673		
y11 and b3	0.607	0.392		
Untreated HeLa cells	23%	10%	28%	39%
b9 and y5	0.064	0.936		
y7 and b5	0.174	0.826		
y11 and b3	0.379	0.621		
TSA-treated HeLa cells	6%	11%	20%	62%

reasons: because lysines may be mono-, di- and tri-methylated and because methylation can also occur on arginine residues. However, our deuteroacetylation strategy has also proven to be an effective strategy for resolving methylated structures as lysine derivatization restricts tryptic digestion to arginine residues, producing analogous longer peptide fragments that contain either/both acetylation and methylation modifications.

Fig. 8(a) is the MALDI-TOF mass spectrum of the yeast histone H3 fragment EIAQDFKTDLR corresponding to residues 73–83. This peptide fragment was obtained from incomplete tryptic digestion of underivatized wild-type histone H3 and reveals peaks corresponding to peptides unmodified, methylated, dimethylated and trimethylated at K₇₉, though at this resolution the latter would not be distinguishable from acetylation. [The presence of trimethylation is shown below in our results using the Orbitrap]. Tryptic digestion of the deuteroacetylated histone H3 fragment in Fig. 8(b) is interesting, because both the unmethylated and mono-methylated species have been derivatized, while the dimethylated and trimethylated (or acetylated) species are not derivatized. Specifically the unmodified peptide seen at *m/z* 1335.85 in Fig. 8(a) now appears at *m/z* 1380.97 as the deuteroacetylated species three mass units above the trimethylated (or acetylated) species at *m/z* 1377.82. Derivatization of the mono-methyl species is observed in Fig. 8(b) at *m/z* 1386.99. The equivalent derivatization of the monomethyl, but not the dimethyl and trimethyl, forms by propionylation has also been observed [23].

Electrospray mass spectra were also obtained for a yeast histone H3 sample containing unmodified and methylated K₇₉. When deuteroacetylated, tryptic digestion produced the methylated/deuteroacetylated and deuteroacetylated species whose MS/MS spectra are shown in Fig. 9, where the high mass accuracy




Fig. 8. MALDI time-of-flight mass spectrum of the yeast histone H3_{73–83} peptide EIAQDFKTDLR showing (a) unmethylated peptides and peptides methylated, dimethylated and trimethylated at lysine 79. (b) Deuteroacetylated peptide showing derivatization of both unmethylated and monomethylated species.

Table 5

Relative abundance of all 16 positional differentially modified peptides with the GKGGKGLGKGGAKR sequence for Histone H4 sample from both untreated HeLa cells and HeLa cells treated with deacetylase inhibitors TSA and NIA (lysine residues in red are acetylated).

Isoforms		No deacetylase inhibitors	With deacetylase inhibitors
0	G K G G K G L G K G G A K R	63%	8%
	G K G G K G L G K G G A K R	2 %	1 %
	G K G G K G L G K G G A K R	1 %	1 %
	G K G G K G L G K G G A K R	4 %	2 %
	G K G G K G L G K G G A K R	22 %	10 %
1 acetylation	G K G G K G L G K G G A K R	0.4 %	0.6 %
	G K G G K G L G K G G A K R	1 %	3 %
	G K G G K G L G K G G A K R	2 %	5 %
	G K G G K G L G K G G A K R	3 %	12 %
	G K G G K G L G K G G A K R	3 %	12 %
2 acetylations	G K G G K G L G K G G A K R	0.2 %	2 %
	G K G G K G L G K G G A K R	0.1 %	3 %
	G K G G K G L G K G G A K R	0.3 %	5 %
	G K G G K G L G K G G A K R	0.4 %	16 %
	G K G G K G L G K G G A K R	0.4 %	16 %
3 acetylations	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
4	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %
	G K G G K G L G K G G A K R	0.3 %	32 %

obtainable from the Orbitrap mass spectrometer is noted for several of the major fragment ions. Interestingly, modification of the basic lysine residue in both cases produces fragmentation patterns that are quite equivalent, suggesting that their comparable ionization and fragmentation behavior may provide an opportunity to make some assessment of the degree of methylation. On the chromatographic time frame (Fig. 10), the retention times are similar, though not identical. Integration of the mass spectra across this range of retention time produces the composite spectrum shown in Fig. 11.

3.5. Methylation of histone H3 peptide KSAPSTGGVKKPHR

A tryptic fragment at m/z 533.6494 corresponds to the expected mass of the triply charged peptide ion from the yeast H3_{27–40} peptide KSAPSTGGVKKPHR carrying a methyl group and three deuterioacetylated lysines. The arginine residue and all three lysines are potential methylation sites. The collision induced dissociation (CID) MS/MS spectrum of the triply charged molecular ion is shown in Fig. 12. All of the observed b-series and y-series ions support methylation at either the K₃₆ or K₃₇ residues. The fragment ion that should distinguish these two possible structures is the singly charged y₄ ion, where the major peak at m/z 582.3545 (Fig. 12 inset) corresponds to methylation of the K₃₆ lysine. The mass accuracy of 0.8 ppm obtainable on the LTQ/Orbitrap mass spectrometer is

excellent and consistent with the mass accuracy in that range for other observed fragment ions. A smaller peak at m/z 596.3589 may also correspond to the y₄ ion of a structure in which the K₃₇ site is methylated. The mass accuracy in this case is 20 ppm, which suggests that this does not correspond to methylation at the K₃₇ site, or that the low signal/noise for this peak precludes obtaining a centroid with sufficient mass accuracy. In either case, methylation in this structure occurs primarily at K₃₆.

Like the previous example, methylated lysines are derivatized by the deuterioacetylation method, which in this case assists in the analysis by preserving these closely spaced lysine residues from cleavage by trypsin.

3.6. Trimethylation of histone H3 peptide KSAPSTGGVKKPHR

The deuterioacetylated histone H3_{27–40} peptide from HeLa cells also reveals a trimethylation site. The MS/MS spectrum of the triply charged molecular ion observed at m/z 527.9819 is shown in Fig. 13, and shows the same characteristic ions corresponding to b₂⁺, b₃⁺ and y₁₁ to y₁₃. These fragment masses are consistent with a structure having two deuterioacetylated lysine residues and a trimethylated lysine residue, and are accurate within 2–3 ppm. In contrast, errors in the mass accuracies when compared with a structure having two deuterioacetylated and one acety-




Fig. 9. Nanospray LTQ/Orbitrap MS/MS spectra of the doubly charged molecular ions of the derivatized peptide EIAQDFKTDLR (a) methylated at lysine 79 and (b) unmethylated. The different masses of the y_7 and y_9 ions distinguish the two species, both of which are deuterioacetylated at lysine 79. Mass accuracy is shown for the major fragment peaks.

lated lysine are of the order of 20–30 ppm, the approximate mass difference between an acetyl and a trimethyl modification. The masses of the b_2^{+1} and b_3^{+1} ions preclude trimethylation on the K_{27} residue, but no fragment ion was observed that would distinguish between trimethylation at the K_{36} and K_{37} sites. Based upon the observation of methylation at K_{36} in the prior exam-




Fig. 10. C18 HPLC single ion chromatograms of derivatized (a) unmethylated and (b) methylated EIAQDFKTDLR obtained by monitoring the doubly charged molecular ions. Retention times of these two species, while not the same, are similar.




Fig. 11. Quantitation of the methylated and unmethylated EIAQDFKTDLR based upon integration of spectra obtained across the retention times of the two species.




Fig. 12. Nanospray LTQ/Orbitrap mass spectrum of the molecular ion of the singly methylated and fully deuterioacetylated H3 peptide KSAPSTGGVKKPHR. Both the fragmentation pattern and mass accuracy are used to determine the location of the methylated lysine. The major y_4 fragment ion at 582.3545 differs 0.8 ppm from the calculated mass 582.3550 of isoform having methylation at the second lysine. A smaller peak at 596.3589 is close to the expected mass for the y_4 ion of the isoform having methylation at the third lysine, but has an error of 20 ppm.




Fig. 13. Nanospray LTQ/Orbitrap MS/MS spectrum of the deuterioacetylated peptide KSAPSTGGVKKPHR + 4Me, distinguished from an acetylated species by the accurate mass measurement.

ple then, the likely structure is KSAPSTGGVK_{3Me}KPHR (observed as K_{dAc}SAPSTGGVK_{3Me}K_{dAc}PHR).

Note that the use of deuterioacetylation can be used to distinguish between this structure and one in which the methylation is distributed, e.g., K_{Me}SAPSTGGVK_{Me}K_{Me}PH. In this case all three of the lysines would have been deuterioacetylated and would result in a much higher mass. Thus, even in the absence of high mass accuracy measurements, deuterioacetylation can in some case be used to distinguish between isomeric forms.

4. Conclusions

There is considerable interest in developing “global” mass spectrometry approaches to assess lysine acetylation, particularly as this type of modification now appears to play significant roles in many cellular processes and species. Other groups such as Zhao et al. [34] have used alternative methods to quantitate lysine acetylation. Their approach utilizes anti-acetyllysine antibodies to enrich for endogenously acetylated proteins and *stable isotope labeling with amino acids in cell culture* (SILAC) for quantitation by mass spectrometry. The sirtuin proteins have been shown to have deacetylase activity in mitochondria [35]. Another group, Kelleher et al. [36], report a method for global histone profiling in response to inhibition or knockdown of human deacetylases using a linear ion trap Fourier transform mass spectrometer. In this approach, histone mixtures are subjected to methionine oxidation prior to RPLC separation to enable resolution of each histone type (H1, H2.B, H2A-1, etc.). The high mass resolution and accuracy of the FTMS then enable one to determine the numbers of acetyl and methyl groups, and the method generally regards these to be the most abundant isoforms, e.g., H4 + 2Me occurs as H4K20, H4 + 2Me + 2Ac occurs as H4K20_{2Me}K12_{Ac}K16_{Ac}, etc. This approach has the advantage of assessing all of the possible modification types, provides some quantitation of the overall modifications, but does not use MS/MS to provide details of positional isomeric forms. Alternatively, top-down methods using electron capture dissociation on the FTMS [13] provide structural verification of the major isoforms, but are not necessarily as quantitative as isotope based methods.

Our interest in the bottom-up approach and chemical derivatization is based upon the possibility for exquisite and facile quantification of acetylation at specific residues, which was established in an earlier study by us of the acetylation of the K₅₆ site in histone H3 for a series of *hst3* and *hst4* mutants [20]. In that study, MALDI mass spectra of protein digests (as shown in Fig. 1) were obtained for histones derived from wild type, deletion mutants

and H184A, N152A and D154A mutants. Using this quantitative method for cells synchronized in G1 phase, we were then able to monitor changes in K₅₆ acetylation to observe the inactivation of Hst3/Hst4p during passage through S Phase in response to NIA [20; Supplementary Material]. Regulation of histone H3 K₅₆ has been shown to be critical in the cell cycle of fungi [37] and is regulated by the fungal histone acetyltransferase Rtt109 [38], a homolog of p300/CBP specific for K₅₆. In addition to the peptide containing K₅₆, expansion of the isotopic displays from the MALDI mass spectra enabled quantitation of the acetylation of lysine residues in K₉STGGK₁₄APR and K₁₈QLASK₂₃AAR in histone H3 [20; Supplementary Material].

In addition to obtaining quantitative information on acetylation at the positional isoform level, the intent is to begin to utilize this bottom-up derivatization method to provide qualitative and semiquantitative analyses for other modifications, particularly methylation, dimethylation and trimethylation. Methylation of the K₇₉ site on histone H3 is of particular importance in cell cycle and replication, and Kelleher et al. [39] have recently used a stable isotope strategy (SILAC) and mass spectrometry to compare the methylation and dimethylation of pre-existing (old) and newly synthesized histones. While it is clear that the deuterioacetylated structures: EIAQDFK_{Me+dAc}TDLR and EIAQDFK_{dAc}TDLR encompassing the K₇₉ site are not chemically and chromatographical identical, as in the case of stable isotope analogs, their similar responses to ionization and mass spectral sensitivity suggest that with the proper peptide analog experiment, analyses intended to observe changes in methylation at this site could be easily calibrated for accurate quantitation.

We conclude by noting that a bottom-up approach using global chemical derivatization of endogenously unmodified lysines is an effective alternative to top-down approaches that is advantageous when quantitative measurements are focused on specific modification sites, and that this approach can be extended to modifications other than acetylation.

Acknowledgments

The authors gratefully acknowledge collaborators Jef Boeke, Ivana Celic, Pamela Meluh and Junbiao Dai (Johns Hopkins) and Alain Verreault (University of Montreal) for providing and preparing mutant yeast histones, and Rocío Montes de Oca and Katherine L. Wilson (Johns Hopkins) for HeLa cell histones.

This project was supported by grant U54 RR020839 from the National Institutes of Health, contract NHLBI N01HV28180 from the National Heart Lung and Blood Institute, and by NIH grant R01GM48646 to K.L.W. The LTQ-Orbitrap mass spectrometer used in this study was purchased through grant S10 RR023025 from the NIH High End Instrumentation Program to R.J.C. Analyses were carried out at the Middle Atlantic Mass Spectrometry Laboratory.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ijms.2011.04.006.

References

- [1] K. Luger, A.W. Mäder, R.K. Richmond, D.F. Sargent, T.J. Richmond, Crystal structure of the nucleosome core particle at 2.8 Å resolution, *Nature* 389 (1997) 251–260.
- [2] P. Cheung, C.D. Allis, P. Sassone-Corsi, Signaling to chromatin through histone modifications, *Cell* 103 (2) (2000) 263–271.
- [3] F. Barlési, G. Giaccone, M.I. Gallegos-Ruiz, A. Loundou, S.W. Span, P. Lefesvre, F.A. Krutz, J.A. Rodriguez, Global histone modifications predict prognosis of resected non-small-cell lung cancer, *J. Clin. Oncol.* 25 (2007) 4358–4364.
- [4] C. Bönsch, S.M. Nieratschker, N.K. Orfanos, S.B. Hake, Chromatin proteomics and epigenetic regulatory circuits, *Expert Rev. Proteomics* 5 (2008) 105–119.

- [5] X. Su, C. Ren, M.A. Freitas, Mass spectrometry-based strategies for characterization of histones and their post-translational modifications, *Expert Rev. Proteomics* 4 (2007) 211–225.
- [6] A.L. Burlingame, X. Zhang, R.J. Chalkley, Mass spectrometric analysis of histone posttranslational modifications, *Methods* 36 (2005) 383–394.
- [7] J.V. Olsen, S.E. Ong, M. Mann, Trypsin cleaves exclusively C-terminal to arginine and lysine residues, *Mol. Cell. Proteomics* 3 (2004) 608–614.
- [8] B.A. Garcia, S. Mollah, B.M. Ueberheide, S.A. Busby, T.L. Muratore, J. Shabanowitz, D.F. Hunt, Chemical derivatization of histones for facilitated analysis by mass spectrometry, *Nat. Protoc.* 2 (4) (2007) 933–938.
- [9] N. Siuti, N.L. Kelleher, Decoding protein modifications using top-down mass spectrometry, *Nat. Methods* 4 (2007) 817–821.
- [10] B.A. Garcia, J. Shabanowitz, D.F. Hunt, Characterization of histones and their post-translational modifications by mass spectrometry, *Curr. Opin. Chem. Biol.* 11 (2007) 66–73.
- [11] L. Jiang, J.N. Smith, S.L. Anderson, P. Ma, C.A. Mizzen, N.L. Kelleher, Global assessment of combinatorial post-translational modification of core histones in yeast using contemporary mass spectrometry: lys4 trimethylation correlates with degree of acetylation on the same H3 tail, *J. Biol. Chem.* 282 (2007) 27923–27934.
- [12] M.T. Boyne II, J.J. Pesavento, C.A. Mizzen, N.L. Kelleher, Precise characterization of human histones in the H2A gene family by top-down mass spectrometry, *J. Proteome Res.* 5 (2006) 248–253.
- [13] N. Siuti, M.J. Roth, C.A. Mizzen, N.L. Kelleher, J.J. Pesavento, Gene-specific characterization of human histone H2B by electron capture dissociation, *J. Proteome Res.* 5 (2006) 233–239.
- [14] N. Young, P. DiMaggio, M. Plazas-Mayorca, R. Baliban, C. Floudas, B. Garcia, High throughput characterization of combinatorial histone codes, *Mol. Cell. Proteomics* 8 (10) (2009) 2266–2284.
- [15] J. Pesavento, Y. Kim, G. Taylor, N. Kelleher, Shotgun annotation of histone modifications: a new approach for streamlined characterization of proteins by top-down mass spectrometry, *J. Am. Chem. Soc.* 126 (11) (2004) 3386–3387.
- [16] C.M. Smith, P.R. Gafken, Z. Zhang, D.E. Gottschling, J.B. Smith, D.L. Smith, Mass spectrometric quantification of acetylation at specific lysines within the amino-terminal tail of histone H4, *Anal. Biochem.* 316 (2003) 23–33.
- [17] C.M. Smith, Quantification of acetylation at proximal lysine residues using isotopic labeling and tandem mass spectrometry, *Methods* 36 (2005) 395–403.
- [18] A.Y. Lee, C.P. Paweletz, R.M. Pollock, R.E. Settlege, J.C. Cruz, J.P. Secrist, T.A. Miller, M.G. Stanton, A.M. Kral, N.D. Ozerova, F. Meng, N.A. Yates, V. Richon, R.C. Hendrickson, Quantitative analysis of histone deacetylase-1 selective histone modifications by differential mass spectrometry, *J. Proteome Res.* 7 (12) (2008) 5177–5186.
- [19] H.C. Beck, E.C. Nielsen, R. Matthiesen, L.H. Jensen, M. Sehested, P. Finn, M. Grauslund, A.M. Hansen, O.N. Jensen, Quantitative proteomic analysis of post-translational modifications of human histones, *Mol. Cell. Proteomics* 5 (7) (2006) 1314–1325.
- [20] I. Celic, H. Masumoto, W.P. Griffith, P. Meluh, R.J. Cotter, J.D. Boeke, A. Verreault, The siruins Hst3 and Hst4p preserve genome integrity by controlling histone H3 lysine 56 deacetylation, *Curr. Biol.* 16 (2006) 1280–1289.
- [21] R.J. Cotter, W. Griffith, C. Jelinek, Tandem time-of-flight (TOF/TOF) mass spectrometry and the curved-field reflectron, *J. Chromatogr. B* 855 (2007) 2–13.
- [22] B.A. Garcia, S. Mollah, B.M. Ueberheide, S.A. Busby, T.L. Muratore, J. Shabanowitz, D.F. Hunt, Chemical derivatization of histones for facilitated analysis by mass spectrometry, *Nat. Protoc.* 2 (2007) 933–938.
- [23] M.D. Plazas-Mayorca, B.M. Zee, N.L. Young, I.M. Fingerman, G. LeRoy, S.D. Briggs, B.A. Garcia, One-pot shotgun quantitative mass spectrometry characterization of histones, *J. Proteome Res.* 8 (2009) 5367–5374.
- [24] B.A. Garcia, J.J. Pesavento, C.A. Mizzen, N.L. Kelleher, Pervasive combinatorial modification of histone H3 in human cells, *Nat. Methods* 4 (2007) 487–489.
- [25] A.H. Wang, N.R. Bertos, M. Vezmar, N. Pelletier, M. Crosato, H.H. Heng, J. Th'ng, J. Han, X. Yang, HDAC4, a human histone deacetylase related to yeast HDA1, is a transcriptional corepressor, *Mol. Cell. Biol.* 19 (1999) 7816–7827.
- [26] N.L. Young, P.A. DiMaggio, M.D. Plazas-Mayorca, R.C. Baliban, C.A. Floudas, B.A. Garcia, High throughput characterization of combinatorial histone codes, *Mol. Cell. Proteomics* 8 (2009) 2266–2284.
- [27] C.R. Jiménez, L. Huang, Y. Qiu, A.L. Burlingame, In-gel digestion of proteins for MALDI-MS fingerprint mapping, *Curr. Protocols Prot. Sci.* 16 (1998) 4.
- [28] S.L. Cohen, B.T. Chait, Influence of matrix solution conditions on the MALDI-MS analysis of peptides and proteins, *Anal. Chem.* 68 (1996) 31–37.
- [29] T.J. Cornish, R.J. Cotter, A curved-field reflectron for improved energy focusing of product ions in time-of-flight mass spectrometry, *Rapid Commun. Mass Spectrom.* 7 (1993) 1037–1040.
- [30] R.J. Cotter, B. Gardner, S. Iltchenko, R.D. English, Tandem time-of-flight mass spectrometry with a curved field reflectron, *Anal. Chem.* 76 (2004) 1976–1981.
- [31] B.A. Garcia, C.M. Barber, S.B. Hake, C. Ptak, F.B. Turner, S.A. Busby, J. Shabanowitz, R.G. Moran, C.D. Allis, D.F. Hunt, Modifications of human histone H3 variants during mitosis, *Biochemistry* 44 (2005) 3202–3213.
- [32] B. Sarg, W. Helliger, H. Talasz, E. Koutzamani, H. Lindner, Histone H4 hyperacetylation precludes histone H4 lysine 20 trimethylation, *J. Biol. Chem.* 279 (2004) 53458–53464.
- [33] N. Suka, Y. Suka, A. Carmen, J. Wu, M. Grunstein, Highly specific antibodies determine histone acetylation site usage in yeast heterochromatin and euchromatin, *Mol. Cell* 8 (2) (2004) 473–479.
- [34] J. Zhang, R. Sprung, J. Pei, X. Tan, S. Kim, H. Zhu, C.F. Liu, N.V. Grishin, Y. Zhao, Lysine acetylation is a highly abundant and evolutionarily conserved modification in *Escherichia coli*, *Mol. Cell. Proteomics* 8 (2) (2009) 215–225.
- [35] B. Schwer, J. Bunkenborg, R.O. Verdin, J.S. Andersen, E. Verdin, Reversible lysine acetylation controls the activity of the mitochondrial enzyme acetyl-CoA synthetase 2, *Proc. Natl. Acad. Sci. U. S. A.* 103 (27) (2006) 10224–10229.
- [36] M. Li, L. Jiang, N.L. Kelleher, Global histone profiling by LC-FTMS after inhibition and knockdown of deacetylases in human cells, *J. Chromatogr. B* 877 (2009) 3885–3892.
- [37] B. Xhemalce, K.M. Miller, R. Driscoll, H. Masumoto, S.P. Jackson, T. Kouzarides, A. Verreault, B. Arcangioli, Regulation of histone H3 lysine 56 acetylation in *Schizosaccharomyces pombe*, *J. Biol. Chem.* 282 (2007) 15040–15047.
- [38] Y. Tang, M.A. Holbert, H. Wurtele, K. Meeth, W. Rocha, M. Gharib, E. Jiang, P. Thibault, A. Verreault, P.A. Cole, R. Marmorstein, Fungal Rtt109 histone acetyltransferase is an unexpected structural homolog of p300/CBP, *Nat. Struct. Mol. Biol.* 15 (2008) 738–745.
- [39] S.M. Sweet, M. Li, P.M. Thomas, K.R. Durbin, N.L. Kelleher, Kinetics of re-establishing H3K79 methylation marks in global human chromatin, *J. Biol. Chem.* 285 (43) (2010), 32778–32786.