

Stationary Axisymmetric Solutions of the Einstein Equations with Rigidly Rotating Perfect Fluid and Nonlinear Charged Sources

Humberto Salazar¹ and Rubén Cordero²

Received September 15, 1995

A class of stationary, rigidly rotating perfect fluids coupled with nonlinear electromagnetic fields was investigated. An exact solution of the Einstein equations with sources for the Carter B(+) branch was found for the equation of state $3p + \epsilon = \text{const}$. We use a structure function for the Born-Infeld nonlinear electrodynamics which is invariant under duality rotations and a metric possessing a four-parameter group of motions. The solution is of Petrov type D and the eigenvectors of the electromagnetic field are aligned to the Debever-Penrose vectors.

1. BORN-INFELD NONLINEAR ELECTRODYNAMICS AND DUALITY ROTATIONS

The basic description of the dynamical equations of nonlinear electrodynamics within general relativity can be done in terms of the null tetrad formalism according to which the metric is given by

$$g = 2e^1 \otimes e^2 + 2e^3 \otimes e^4, \quad e^2 = \overline{e^1} \quad (1)$$

where the $e^a \in \Lambda^1$ fulfill the Cartan structure equations

$$de^a = e^b \wedge \Gamma_b^a = \Gamma_{bc}^a e^b \wedge e^c \quad (2)$$

and $\Gamma_b^a \in \Lambda^1$ satisfy the second structure equations

$$d\Gamma_b^a + \Gamma_s^a \wedge \Gamma_b^s = \frac{1}{2} R_{bcd}^a e^c \wedge e^d \quad (3)$$

The Riemann curvature components R_{bcd}^a may be replaced by the Weyl conformal tensor components, which are characterized by five complex curva-

¹FCFM, Benemérita Universidad Autónoma de Puebla, A.P. 1152, 72000 Puebla, México.

²Departamento de Física, CINVESTAV-IPN, A.P. 14-740, 07000 México, D.F., México.

ture coefficients $C^{(a)}$, and the components of the traceless Ricci tensor $C_{ab} = R_{ab} - \frac{1}{4}g_{ab}R$, where $R_{ab} = R_{abs}$ and $R = R_a^a$. Nonlinear theories of Born-Infeld (BI) type are theories with a Hamiltonian function \mathcal{H} depending on the invariants of the skew-symmetric tensor P_{ab} ,

$$P = \frac{1}{4}P^{ab}P_{ab}, \quad Q = \frac{1}{4}\check{P}^{ab}P_{ab}, \quad \check{P}^{ab} = -\frac{1}{2}\epsilon^{abcd}P_{cd}$$

where ϵ^{abcd} is the Levi-Civita symbol with $\epsilon^{1234} = 1$.

The skew-symmetric tensor F_{ab} can be defined by the material equations

$$F_{ab} = \mathcal{H}_P P_{ab} + \mathcal{H}_Q \check{P}_{ab}$$

$$\mathcal{H}_P = \frac{\partial \mathcal{H}}{\partial P}, \quad \mathcal{H}_Q = \frac{\partial \mathcal{H}}{\partial Q} \tag{4}$$

We select the null tetrad in such a manner that out of all independent components of the electromagnetic field tensors F_{ab} (corresponding to \mathbf{E} and \mathbf{B}) and P_{ab} (corresponding to \mathbf{D} and \mathbf{H}), only they are different from zero:

$$P_{34} = D, \quad P_{12} = iH, \quad F_{34} = E, \quad F_{12} = iB \tag{5}$$

where $D, H, E,$ and B are real. The above selection for F_{ab} and P_{ab} can be made simultaneously by virtue of the material equations. The invariants of F_{ab} and P_{ab} read

$$\frac{1}{4}P^{ab}P_{ab} + \frac{1}{4}\check{P}^{ab}P_{ab} = P + Q = -\frac{1}{2}(D + iH)^2 \neq 0$$

$$\frac{1}{4}F^{ab}F_{ab} + \frac{1}{4}\check{F}^{ab}F_{ab} = F + G = -\frac{1}{2}(E + iB)^2 \neq 0 \tag{6}$$

Thus (D, H) and (E, B) can be interpreted as independent parameters of the complex invariant of the electromagnetic field. We now can introduce by a Legendre transformation of the Hamiltonian $\mathcal{H}(P, Q)$ or $\mathcal{H}(D, H)$ a new structure function for the nonlinear electrodynamics given by (Salazar *et al.*, 1987)

$$M(D, B) = BH - \mathcal{H}(D, H) \tag{7}$$

In the following we use the same notation as Salazar *et al.* (1987).

Furthermore, we restrict our structure function M to be invariant under duality rotations, i.e.,

$$M(D', B') = M(D, B) \quad \text{for} \quad D' + iB' = e^{i\alpha}(D + iB) \tag{8}$$

The last condition can be easily seen to constrain the function M to be a function of the variable $(D^2 + B^2)$ only:

$$M = b^2f(X), \quad X \equiv \frac{1}{2b^2}(D^2 + B^2), \quad b = \text{const} \tag{9}$$

The original Born–Infeld theory given by the Hamiltonian function

$$\mathcal{H} = b^2 - \sqrt{b^4 - 2b^2P + Q^2} \tag{10}$$

belongs to this class of theories invariant under duality rotations and corresponds to

$$f(X) = \sqrt{1 + 2X} - 1 \tag{11}$$

2. EINSTEIN–BORN–INFELD EQUATIONS WITH PERFECT FLUIDS

In this work we are concerned with solutions to the Einstein–Born–Infeld equations with a perfect fluid

$$\begin{aligned} R_{ab} - \frac{1}{2}g_{ab}R &= -T_{ab} \\ T_{ab} &= (p + \epsilon)u_a u_b + pg_{ab} - 8\pi E_{ab} \\ u_a u^a &= -1 \\ p + \epsilon &> 0 \\ 4\pi E_{ab} &= \mathcal{H}_p(-P_{as}P_b^s + g_{ab}P) + (P\mathcal{H}_p + Q\mathcal{H}_Q - \mathcal{H})g_{ab} \\ \check{F}_{ab; a} &= 0 \\ P^{ba; a} &= 4\pi J^b \end{aligned} \tag{12}$$

where E_{ab} is the energy-momentum tensor of the nonlinear electromagnetic field, u_a is the fluid four-velocity, p is the fluid pressure, and ϵ is the energy density. We shall consider the Carter type D metric with conformal factor and having a four-parameter group of symmetries. We shall work with standard gravitational units so chosen that the gravitational constant G and the velocity of light c are equal to unity. Let now ϵ be a dimensionless constant, l be a constant of dimension of length, and $x^\mu := \xi, \bar{\xi}, r, \tau$ and $x^\mu := u, v, r, \sigma$ be two coordinate charts: ξ is complex, while the remaining coordinates and constants are real. The $r, \tau,$ and σ are of dimension of length, while $\xi, u,$ and v are dimensionless. Given these ingredients, we construct first the two-dimensional Riemannian spaces of constant curvature

$$\Lambda^1 \otimes \Lambda^1 \ni dl^{2(\pm)} := \begin{cases} 4 \frac{d\xi \otimes d\bar{\xi}}{(1 + \epsilon\xi\bar{\xi})^2} \\ 4 \frac{du \otimes dv}{(1 + \epsilon uv)^2} \end{cases}$$

given in terms of stereographic coordinates $(\xi, \bar{\xi})$ and the associated real one-forms of dimension of length

$$\Lambda^1 \ni \pi^\pm := \begin{cases} d\tau + 2il \left(\frac{\bar{\xi}d\xi - \xi d\bar{\xi}}{1 + \varepsilon\xi\bar{\xi}} \right) \\ d\sigma + 2l \left(\frac{vdu - u dv}{1 + \varepsilon\xi\bar{\xi}} \right) \end{cases}$$

Then, introducing two real analytic functions $N = N(r)$ and $F^{(\pm)} = F^{(\pm)}(r)$ (of dimension $[\text{length}]^2$), we define now the four-dimensional Riemannian space-time of signature $(+++ -)$

$$B^{(\pm)} : ds^2 := N dl^{2(\pm)} + \frac{N}{F^{(\pm)}} dr \otimes dr \mp \frac{F^{(\pm)}}{N} \pi^{(\pm)} \otimes \pi^{(\pm)} \quad (13)$$

It can be easily shown that the (Carter, 1968) separable $B^{(\pm)}$ branches of type D can be brought—without any loss of generality—to the form of (13). In the considered representation of the $B^{(\pm)}$ metrics, only the sign of the parameter ε is relevant.

We now explain why we consider (13) as the optimal coordinatization of the Carter $B^{(\pm)}$ branches for our purpose. We observe first that a formal transformation

$$z \rightarrow -u, \quad \bar{z} \rightarrow -v, \quad \tau \rightarrow i\sigma, \quad F^{(+)} \rightarrow F^{(-)} \quad (14)$$

which obviously implies $dl^{2(+)} \rightarrow dl^{2(-)}$ and $\pi^{(+)} \rightarrow i\pi^{(-)}$, brings the $B^{(+)}$ metric into the $B^{(-)}$ metric. We will see that this leads to a useful computational advantage, the natural tetrads; the connections and curvatures of $B^{(\pm)}$ metrics can be treated via perfectly parallel computations. Second, we notice that our coordinatization of the $B^{(\pm)}$ metrics allows us to give a unified description of their minimal subgroup of symmetries, which are four-dimensional Lie groups. Next, taking the $B^{(+)}$ metrics, we investigate the advantages of our coordinatization from the point of view of the Debney–Kerr–Schild formalism described in the first section. With the metrics (13) a natural choice for the null tetrads is correspondingly

$$-\frac{\sqrt{2N}}{1 + \varepsilon\xi\bar{\xi}} \begin{Bmatrix} d\xi \\ d\bar{\xi} \end{Bmatrix} = \begin{Bmatrix} e^1 \\ e^2 \end{Bmatrix}$$

$$\frac{1}{\sqrt{2}} \left[\left(\sqrt{\frac{N}{F}} \right) dr \pm \sqrt{\frac{F}{N}} \pi \right] = \begin{Bmatrix} e^3 \\ e^4 \end{Bmatrix}$$

whereas the connection forms are given by

$$\Gamma_{42} = \sqrt{\frac{F}{2N^3}} (il - \frac{1}{2}\dot{N})e^1 \tag{15}$$

$$\Gamma_{31} = \sqrt{\frac{F}{2N^3}} (il - \frac{1}{2}\dot{N})e^2 \tag{16}$$

where dots denote the r derivative, and

$$\Gamma_{12} + \Gamma_{34} = \frac{\epsilon}{\sqrt{2N^3}} [\xi e^2 - \bar{\xi} e^1] - \frac{1}{\sqrt{2}} \left[\left(\sqrt{\frac{F}{N}} \right)^* + il \sqrt{\frac{F}{N^3}} \right] (e^3 - e^4) \tag{17}$$

From the second structure equations

$$d\Gamma_{42} + \Gamma_{42} \wedge (\Gamma_{12} + \Gamma_{34}) = \gamma e^3 \wedge e^1 + \delta e^4 \wedge e^1 \tag{18}$$

$$d\Gamma_{31} + (\Gamma_{12} + \Gamma_{34}) \wedge \Gamma_{31} = \delta e^3 \wedge e^2 + \gamma e^4 \wedge e^2 \tag{19}$$

$$d(\Gamma_{12} + \Gamma_{34}) + 2\Gamma_{42} \wedge \Gamma_{31} = \beta e^1 \wedge e^2 + \alpha e^3 \wedge e^4 \tag{20}$$

where

$$\alpha = \left[\frac{1}{2} \left(\frac{F}{N} \right)^* + il \left(\frac{F}{N^2} \right) \right] \tag{21}$$

$$\beta = \frac{F}{N^3} (il - \frac{1}{2}\dot{N})^2 - \frac{\epsilon}{N} + 2il \sqrt{\frac{F}{N^3}} \left[\left(\sqrt{\frac{F}{N}} \right)^* + il \sqrt{\frac{F}{N^3}} \right] \tag{22}$$

$$\begin{aligned} \gamma = \frac{1}{2} \frac{F^{1/2}}{N} \left\{ \frac{2}{N^{1/2}} (il - \frac{1}{2}\dot{N}) \left[\left(\sqrt{\frac{F}{N}} \right)^* + il \sqrt{\frac{F}{N^3}} \right] \right. \\ \left. - \frac{F^{1/2}}{N} \left(il \frac{\dot{N}}{N} + \ddot{N} + \frac{1}{2} \frac{\dot{N}^2}{N} \right) \right\} \end{aligned} \tag{23}$$

$$\delta = -\frac{F}{8N\dot{N}} \left(\frac{4l^2 + \dot{N}^2}{N} \right)^* \tag{24}$$

we can obtain the Weyl coefficients $C^{(a)}$, the traceless Ricci tensor C_{ab} , and the scalar curvature R :

$$C^{(3)} = \frac{1}{3}(\alpha + \beta + 2\gamma) \tag{25}$$

$$R = 2(\alpha + \beta - 4\gamma) \tag{26}$$

$$C_{12} = \frac{1}{2}(\beta - \alpha) = -C_{34} \tag{27}$$

$$C_{33} = -2\delta = C_{44} \tag{28}$$

Other curvature components are equal to zero.

We shall consider a rigidly rotating perfect fluid in the comoving frame such that the four-velocity in tetrad and coordinate components reads

$$u_1 = 0 = u_2; \quad u_3 = -u_4 = 1/\sqrt{2} \quad (29)$$

$$u^\mu = \sqrt{\frac{N}{F}} \delta_r^\mu \quad (30)$$

Then the Einstein equations coupled with BI theories invariant under duality rotations and perfect fluid become (Salazar *et al.*, 1987)

$$R = -8b^2f(X) + 8b^2Xf^\nabla(X) + 3p - \epsilon \quad (31)$$

$$C_{12} = -2b^2Xf^\nabla(X) - \frac{1}{4}(p + \epsilon) \quad (32)$$

$$C_{33} = -\frac{1}{2}(p + \epsilon) \quad (33)$$

denoting the derivative of $f(X)$ with respect to X by superscript ∇ .

From the definition

$$J^\mu = \rho u^\mu \quad (34)$$

we can infer that X depends only on r

$$X = X(r) \quad (35)$$

and the Born–Infeld equations read

$$[N(B + iD)]' - 2ilf^\nabla(X)(B + iD) = 4\pi iNJ^\nabla \quad (36)$$

Solving equations (31)–(36), we arrive at the equations for the pressure p and the energy density ϵ :

$$p = 2b^2f(X) - \frac{\epsilon}{N} + \frac{F\dot{N}}{2N^2} - \frac{F}{4N^3}(4l^2 + \dot{N}^2) \quad (37)$$

$$\epsilon = -2b^2f(X) + \frac{\epsilon}{N} - \frac{F\dot{N}}{2N^2} + \frac{3F}{4N^3}(4l^2 + \dot{N}^2) - \frac{F\dot{N}}{N^2} \quad (38)$$

$$p + \epsilon = -\frac{F}{2N\dot{N}} \left(\frac{4l^2 + \dot{N}^2}{N} \right)' \neq 0 \quad (39)$$

and for the electromagnetic field

$$B = \frac{a_0 - 2lA}{N} b, \quad D = \frac{Ab}{f^\nabla(X)}, \quad A = A(r) \quad (40)$$

$$a_0, b = \text{const}$$

$$f^{\sigma}(X) = \dot{A} \left[2X - \left(\frac{a_0 - 2lA}{N} \right)^2 \right]^{1/2}, \quad X = X(A) \tag{41}$$

the remaining Einstein equation is

$$-2b^2 X f^{\sigma}(X) = \left[-2\varepsilon - \ddot{F} + \frac{2\dot{F}\dot{N}}{N} - \frac{F(4l^2 + \dot{N}^2)}{N^2} \right] / 4N \tag{42}$$

For every set of functions A, F, N satisfying equation (42) we have an analytic solution of the Einstein–Born–Infeld equations coupled with a perfect fluid. Energy density and pressure are given by (37) and (38) and the electromagnetic field is given by (40).

3. SOME EXPLICIT EXAMPLES

3.1. Solution Without Electromagnetic Field

When we switch off the electromagnetic field $X = 0, A = a_0/2l$ we recover from equation (42) the equation given by Kramer *et al.* (1980),

$$-2\varepsilon - \ddot{F} + 2 \frac{\dot{F}\dot{N}}{N} - F \frac{(4l^2 + \dot{N}^2)}{N^2} = 0 \tag{43}$$

3.2. Linear Electromagnetic Field

When

$$f(X) = X \tag{44}$$

equation (42) reduces to

$$(\dot{A})^2 + \frac{(a_0 - 2lA)^2}{N^2} + \frac{1}{4N} \left[-2\varepsilon - \ddot{F} + 2 \frac{\dot{F}\dot{N}}{N} - F \frac{(4l^2 + \dot{N}^2)}{N^2} \right] = 0 \tag{45}$$

This solution was reported by García and Tellez (1992).

3.3. Linear Electromagnetic Field Coupled with Dust

When we couple dust to the gravitational and linear electromagnetic field, characterized by the condition $p = 0$, then

$$F = \varepsilon r^2 + C_1 r + C_2, \quad C_1, C_2 = \text{const} \tag{46}$$

and equation (42) reduces to

$$\dot{A}^2 + \frac{(a_0 - 2lA)^2}{N^2} - \frac{1}{2N} \dot{F}\dot{N} + \frac{F}{4N^3} (4l^2 + \dot{N}^2) + \frac{\varepsilon}{N} = 0 \tag{47}$$

This solution is not reported in the literature.

3.4. Original Born–Infeld Theory

For the original BI theory described by $f(X) = (1 + 2X)^{1/2} - 1$, equation (42) reduces to

$$b^2 \frac{(\dot{A})^2 + (a_0 - 2IA)^2/N^2}{\{[1 - (\dot{A})^2][1 + (a_0 - 2IA)^2/N^2]\}^{1/2}} + \frac{1}{4N} \left[-2\epsilon - \dot{F} + \frac{2\dot{F}\dot{N}}{N} - \frac{F[4I^2 + (\dot{N})^2]}{N^2} \right] = 0 \tag{48}$$

This branch of solutions has been not reported in the literature, except for $A = \text{cte} \neq a_0/(2I)$ (Bretón, 1989).

3.5. Born–Infeld Theory with Perfect Fluid and Given Equation of State

From equations (37) and (38) we arrive at the general equation

$$(3p + \epsilon)' = 4b^2 f''(X) \frac{(XN^2)''}{N^2} - \frac{F}{N^2} \left[\dot{N} - \frac{4I^2 + (\dot{N})^2}{N} \right]' \tag{49}$$

with the ansatz

$$X = \frac{\epsilon_0^2 + g_0^2}{N^2}, \quad \epsilon_0, g_0 = \text{const} \tag{50}$$

and the equation of state

$$3p + \epsilon = \text{cte} = \gamma_0 \tag{51}$$

Equation (49) reduces to

$$\left[\dot{N} - \frac{4I^2 + \dot{N}^2}{N} \right]' = 0 \tag{52}$$

with general solution

$$N = \beta_0 + \left[\frac{4I^2}{\beta_1} + (\beta_0)^2 \right]^{1/2} \cosh \sqrt{\beta_1}(r - \beta_2) \\ \beta_0, \beta_1, \beta_2 = \text{const} \tag{53}$$

Then from the equation of state we obtain for F

$$F = \dot{N} \left(F_0 + \int^r \frac{N}{(\dot{N})^2} \{ [\gamma_0 - 4b^2 f(N^{-2})]N + 2\epsilon \} dy \right) \tag{54}$$

On the other hand, from

$$X = \frac{\epsilon_0^2 + g_0^2}{N^2} \quad (55)$$

we have two possibilities for the electromagnetic field $B + iD$:

(i) Magnetic solution

$$\begin{aligned} A &= \text{const} \neq \frac{a_0}{2l} \\ B &= \frac{g_0}{N} \\ D &= 0 \\ 4\pi J^r &= \frac{2lg_0}{N^2} f^r \left(\frac{g_0^2}{N^2} \right) \end{aligned} \quad (56)$$

(ii) Generalized NUT solution. For $A \neq \text{cte}$,

$$\begin{aligned} B + iD &= \frac{2b^2(\epsilon_0^2 + g_0^2)}{N} e^{i\phi} \\ \phi &= 2l \int^r \frac{1}{N} f^r \left(\frac{1}{N^2} \right) dy \\ J^r &= 0 \end{aligned} \quad (57)$$

Case (ii) generalizes the NUT type D solution of the Einstein–Born–Infeld equations (Salazar *et al.*, 1987) when we add a rigidly rotating perfect fluid.

4. CONCLUSIONS

We have obtained a wide set of solutions of the Einstein equations coupled with a rigidly rotating perfect fluid and linear and nonlinear electromagnetic field. One of them, given by equation (57), generalizes the NUT solution in the case of Born–Infeld and perfect fluid sources satisfying the equation of state $3p + \epsilon = \text{const}$.

ACKNOWLEDGMENTS

This work was supported in part by CONACyT.

REFERENCES

- Bretón, B. N. (1989). *Journal of Mathematical Physics*, **30**, 2607.
- Carter, B. (1968). *Communications in Mathematical Physics*, **10**, 280.
- García, D. A., and Tellez, J. (1992). *Journal of Mathematical Physics*, **33**, 2254.
- Kramer, D., *et al.* (1980). *Exact Solutions of Einstein's Field Equations*, Cambridge University Press, Cambridge.
- Salazar, H. I., García, D. A., and Plebanski, J. F. (1987). *Journal of Mathematical Physics*, **28**, 2171.