

Subdirect Decompositions of Lattice Effect Algebras

Zdenka Riečanová¹

Received March 13, 2003

We prove a theorem about subdirect decompositions of lattice effect algebras. Further, we show how, under these decompositions, blocks, sets of sharp elements and centers of those effects algebras are decomposed. As an application we prove a statement about the existence of subadditive state on some block-finite effect algebras.

KEY WORDS: effect algebras; central elements; blocks; states; subdirect decompositions.

1. INTRODUCTION AND BASIC NOTIONS

In general, an effect algebra is a partial algebra with two constants 0, 1 and a partial binary operation \oplus (the orthogonal sum) satisfying very simply axioms introduced by Foulis and Bennett (1994). A model for an effect algebra is the standard effect algebra $\mathcal{E}(H)$ of positive self-adjoint operators dominated by the identity on the Hilbert space H .

Definition 1.1. (Foulis and Bennett, 1994). A structure $(E; \oplus, 0, 1)$ is called an *effect algebra* if 0, 1 are two distinguished elements and \oplus is a partially defined binary operation on E which satisfies the following conditions for any $a, b, c, \in E$:

- (Ei) $b \oplus a = a \oplus b$ if $a \oplus b$ is defined,
- (Eii) $(a \oplus b) \oplus c = a \oplus (b \oplus c)$ if one side is defined,
- (Eiii) for every $a \in E$ there exists a unique $b \in E$ such that $a \oplus b = 1$ (we put $a' = b$),
- (Eiv) if $1 \oplus a$ is defined then $a = 0$.

We often denote the effect algebra $(E; \oplus, 0, 1)$ briefly by E . In every effect algebra E we can define the partial operation \ominus and the partial order \leq by putting

$$a \leq b \text{ and } b \ominus a = c \text{ iff } a \oplus c \text{ is defined and } a \oplus c = b.$$

¹Department of Mathematics, Faculty of Electrical Engineering and Information Technology, Slovak Technical University, Ivkovičova 3, 812 19 Bratislava, Slovak Republic; e-mail: zriecan@elf.stuba.sk.

Since $a \oplus c = a \oplus d$ implies $c = d$, the \ominus and the \leq are well defined. If E with the defined partial order is a lattice (a complete lattice) then $(E; \oplus, 0, 1)$ is called a *lattice effect algebra* (a *complete effect algebra*). For more details we refer the reader to Dvurečenskij and Pulmannová (2000) and the references given there.

Definition 1.2. Let $(E; \oplus, 0, 1)$ be an effect algebra. $Q \subseteq E$ is called a subeffect algebra of E iff

- (i) $1 \in Q$,
- (ii) if $a, b, c \in E$ with $a \oplus b = c$ and out of a, b, c at least two elements are in Q then $a, b, c \in Q$.

Note that if Q is a subeffect algebra of E then Q with inherited operation \oplus is an effect algebra in its own right.

Definition 1.3. Let $(E; \oplus_E, 0_E, 1_E)$ and $(F; \oplus_F, 0_F, 1_F)$ be effect algebras. A bijective map $\varphi: E \rightarrow F$ is called an *isomorphism* if

- (i) $\varphi(1_E) = 1_F$,
- (ii) for all $a, b \in E: a \leq_E b'$ iff $\varphi(a) \leq_F (\varphi(b))'$ in which case $\varphi(a \oplus_E b) = \varphi(a) \oplus_F \varphi(b)$.

We write $E \cong F$. Sometimes we identify E with $F = \varphi(E)$. If $\varphi: E \rightarrow F$ is an injection with properties (i) and (ii) then φ is called an *embedding*.

Note, that Kôpka (1992) introduced to effect algebras equivalent in some sense structures called D -posets, in which the operation of difference of fuzzy sets is the primary operation. For the connections we refer to Dvurečenskij and Pulmannová (2000) Kôpka (1992), and Riečanová (1999).

Finally, note that lattice effect algebras generalize orthomodular lattices and MV-algebras, (including Boolean algebras) (Dvurečenskij and Pulmannová, 2000; Hájek, 1998; Höhle and Klement, 1995; Kôpka and Chovanec, 1995).

For every central element z of a lattice effect algebra E , the interval $[0, z]$ with the \oplus operation inherited from E and the new unity z is a lattice effect algebra in its own right. We are going to prove a statement about decompositions of E into subdirect products of such intervals $[0, z]$.

2. COMPATIBILITY, BLOCKS AND CENTRAL ELEMENTS

Recall that elements a, b of a lattice effect algebra $(E; \oplus, 0, 1)$ are called *compatible* (written $a \leftrightarrow b$) iff $a \vee b = a \oplus (b \ominus (a \wedge b))$ (see Kôpka and Chovanec, 1995). $P \subseteq E$ is a *set of pairwise compatible elements* if $a \leftrightarrow b$ for all $a, b \in P$. For $a \in E$ and $Q \subseteq E$ we will write $a \leftrightarrow Q$ iff $a \leftrightarrow q$ for all $q \in Q$. $M \subseteq E$ is called a *block* of E iff M is a maximal subset of pairwise compatible elements. Every block of a lattice effect algebra E is a subeffect algebra and a sub-lattice of

E and E is a union of its blocks (Riečanová, 2000a). Lattice effect algebra with a unique block is called an *MV-effect algebra*. Every block of a lattice effect algebra is an MV-effect algebra in its own rights. In Riečanová (2000a) it was proved that every block M of a lattice effect algebra E is closed with respect to all existing infima and suprema of subsets of M . We say than M is a *full sub-lattice* of E .

An element z of an effect algebra E is called central if $x = (x \wedge z) \vee (x \wedge z')$ for all $x \in E$. The center $C(E)$ of E is the set of all central elements of E , (Greechie *et al.*, 1995). If E is a lattice effect algebra then $z \in E$ is central if $z \wedge z' = 0$ and $z \leftrightarrow x$ for all $x \in E$, Riečanová (1999). Thus, in a lattice effect algebra E , $C(E) = B(E) \cap S(E)$, where $B(E) = \cap\{M \subseteq E \mid M \text{ is a block of } E\}$ is called a *compatibility center* of E and $S(E) = \{z \in E \mid z \wedge z' = 0\}$ is the set of all *sharp* elements of E . Evidently, $B(E) = \{x \in E \mid x \leftrightarrow y \text{ for all } y \in E\}$. In every lattice effect algebra E , $S(E)$ is an orthomodular lattice (Jenča and Riečanová, 1999) and $B(E)$, is an MV-effect algebra. Hence $C(E)$ is a Boolean algebra (Greechie *et al.*, 1995). Moreover, $S(E)$, $B(E)$, and $C(E)$ are full sub-lattices of a lattice effect algebra E , which means that they are closed with respect to all infima and suprema existing in E (Riečanová, 2001b). It follows that if E is a complete effect algebra then $S(E)$, $B(E)$, $C(E)$, and every block of E are also complete. Further, $B(E)$, $S(E)$, and $C(E)$ are subeffect algebras of E (Jenča and Riečanová, 1999; Riečanová, 2001b).

*In every lattice effect algebra E , if $z \in C(E)$ then the interval $[0, z]$ is a lattice effect algebra with the new unit z and the partial operation \oplus inherited from E . It is because for $x, y \leq z$ with $x \oplus y$ defined in E we have $x \oplus y \leq z$ (Greechie *et al.*, 1995). Further, $z = (x \oplus x') \wedge z = (x \oplus z) \oplus (x' \oplus z)$ for all $x \in E$. It follows that for $y \leq z$ we have $z = y \oplus (y' \wedge z)$ and hence $z \ominus y = y' \wedge z$ (Riečanová, 2003b).*

Our observations about subdirect products of lattice effect algebras are based on the following Theorem 2.1 from Jenča and Riečanová (1999), reprinted also in the book by Dvurečenskij and Pulmannová (2000, p. 107).

Lemma 2.1. (Jenča and Riečanová, 1999). *Let E be a lattice effect algebra. Assume $b \in E$ and $A \subseteq E$ are such that $\bigvee A$ exists in E and $b \leftrightarrow a$ for all $a \in A$. Then*

- (i) $b \leftrightarrow \bigvee A$
- (ii) $\bigvee\{b \wedge a \mid a \in A\}$ exists and equals $b \wedge (\bigvee A)$.

Another useful statements will be about central elements.

Lemma 2.2. (Riečanová, 2003b). *Let u, v with $u \leq v'$ be elements of a lattice effect algebra E and $z \in C(E)$, then $(u \oplus v) \wedge z = (u \wedge z) \oplus (u \wedge z)$.*

Lemma 2.3. *Let E be a lattice effect algebra and $D \subseteq C(E)$ with (1) $\bigvee D = 1$ and (2) $d_1 \wedge d_2 = 0$ for all $d_1 \neq d_2, d_1, d_2 \in D$. Then the following conditions are equivalent for $u, v \in E$:*

- (i) $u \leftrightarrow v$,
- (ii) $\forall d \in D: u \wedge d \leftrightarrow u \wedge d$,
- (iii) $\forall d_1, d_2 \in D: u \wedge d_1 \leftrightarrow u \wedge d_2$.

Proof: (i) \implies (ii) Because $D \leftrightarrow E$. Further (ii) \implies (iii) as for every $d_1 \neq d_2, d_1, d_2 \in D$ we have $d_1 \leq d_2'$ which gives $u \wedge d_1 \leq d_1 \leq d_2' \leq d_2' \vee u' = (u \wedge d_2)'$ and hence $u \wedge d_1 \leftrightarrow v \wedge d_2$. By Lemma 2.1 we obtain (iii) \implies (i), as for each $d_1 \in D$ we have $v \wedge d_1 \leftrightarrow u = \bigvee \{u \wedge d \mid d \in D\}$ and hence $v = \{\bigvee \{v \wedge d \mid d \in D\} \leftrightarrow u$. \square

Lemma 2.4. *Let E be a lattice effect algebra and $d \in C(E)$. Then*

- (i) if M is a block of E then $M \cap [0, d] = \{x \wedge d \mid x \in M\} \leftrightarrow u$.
- (ii) elements $u, v \in [0, d]$ are compatible in $[0, d]$ iff $u \leftrightarrow v$ in E .
- (iii) $M_d \subseteq [0, d]$ is block od $[0, d]$ iff there exists a block M of E with $M \cap [0, d] = M_d$.
- (iv) $S([0, d]) = \{x \wedge d \mid x \in S(E)\} = S(E) \cap [0, d]$.
- (v) $B([0, d]) = \{x \wedge d \mid x \in B(E)\} = B(E) \cap [0, d]$.
- (vi) $C([0, d]) = \{x \wedge d \mid x \in C(E)\} = C(E) \cap [0, d]$.

Proof:

- (i) Let $x \in M$. Because $d \leftrightarrow E$ and $x \leftrightarrow M$ we obtain $x \wedge d \leftrightarrow M$, which, by the maximality of blocks, gives $x \wedge d \in M$. Conversely, $y \in M \cap [0, d]$ implies $y = y \wedge d \in M$.
- (ii) This follows from the fact that $[0, d]$ is a sublattice of E which is closed with respect to \oplus .
- (iii) If M_d is a block of $[0, d]$ then, by (ii) M_d is a pairwise compatible set in E and hence by (Riečanová, 2000a) there exists a block M of E such that $M_d \subseteq M$. Hence $M_d \subseteq M \cap [0, d]$, which, by (i) and (ii) and the maximality of blocks, gives $M_d = M \cap [0, d]$. Conversely, if M is a block of $M \cap [0, d]$ is a pairwise compatible set in $[0, d]$, by (i) and Lemma 2.3. Further, if $y \in [0, d]$ such that $y \leftrightarrow M \cap [0, d]$ then $y = y \wedge d \leftrightarrow x \wedge d$ for all $x \in M$ by (i). Further, for all $x \in M$ and $d_1 \neq d, d_1 \in D$ we have $y \leq d \leq d_1' \leq d_1' \vee x' = (d_1 \wedge x)'$ and hence $y \leftrightarrow d_1 \wedge x$. By Lemma 2.1 we obtain $y \leftrightarrow x$ for all $x \in M$. By the maximality of blocks we obtain $y \in M$, hence $y \in M \cap [0, d]$. We conclude that $M \cap [0, d]$ is a block od $[0, d]$.

- (iv) If $x \in S(E) \cap [0, d]$ then $x \wedge x' = 0$ and $x \leq d$ which gives $(x \wedge d) \wedge (x' \wedge d) \leq x \wedge x' = 0$ and hence $x = x \wedge d \in S[0, d]$. If $x \in S[0, d]$ then $x = x \wedge d$ and $x \wedge x' = (x \wedge d) \wedge x' = x \wedge (x' \wedge d) = 0$ which gives $x \in S(E) \cap [0, d]$.
- (v) $B(E) \cap [0, d] = \bigcap \{M \cap [0, d] \mid M \text{ is a block of } E\} = \bigcap \{M_d \mid M_d \text{ is a block of } [0, d]\} = B([0, d])$. Moreover, $x \in B(E) \cap [0, d]$ implies $x = x \wedge d$. Conversely, for $x \in B(E)$ we have $x \wedge d \leftrightarrow E$ and hence $x \wedge d \in B([0, d])$.
- (vi) $C(E) \cap [0, d] = B(E) \cap S(E) \cap [0, d] = B([0, d]) \cap S([0, d])$, by (iv) and (v). Hence $C([0, d]) = C(E) \cap [0, d]$. Moreover, $x \in C(E) \cap [0, d]$ implies $x = x \wedge d$. Conversely, $x \in C(E)$ implies $x \wedge d \leftrightarrow E$ and $(x \wedge d) \wedge (x' \wedge d) = 0$ which gives $x \wedge d \in C([0, d])$.

3. SUBDIRECT DECOMPOSITIONS OF LATTICE EFFECT ALGEBRAS

A *direct product* of a family of effect algebras $\{E_x \mid x \in H\}$, $H \neq \emptyset$ is the Cartesian product $\prod \{E_x \mid x \in H\}$ with componentwise defined operations $\hat{\oplus}, \hat{0}, \hat{1}$, which means that $(a_x)_{x \in H} \hat{\oplus} (b_x)_{x \in H} = (a_x \oplus_x b_x)_{x \in H}$ iff $a_x \oplus_x b_x$ is defined in E_x for all $x \in H$. Further, $\hat{0} = (0_x)_{x \in H}$ and $\hat{1} = (1_x)_{x \in H}$. Evidently, for each $x_i \in H$, the natural projection pr_{x_i} of $\prod \{E_x \mid x \in H\}$ onto E_{x_i} , is a homomorphism. $\prod \{E_x \mid x \in H\}$ is a direct product decomposition of an effect algebra E if there is an isomorphism $\varphi : E \rightarrow \prod \{E_x \mid x \in H\}$.

A *subdirected product* of a family $\{E_x \mid x \in H\}$ of effect algebras, $H \neq \emptyset$ is a subeffect algebra Q of the direct product $\prod \{E_x \mid x \in H\}$ such that each restriction $pr_{x_i|_Q}$, $x_i \in H$, of the natural projection pr_{x_i} to Q is onto E_{x_i} . Q is a *subdirect product decomposition* of an effect algebra E if there exists an isomorphism $\varphi : E \rightarrow Q$.

Theorem 3.1. *Let $(E; \oplus, 0, 1)$ be a lattice effect algebra and $D \subseteq C(E)$ with (1) $\bigvee D = 1$ and (2) $d_1 \wedge d_2 = 0$ for all $d_1 \neq d_2; d_1, d_2 \in D$. Then:*

- (i) E is isomorphic to a subdirect product of the family of effect algebras $\{[0, d] \mid d \in D\}$.
- (ii) Up to isomorphism, E is a sublattice of $\hat{E} = \prod \{[0, d] \mid d \in D\}$.
- (iii) If, moreover, E is complete or D is finite then $E \cong \hat{E}$.

Proof:

- (i) By Lemma 2.1, for every $u \in E$ we have $u = u \wedge (\bigvee D) = \bigvee \{u \wedge d \mid d \in D\}$. Let $Q = \{(u \wedge d)_{d \in D} \mid u \in E\}$ and the map $\varphi : E \rightarrow Q$ be defined by the formula

$$\varphi(u) = (u \wedge d)_{d \in D}, \quad \text{for all } u \in E.$$

Let us show that Q is a subeffect algebra of $\hat{E} = \prod\{[0, d] \mid d \in D\}$ and φ is an isomorphism. Because operations $\hat{\oplus}$, $\hat{0}$ and $\hat{1}$ in \hat{E} are defined componentwise, we obtain that for every $x = (v \wedge d)_{d \in D} \in Q$ we have $x' = (u' \wedge d)_{d \in D} \in Q$ and $x \hat{\oplus} x' = ((u \wedge d) \oplus (u' \wedge d))_{d \in D} = ((u \oplus u') \wedge d)_{d \in D} = (1 \wedge d)_{d \in D} = \hat{1} \in Q$. Thus $\varphi(u') = (\varphi(u))'$. Because the partial order in \hat{E} is derived from $\hat{\oplus}$, hence it is also defined componentwise, we have for $y = (v \wedge d)_{d \in D} \in Q$ that $y \leq x'$ iff $v \leq u'$ iff $\varphi(v) \leq (\varphi(u))'$ and $\varphi(u) \hat{\oplus} \varphi(v) = x \hat{\oplus} y = (u \wedge d)_{d \in D} \hat{\oplus} (v \wedge d)_{d \in D} = ((u \wedge d) \oplus (v \wedge d))_{d \in D} = ((u \oplus v) \wedge d)_{d \in D} = \varphi(u \oplus v) \in Q$. Further, we can easily see that for $u, v \in E$ we have $\varphi(u) = \varphi(v)$ iff $u \wedge d = v \wedge d$ for all $d \in D$ iff $u = v$ by Lemma 2.1. Moreover, the restriction $pr_d|_Q$ of the natural projection pr_d to Q is onto $[0, d]$, hence Q is a subdirect product of \hat{E} .

- (ii) Because, up to isomorphism $[0, d] \subseteq E \subseteq \hat{E}$, for every $x = (u \wedge d)_{d \in D} \in Q$ we have $\bigvee\{u \wedge d \mid d \in D\} = u \in E$, we can conclude that, up to isomorphism E is a sublattice of \hat{E} , by Lemma 2.1.
- (iii) If D is a finite set or E is a complete lattice then for every $(u_d)_{d \in D} \in \hat{E}$ we have $\bigvee\{u_d \mid d \in D\} = u \in E$.

Corollary 3.2. *Under the assumptions of Theorem 1, if, moreover E is complete or D is finite then*

- (i) M is a block of E iff $M \cong \prod\{M_d \mid d \in D, M_d \text{ is a block of } E\} = \prod\{M \cap [0, d] \mid d \in D\}$.
- (ii) $S(E) \cong \prod\{S([0, d]) \mid d \in D\}$.
- (iii) $B(E) \cong \prod\{B([0, d]) \mid d \in D\}$.
- (iv) $C(E) \cong \prod\{C([0, d]) \mid d \in D\}$.

Proof: We have shown in the proof of Theorem 3.1 that the map $\varphi : E \rightarrow \prod\{[0, d] \mid d \in D\}$ defined by $\varphi(u) = (u \wedge d)_{d \in D}$, for all $u \in E$, is an isomorphism.

- (i) The statement follows by Lemmas 1 and 2 and the maximality of blocks, under which, the restriction $\varphi|_M$ is the wanted isomorphism.
- (ii) By Lemma 2.4, (iv), the restriction $\varphi|_{S(E)}$ is the wanted isomorphism.
- (iii) By Lemma 2.4, (v), the restriction $\varphi|_{B(E)}$ is the wanted isomorphism.
- (iv) By Lemma 2.4, (vi), the restriction $\varphi|_{C(E)}$ is the wanted isomorphism.

In the next theorem and subsequently we use the following notation: For an arbitrary mapping $\varphi : X \rightarrow Y$ and $A \subseteq Y$, the *inverse image* of A denoted by $\varphi^{-1}(A)$ is the set

$$\varphi^{-1}(A) = \{x \in X \mid \varphi(x) \in A\} \subseteq X.$$

Clearly, if φ is a bijection this is identical with the image of A by the inverse mapping φ^{-1} .

Theorem 3.3. *Under the assumptions of Theorem 3.1, if φ_D is the embedding of E into $\prod\{[0, d] \mid d \in D\}$ defined for all $x \in E$ by $\varphi_D(x) = (x \wedge d)_{d \in D}$, then*

- (i) M is a block of E iff there are blocks M_d of $[0, d]$, $d \in D$ such that $M = \varphi_D^{-1}(\prod\{M_d \mid d \in D\})$.
- (ii) $S(E) = \varphi_D^{-1}(\prod\{S([0, d]) \mid d \in D\})$,
- (iii) $B(E) = \varphi_D^{-1}(\prod\{B([0, d]) \mid d \in D\})$,
- (iv) $C(E) = \varphi_D^{-1}(\prod\{C([0, d]) \mid d \in D\})$.

Proof:

- (i) By Lemmas 2.3 and 2.4, M is a maximal pairwise compatible set of elements of E_{iff} for each $d \in D$, $\{u \wedge d \mid u \in M\}$ is a maximal pairwise compatible set in $[0, d]$. Thus the statement follows by the definition of φ_D and blocks.
- (ii)–(iv) These follow by Lemma 2.4 and the definition of φ_D .

Note that Theorem 3.3 and Lemma 2.4 actually show that every block M of E can be subdirectly decomposed by family $\{M_d \mid d \in D\}$ of blocks of $[0, d]$. Similar statements are true for the set of all sharp elements of E , compatibility center of E and center of E .

It is known that on every MV-algebra there exists a state. Moreover, every state on an MV-algebra is subadditive (Höhle and Klement, 1995). Unfortunately, there are even finite (lattice) effect algebras and orthomodular lattices admitting no states (Riečanová, 2001c and Greechie, 1971). Some positive results were given e.g. in Riečanová (2002). Further, positive results we can obtain as applications of decompositions of effect algebras.

Definition 3.4. A map $\omega : E \rightarrow [0, 1] \subseteq R$ is called a *state on the effect algebra* E if (i) $\omega(1) = 1$ and (ii) if $a, b \in E$ with $a \leq b'$ then $\omega(a \oplus b) = \omega(a) + \omega(b)$. If, moreover, E is lattice ordered then ω is called *subadditive* if $\omega(a \vee b) \leq \omega(a) + \omega(b)$, for all $a, b \in E$.

Note that a state on a lattice effect algebra need not be subadditive, (Riečanová, 2002).

Lemma 3.5. *Let E be a lattice effect algebra. A (subadditive) state on E exists iff there is a nonzero $d \in C(E)$ such that there exists a (subadditive) state on $[0, d]$.*

Proof: If $d \in C(E)$, $d \neq 0$ then for $x, y \in E$ with $x \leq y'$ we have $(x \oplus y) \wedge d = (x \wedge d) \oplus (y \wedge d)$ by Lemma 2.4. Moreover, for every $u, v \in E$, by Lemma 2.3 we

have $(u \vee v) \wedge d = (u \wedge d) \vee (v \wedge d)$. It follows that if $m : [0, d] \rightarrow [0, 1] \subseteq R$ is a (subadditive) state on $[0, d]$ then $\omega : E \rightarrow [0, 1] \subseteq R$ defined for all $x \in E$ by $\omega(x) = m(x \wedge d)$ is a (subadditive) state on E . The converse is clear, as $1 \in C(E)$.

Theorem 3.6. *Let E be a lattice effect algebra with exactly n blocks and the center $C(E) \neq \{0, 1\}$. If n is a prime number then:*

- (i) *For every $d \in C(E) \setminus \{0, 1\}$, at least one of the effect algebras $[0, d]$ and $[0, d']$ is an MV-effect algebra.*
- (ii) *There exists a subadditive state on E .*

Proof:

- (i) Let $d \in C(E) \setminus \{0, 1\}$. By Theorem 3.1, $E \cong [0, d] \times [0, d']$. Moreover, by Corollary 3.2, (i) we have $n = k_1 \cdot k_2$ under which k_1 and k_2 are numbers of blocks of $[0, d]$ and $[0, d']$ respectively. As n is a prime number, we conclude that at least one of k_1 and k_2 equals to 1.
- (ii) Assume that $[0, d]$ is an MV-effect algebra. Then there exists a subadditive state m on $[0, d]$, as $[0, d]$ can be organized into an MV-algebra. By Lemma 3.5, there exists a subadditive state on E .

ACKNOWLEDGMENT

This research is supported by grant G-1/0266/03 of the MŠ SR, Slovakia.

REFERENCES

- Dvurečenskij, A. and Pulmannová, S. (2000). *New Trends in Quantum Structures*, Kluwer, Dordrecht.
- Foulis, D. and Bennett, M. K. (1994). Effect algebras and unsharp quantum logics. *Foundations of Physics*, **24**, 1331–1352.
- Grätzer, G. (1998). *General Lattice Theory*, 2nd edn., Birkhäuser Verlag, Basel.
- Greechie, R. J. (1971). Orthomodular lattices admitting no states. *Journal of Combinatorial Theory A*, **10**, 119–132.
- Greechie, R. J., Foulis, D., and Pulmannová, S. (1995). The center of an effect algebra. *Order*, **12**, 91–106.
- Hájek, P. (1998). *Mathematics of Fuzzy Logic*, Kluwer, Dordrecht.
- Höhle, U. and Klement, E. P. (eds.) (1995). *Non-Classical Logics and Their Applications to Fuzzy Subsets*, Kluwer, Dordrecht.
- Jenča, G. and Riečanová, Z. (1999). On sharp elements in lattice ordered effect algebras. *BUSEFAL* **80**, 24–29.
- Kôpka, F. (1992). D-posets of fuzzy sets. *Tatra Mountains Mathematical Publications*, **1**, 83–87.
- Kôpka, F. and Chovanec, F. (1995). Boolean D-posets. *International Journal of Theoretical Physics*, **34**, 1297–1302.
- Riečanová, Z. (1999). Subalgebras, intervals and central elements of generalized effect algebras. *International Journal of Theoretical Physics*, **38**, 3209–3220.

- Riečanová, Z. (2000a). Generalization of blocks for D-lattices and lattice ordered effect algebras, *International Journal of Theoretical Physics*, **39**, 231–237.
- Riečanová, Z. (2000b). Archimedean and block-finite effect algebras. *Demonstratio Mathematica*, **33**, 443–452.
- Riečanová, Z. (2001b). Orthogonal sets in effect algebras. *Demonstratio Mathematica* **34**, 525–532.
- Riečanová, Z. (2001c). Proper effect algebras admitting no states. *International Journal of Theoretical Physics*, **40**, 1683–1691.
- Riečanová, Z. (2002). Smearings of states defined on sharp elements onto effect algebras. *International Journal of Theoretical Physics*, **41**, 1511–1524.
- Riečanová, Z. (2003a). Distributive atomic effect algebras, *Demonstratio Mathematica*, **36**, 247–259.
- Riečanová, Z. (2003b). Continuous lattice effect algebras admitting order continuous states. *Fuzzy Sets and Systems*, **136**, 41–54.