

Dissipation of System and Atom in Two-Photon Jaynes–Cummings Model with Degenerate Atomic Levels

Yan-Qing Guo,^{1,2} Ling Zhou,¹ and He-Shan Song¹

Received December 5, 2004; accepted March 30, 2005

A method of perturbative expansion of master equation is employed to study the dissipative properties of system and of atom in the two-photon Jaynes–Cummings model (JCM) with degenerate atomic levels. The numerical results show that the degeneracy of atomic levels prolongs the period of entanglement between the atom and the field. The asymptotic value of atomic linear entropy is apparently increased by the degeneration. The amplitude of local entanglement and disentanglement is suppressed. The better the initial coherence property of the degenerate atom, the larger the coherence loss.

KEY WORDS: Jaynes–Cummings model; degenerate atomic level; master equation.

1. INTRODUCTION

A main obstacle of quantum information is the decoherence of qubits. A quantum system inevitably interacts with its surrounding environment, and the interaction between the quantum system and its surroundings could lead to decoherence. The theory of environment dissipation induced decoherence is addressed not only in quantum information field (Davidovich *et al.*, 1996; Duan and Guo, 1998; Peixoto and Nemes, 1999; Zurek, 1991) but also in other fields, like brain process (Hagan, 2000; Tegmark, 2000).

As a candidate for quantum information, cavity QED has been widely studied in following aspects: quantum network (Cirac *et al.*, 1996; Pellizzari, 1997), producing entanglement (Cirac and Zoller, 1994a; Gerry, 1996; Zheng and Guo, 2001) and quantum teleportation (Cirac and Parkins, 1994; Davidovich *et al.*, 1994; Zheng and Guo, 2001), especially in studying the dissipative dynamics of a quantum system (Brune *et al.*, 1996; Davidovich *et al.*, 1996; Duan and Guo, 1998; Peixoto and Nemesbut, 1999). This is because the dissipation of cavity QED could

¹Department of Physics, Dalian University of Technology, Liaoning Dalian 116024, P.R. China.

²To whom correspondence should be addressed at Department of Physics, Dalian University of Technology, Liaoning Dalian 116024, P.R. China; e-mail: yqguohtp@yahoo.com.cn.

be monitored in experiment (Brune *et al.*, 1996; Davidovich *et al.*, 1996; Duan and Guo, 1998), and this is valuable not only to the candidate of quantum information but also to the research of decoherence theory. In Ref. (Peixoto and Nemes, 1999), the dissipative dynamics of the JCM in the dispersive approximation has been studied, and the influence of dissipation on the entanglement of the two subsystems was investigated. In Ref. (Zhou *et al.*, 2001), the authors studied the field state dissipative dynamics of two-photon Jaynes–Cummings Model (JCM) and showed that the cavity dissipation affected the coherence properties of the field. In these studies, the atom is a single pure two-level one. While, in some case, atomic level is degenerate in the projections of the angular momenta on the quantization axis (Reshetov, 2000). In Refs. (Zhou *et al.*, 2001, 2002), the authors studied the case of degenerate atomic levels and showed that the revival period of the atom population inversion became longer than that of the original JCM. In Ref. (Zhou *et al.*, 2003), our group has studied the effect of degenerate atomic levels on the field state dissipation in two-photon JCM. The initial degenerate atomic states are found to increase the period of entanglement of atom and field, and the coherence properties of field is affected by the reservoir qualitatively.

The interaction between an atom and a single-mode field *via* two-photon process is a nonlinear one. Giving the explicit solution of the dissipative dynamics of atom in a nonlinear process is very difficult. So the atomic decoherence property in a two-photon JCM with degenerate atomic levels is not studied before, despite that the field decoherence property has been studied (Zhou *et al.*, 2001, 2003). However, the atom is the information depositor in a quantum information system (Fleischhauer and Dluikiin, 2000); it is more important to study the atomic decoherence property than that of the field. In the present paper, considering degenerate atomic levels and using the method of perturbative expansion for master equation, we study the dissipative dynamics of system and of atom in two-photon JCM. We show that the period of entanglement between atom and field is prolonged explicitly. In case of degenerate atomic levels, contrasting to the case of two levels atom, the atomic asymptotic value of linear entropy is increased clearly in a same dissipative cavity. This asymptotic value is the largest when the atom is initially in an equal probability superposition degenerate state. Moreover, the numerical results show that the degenerate atomic levels increase the mixture degree of nonlinear situation of two-photon JCM.

2. PERTURBATIVE SOLUTION OF MASTER EQUATION

Let us consider an atom with degenerated levels interacts with a single-mode electromagnetic field at zero temperature as usual in a high- Q cavity. The dynamics of the global density operator $\hat{\rho}$ in interaction representation is governed by the

master equation

$$\frac{d}{dt}\hat{\rho} = -i [H_{\text{eff}}, \hat{\rho}(t)] + \kappa(2a\hat{\rho}a^\dagger - aa^\dagger\hat{\rho} - \hat{\rho}aa^\dagger), \tag{1}$$

where κ is the damping constant of the cavity. The effective Hamiltonian H_{eff} in two-photon JCM with atomic degeneracy is nonlinear. Thus, Equation (1) is a nonlinear differential equation. Completely solving the equation with the usual method used in the study of original JCM is very difficult. In order to study the dissipative dynamics of two-photon JCM, we can solve the master equation approximately (Yi *et al.*, 2000). While, here, we expand $\hat{\rho}(t)$ in powers of κ but not t as

$$\hat{\rho}(t) = \hat{\rho}^0(t) + \frac{\partial\hat{\rho}(t)}{\partial\kappa}\kappa + \frac{1}{2}\frac{\partial^2\hat{\rho}(t)}{\partial\kappa^2}\kappa^2 + 0(\kappa^2). \tag{2}$$

In a high- Q (quality factor of cavity) cavity, $\kappa \ll 1$, the effectiveness of the expansion is assured and the reservation of only fore 3 terms is satisfying.

From Equation (1), each order of differential of $\hat{\rho}(t)$ can be determined by using iterative method as

$$\frac{d}{dt}\hat{\rho}^{(n)}(t) = -i [\hat{H}_{\text{eff}}, \hat{\rho}^{(n)}(t)] + (2a\hat{\rho}^{(n-1)}(t)a^\dagger - aa^\dagger\hat{\rho}^{(n-1)}(t) - \hat{\rho}^{(n-1)}(t)aa^\dagger), \tag{3}$$

with $n = 1, 2, 3 \dots$, and $\hat{\rho}^{(n)}(t) = \partial^n \rho(t) / \partial \kappa^n$. The solution of Equation (3) is

$$\hat{\rho}^{(n)}(t) = e^{-i\hat{H}_{\text{eff}}t} \left\{ \int_0^t dt e^{i\hat{H}_{\text{eff}}t} [2a\hat{\rho}^{(n-1)}(t)a^\dagger - aa^\dagger\hat{\rho}^{(n-1)}(t) - \hat{\rho}^{(n-1)}(t)aa^\dagger] e^{-i\hat{H}_{\text{eff}}t} \right\} e^{i\hat{H}_{\text{eff}}t}. \tag{4}$$

So, $\hat{\rho}^{(1)}(t)$ can be calculated from an integer of $\hat{\rho}^{(0)}(t)$, and $\hat{\rho}^{(2)}(t)$ from $\hat{\rho}^{(1)}(t)$, ... Thus, we can get the precise value of $\hat{\rho}(t)$ at least in principle.

We assume that the initial states of the atom–field system can be written as

$$\Psi(0) = \left(\sum_{m_b} \frac{e}{\sqrt{2J_b + 1}} |J_b, m_b\rangle + \sum_{m_c} \frac{f}{\sqrt{2J_c + 1}} |J_c, m_c\rangle \right) \otimes |\alpha\rangle, \tag{5}$$

where J_b, J_c denote the values of the total electronic angular momenta of resonant levels b, c respectively, and m_b, m_c represent their projections on the polarized direction. $|\alpha\rangle$ is the coherence field state of cavity; it can be expanded as $|\alpha\rangle = \exp(-|\alpha|^2/2) \sum_n \alpha/\sqrt{n!} |n\rangle = \sum_n F_n |\alpha\rangle$. From Equation (5), we easily obtain the initial density of the system, i.e. $\hat{\rho}^{(0)}(0)$, and further obtain the evolution of $\hat{\rho}^{(0)}(0)$. In interaction representation, the effective Hamiltonian of the two-photon JCM within rotating-wave approximation including degenerate atomic levels has

been obtained in Ref. (Zhou *et al.*, 2003) as

$$\hat{H}_{\text{eff}} = \Omega[(a^\dagger a + 1)(a^\dagger a + 2)R_b - (a^\dagger a - 1)R_c], \quad (6)$$

where $R_\mu = \sum_{m\mu} \alpha_{m\mu}^2 (|J_\mu, m_\mu\rangle \langle J_\mu, m_\mu|)$, $\mu = b, c$. Ω is the Rabi frequency, which measures the coupling between atom and field. $\alpha_{m\mu}$ is defined as

$$\alpha_{m\mu} = (-1)^{J_b - m} \begin{pmatrix} J_b & 1 & J_c \\ -m & 0 & m \end{pmatrix}.$$

The linear entropy is defined as

$$S = 1 - \text{Tr}(\hat{\rho}^2), \quad (7)$$

which can be used to measure the coherence lose and the purity degree of the state. For a pure state $S = 0$, otherwise, $S > 0$, corresponding to a mixture.

Substitute $\hat{\rho}(t)$, $\hat{\rho}_a(t)$ into Equation (9). We get the linear entropy of system S and of atom S_{da} . Here, we use letter d to denote atomic linear entropy in JCM in case of degenerate atomic levels. The linear entropy S_{da} then depends on a set of parameters ($e, f, |\alpha|^2, J_b, J_c, \kappa/\Omega, \Omega t$). In next section, typical values of the parameters are given to discuss the behavior of S , S_{da} , and from the appearance of the linear entropy, the decoherence of the system and the atom and the disentanglement of atom and field are studied.

3. RESULTS AND DISCUSSION

In experiment (Brune *et al.*, 1987), the resonant atomic levels b, c were usually the Rydberg states of the atom with angular momenta $J_b = 3/2, J_c = 3/2$ or $J_c = 5/2$. By the support of these experiment data, we choose $J_b = J_c = 3/2$, which emerges $\alpha_{1/2} = \alpha_{-1/2} = 1/2\sqrt{15}, \alpha_{3/2} = \alpha_{-3/2} = 3/2\sqrt{15}$.

With same set of parameters, we plot the evolution of atomic linear entropy with and without degeneration to study the effect of atomic degeneracy on the dissipative dynamics of atom in Fig. 1.

Without degeneration, the atomic is a pure two-level one. Under this case, the linear entropy has been discussed in Ref. (Zhou *et al.*, 2004). We directly employ the result in that paper to show the effect of the atomic degeneracy. In the process of atomic linear entropy evolution, Fig. 1 shows that the behavior of $S_a(t)$ (atomic linear entropy in non-degenerate case) and S_{da} both present local maximum and minimum. This corresponds to the entanglement and disentanglement respectively of atom and field. In Ref. (Peixoto and Nemes, 1999), the authors have shown that disentanglement took place at instants $t_d = n\pi/\omega$. Here, contrasting to the case of non-degeneration (Zhou *et al.*, 2004), the period of entanglement and disentanglement in case of degenerate atomic levels is obviously prolonged. And the amplitude of local entanglement and disentanglement is suppressed. We can also observe that the two asymptotic values are obviously different, i.e. $S_{da}(\infty) >$




Fig. 1. Linear entropy of atom as a function of Ωt , S_{da} with degenerate atomic levels (solid line), S_a with two levels atomic state (dotted line), where $e = f = 1/\sqrt{2}$, $\kappa/\Omega = 0.02$, $|\alpha|^2 = 0.5$.

$S_a(\infty)$. The linear entropy is seen as a measure of the coherence loss. If the atom is initially in a degenerate state written in Equation (5), the coherence property is better than that of without degeneracy. When coherence loss finished completely, the better the initial coherence property the bigger the coherence loss. We will show the effect of initial coherence property on the coherence loss in Fig. 2. By comparing the atomic linear entropy with and without degeneracy, we find that the degeneracy increases the maximum of the asymptotic value of the linear




Fig. 2. Linear entropy of atom as a function of Ωt , where $\kappa/\Omega = 0.02$, $|\alpha|^2 = 0.5$, solid line with $e = f = 1/\sqrt{2}$, dotted line with $e = 0.4$.




Fig. 2. Continued.

entropy. Thus, the degeneracy of the atom on one hand increases the period of entanglement, and on the other hand increases the atomic maximum coherence loss. The result has not been given before.

The dependence of the linear entropy on the photon number is shown in Fig. 3. It is clear that the period of entanglement does not depend on the average photon number $|\alpha|^2$. And obviously, the larger the average photon number, the larger the asymptotic value of the curves. The similar results have been shown in Ref. (Peixoto and Nemes, 1999). In that paper, the asymptotic values of the linear




Fig. 3. (a) Linear entropy of the system (atom–field) as a function of Ωt , where $e = f = 1/\sqrt{2}$, $\kappa/\Omega = 0.01$, solid line with $|\alpha|^2 = 1$, dotted line with $|\alpha|^2 = 0.5$. (b) Linear entropy of the atom as a function of Ωt , where $e = f = 1/\sqrt{2}$, $\kappa/\Omega = 0.02$, solid line with $|\alpha|^2 = 1$, dotted line with $|\alpha|^2 = 0.5$.

entropy never exceed $\frac{1}{2}$, while here, the asymptotic values can do. The average photon number $|\alpha|^2$ is a measure of its classicality. The entanglement of atom and field proportionally contributes to S_{da} . Therefore, the larger the $|\alpha|^2$ is the more rapidly the atom and the system lose their coherence. These properties are also observed in Ref. (Peixoto and Nemes, 1999; Zhou *et al.*, 2002).

Now, we take different initial atomic degenerate states to show their effect on the disentanglement and decoherence of atom. In Fig. 2, when the atom is initially in degenerate state with $e = f = 1/\sqrt{2}$, the asymptotic value is the largest. In this case, the initial coherence of atom is the best; when dissipation finally finishes, the coherence loss is the most and the linear entropy is the largest. Figure 2 also shows that the amplitude of local maximum and minimum of the curves is the largest in this case. Because of the best initial coherence property of atom, the atom would best entangle with field. So, the better the initial atomic coherence the larger the amplitude of local maximum and minimum. Therefore, the better the initial atomic coherence properties, the larger the asymptotic value of S_{da} and the larger the amplitude of local maximum and minimum of S_{da} .

4. CONCLUSION

We have studied the dissipative dynamics of system and atom in two-photon JCM with degenerate atomic levels by employing the perturbative expansion of master equation. The results show that the degeneracy prolongs the period of entanglement and disentanglement of atom and field qualitatively. This provides us a way in preparing a long time surviving entangled quantum states needed in experiment (Cirac and Zoller, 1994b; Kudryavtsev and Knight, 1993; Phoenix and Barnett, 1993). The degeneracy also increases the asymptotic value of the linear entropy. This means the coherence loss is increased by the degeneracy. Comparing to the non-degenerate case, the amplitude of local maximum and minimum of the linear entropy is suppressed due to the degeneracy. The more average photon number affects the cavity dissipation by increasing the asymptotic value of the linear entropy of the atom and the system and making them lose their coherence more rapidly than usual. The better the initial coherence property of the atom the larger the asymptotic value of the linear entropy, i.e. the larger the coherence loss.

REFERENCES

- Brune, M. *et al.* (1996). *Physical Review Letters* **77**, 4887.
- Brune, M., Raimond, J. M. Goy, P., Davidovich, L., and Haroche, S. (1987). *Physics Letters* **59**, 1899.
- Cirac, J. I. and Parkins, A. S. (1994). *Physics Review A* **50**, R4441.
- Cirac, J. I. and Zoller P. (1994a). *Physics Review A* **50**, R2799.
- Cirac, J. I. and Zoller, P. (1994b). *Physics Review A* **50**, R4887.
- Cirac, J. I., Zoller, P., Kimble, H. J., and Mabuchi, H. (1996). *Physical Review Letters* **78**, 3321.

- Davidovich, L., Brune, M., Raimond, J. M., and S Haroche, S. (1996). *Physics Review A* **53**, 1295.
- Davidovich, L., Zagury, N., Brune, M., Raimond, J. M., and Haroche, S. (1994). *Physics Review A* **50**, R895.
- Duan, L. M. and Guo, G. C. (1998). *Physics Review A* **57**, 2399.
- Fleischhauer, M. and Dlukin, M. (2000). *Physical Review Letters* **84**, 5094.
- Gerry, C. C. (1996). *Physics Review A* **53**, 2857.
- Hagan, S., Hameroff, S. R., and Tuszyński, J. S. (2000). *eprint quant-ph/0005025*.
- Kudryavtsev I. K. and Knight, P. L. (1994). *Journal of Modern Optics* **40**, 1673.
- Peixoto, J. G. and Nemes, M. C. (1999). *Physics Review A* **59**, 3918.
- Pellizzari, T. (1997). *Physical Review Letters* **79**, 5242.
- Phoenix, S. J. and Barnett, S. M. (1993). *Journal of Modern Optics* **40**, 979.
- Reshetov V. A. (2000). *eprint physics/0002047*.
- Tegmark, M. (2000). *Physics Review E* **61**, 4194.
- Yi, X. X., Li, C., and Su, J. C. (2000). *Physics Review A* **62**, 013819.
- Zheng, S. B. and Guo, G. C. (2001). *Physical Review Letters* **85**, 2392.
- Zhou, L., Guo, Y. Q., Song, H. S., and Li, C. (2004). *Communications on Theoretical Physics* **41**, 111.
- Zhou, L., Song, H. S., and Li, Y. (2001). *Chinese Physics* **10**, 413.
- Zhou, L., Song, H. S., Li C., and Guo, Y. Q. (2003). *Chinese Physics* **12**, 45.
- Zhou, L., Song H. S., and Luo, Y. X. (2002). *Journal of Optics B* **4**, 103.
- Zhou, L., Song, H. S., Luo Y. X., and Li, C. (2001). *Physics Letters A* **284**, 156.
- Zurek, W. H. (1991). *Physics Today* **44**, 36.