

Homotopic Classification of Yang–Mills Vacua Taking into Account Causality

Gábor Etesi¹

Received May 19, 2006; accepted July 25, 2006
Published Online: January 31 2007

Existence of θ -vacuum states in Yang–Mills theories defined over asymptotically flat space-times examined taking into account not only the topology but the complicated causal structure of these space-times, too. By a result of Galloway apparently causality makes all vacuum states, seen by a distant observer, homotopically equivalent making the introduction of θ -terms unnecessary.

But a more careful analysis shows that certain twisted classical vacuum states survive even in this case eventually leading to the conclusion that the concept of “ θ -vacua” is meaningful in the case of general Yang–Mills theories. We give a classification of these vacuum states based on Isham’s results showing that the Yang–Mills vacuum has the same complexity as in the flat Minkowskian case hence the general CP-problem is not more complicated than the well-known flat one. We also construct the θ vacua rigorously via geometric quantization.

KEY WORDS: θ -vacua; asymptotical flatness; causality; topological censorship.

PACS numbers: 11.15, 11.30.E, 04.20.G, 04.70

1. INTRODUCTION: THE MINKOWSKIAN YANG–MILLS THEORY

The famous solution of the long-standing $U(1)$ -problem in the Standard Model via instanton effects was presented by ’t Hooft about three decades ago (’t Hooft, 1976, 1986). This solution demonstrated that *instantons* i.e., finite-action self-dual solutions of the *Euclidean* Yang–Mills-equations discovered by Belavin *et al.* (1975) should be taken seriously in gauge theories. Another problem arose in these models over the *Minkowskian* space-time, however: if instantons really exist, they induce a P-hence CP-violating so-called θ -term in the effective Yang–Mills action. But according to accurate experimental results, such a CP-violation cannot occur in QCD, for instance. The most accepted solution to this problem is the so-called *Peccei–Quinn mechanism* (Peccei and Quinn, 1977). A consequence

¹ Department of Geometry, Mathematical Institute, Budapest University of Technology and Economics, H ép., Egrý József u. 1., Budapest, H-1111 Hungary; e-mail: etesi@math.bme.hu.

of this mechanism is the existence of a light particle, the so-called *axion*. This particle has not been observed yet, however.

The question naturally arises whether or not such problematic θ -term must be introduced over more generic space-times. The aim of our paper is to claim that the answer is yes.

First, let us summarize the vacuum structure of a gauge theory over Minkowski space-time following basic text books (Cheng and Li, 1984; Kaku, 1993). Let E be a complex vector bundle over an oriented and time oriented Lorentzian manifold (M, g) belonging to a finite dimensional complex representation of G . Without loss of generality we choose the gauge group G to be a compact Lie group. Consider a G -connection ∇ on this bundle with curvature F_∇ ; we take the usual Yang–Mills action (by fixing the coupling to be 1):

$$S(\nabla, g) = -\frac{1}{8\pi^2} \int_M \text{tr}(F_\nabla \wedge *F_\nabla), \quad (1)$$

whose Euler–Lagrange equations are

$$d_\nabla F = 0, \quad d_\nabla *F = 0.$$

Here $*$ is the Hodge operation induced by the orientation and the metric on M . In our present case $M = \mathbb{R}^4$ and usually the metric g is fixed and supposed to be the Minkowskian one on \mathbb{R}^4 . Moreover all G -bundles E are trivial consequently by choosing a particular frame on E , the connection ∇ can be identified globally with a g -valued 1-form A .

The simplest solution is the vacuum i.e., a flat connection: $F_\nabla = 0$. By simply connectedness of \mathbb{R}^4 such gauge fields can be written in the form $A = f^{-1}df$, where $f : \mathbb{R}^4 \rightarrow G$ is a smooth function.

But by the existence of a global temporal gauge on \mathbb{R}^4 (in this gauge flat connections are independent of the “time” variable) it is enough to consider the restriction of f to a spacelike submanifold of Minkowski space-time i.e., $f : \mathbb{R}^3 \rightarrow G$. Minkowski space-time is asymptotically flat as well, so there is a point i^0 called spacelike infinity. This point represents the “infinity of space” hence can be added to \mathbb{R}^3 completing it to the three-sphere $\mathbb{R}^3 \cup \{i^0\} = S^3$. It is well-known that vacuum fields (possibly after a null-homotopic gauge-transformation around i^0) extend to the whole S^3 consequently classical vacua are classified by maps $f : S^3 \rightarrow G$. These maps up to homotopy are given by elements of $\pi_3(G)$. For typical compact Lie groups $\pi_3(G)$ is not trivial. This fact can be interpreted as classical vacua are separated from each other by energy barriers of finite height i.e., it is impossible to deform two vacua of different winding numbers into each other only through vacuum states. Hence homotopy equivalence reflects the *dynamical structure* of the theory.

On the other hand, vacua are also acted upon by the gauge group. For simplicity assume $G \cong SU(2)$. In this case $\pi_3(SU(2)) \cong \mathbb{Z}$. If f_1, f_2 are vacua of winding numbers n_1, n_2 respectively, there is a gauge transformation $g : S^3 \rightarrow SU(2)$ of winding number $n_2 - n_1$ satisfying $gf_1 = f_2$. Consequently we can see that the concept of *dynamical equivalence* of vacua reflecting the *dynamics* of the theory (i.e., the *homotopy equivalence* of maps $f : S^3 \rightarrow SU(2)$) is different from that of *symmetry equivalence* of vacua representing the *symmetry* of the gauge theory (i.e., the *gauge equivalence* of the above maps).

To avoid this discrepancy, we proceed as follows. Suppose we have constructed the Hilbert space $\mathcal{H}_{\mathbb{R}^4}$ of the corresponding quantum gauge theory. If $|n\rangle \in \mathcal{H}_{\mathbb{R}^4}$ denotes the quantum vacuum state belonging to a classical vacuum f of winding number n , the simplest way to construct a state which is invariant (up to phase) under both dynamical (i.e., homotopy) and symmetry (i.e., gauge) equivalence is to formally introduce the quantum state

$$|\theta\rangle := \sum_{n=-\infty}^{\infty} e^{in\theta} |n\rangle \in \mathcal{H}_{\mathbb{R}^4}, \quad \theta \in \mathbb{R}. \quad (2)$$

These formal sums are referred to as “ θ -vacua.”

From the physical point of view, the introduction of θ -vacua is also necessary. Although the vacuum states of different winding numbers are separated classically, they can be joined semi-classically i.e., by a tunneling induced by non-trivial instantons of the corresponding *Euclidean* gauge theory. Indeed, as it is well known, the $SU(2)$ instanton number is an element $k \in H^4(S^4, \mathbb{Z}) \simeq \mathbb{Z}$ (here S^4 is the one-point conformal compactification of the Euclidean flat \mathbb{R}^4). Note that the notion of “instanton number” comes from a very different compactification compared with the derivation of “vacuum winding number”). If two vacua, $|n_1\rangle, |n_2\rangle$ ($n_1, n_2 \in \pi_3(SU(2)) \simeq \mathbb{Z}$) are given then there is an instanton of instanton number $n_2 - n_1 \in H^4(S^4, \mathbb{Z}) \simeq \mathbb{Z}$ tunneling between them in temporal gauge (Cheng and Li, 1984; Kaku, 1993). In other words the true vacuum states are linear combinations of the vacuum states of unique winding numbers yielding again (2).

But the value of θ cannot be changed in any order of perturbation i.e., it should be treated as a physical parameter of the theory; this implies that tunnelings induce the effective term

$$\frac{\theta}{8\pi^2} \int_{\mathbb{R}^4} \text{tr}(F_{\nabla} \wedge F_{\nabla})$$

in addition to the action (1). But it is not difficult to see that such a term violates the parity symmetry P of the theory resulting in the violation of the CP-symmetry.

In summary, we have seen that there are at least three different ways to introduce θ -parameters in Yang–Mills theories *over Minkowskian space-time*:

- (i) θ is introduced to fill in the gap between the notions of dynamical (i.e., homotopy) and symmetry (i.e., gauge) equivalence of Yang–Mills vacua. This approach is pure mathematical in its nature;
- (ii) θ must be introduced because by instanton effects vacua of definite winding numbers are superposed in the underlying semi-classical Yang–Mills theory;
- (iii) θ must be introduced by “naturalness arguments” i.e., nothing prevents us to extend the Yang–Mills action at the full quantum level by a P -violating term $\text{tr}(F_{\nabla} \wedge F_{\nabla})$ with coupling constant θ .

There is a correspondence between the above three characterizations of the θ in *flat Minkowskian space-time* but in the case of general space-times, clear and careful distinction must be made until a relation or correspondence between the three notions is established. Clearly, (i) is related to the *topology* of the space-time and the gauge group hence it is relatively easy to check whether or not it remains valid in the general case. Concept (ii) is related to the semi-classical structure of the general Yang–Mills theory especially to the existence of instanton solutions in the Wick-rotated theory and their relationship with vacuum tunneling. The validity of concept (iii) is the most subtle one: we need lot of information on the global non-perturbative aspects of a general quantum Yang–Mills theory to check if any θ -term survives quantum corrections. In the present state of affairs, having no adequate general theory of Wick rotation, instantons and their physical interpretation, non-perturbative aspects of general Yang–Mills theories etc., we can examine only the validity of concept (i) in the general case. Its validity or invalidity may serve as a good indicator for the existence and role of θ -terms in general Yang–Mills theories.

The analysis of the vacuum structure of general Yang–Mills theories over a space-time (M, g) from the point of view of (i) was carried out by Isham *et al.* (Deser *et al.*, 1980; Isham, 1983; Isham and Kunstatter, 1982; Isham and Kunstatter, 1981). In these papers Isham *et al.* argue that in the general case concept (i) for introducing θ -terms still continues to hold due to the complicated topology of the spatial surface $S \subset M$ and the gauge group G (Isham, 1983). The classical vacuum structure of these theories becomes more complicated and we cannot avoid the introduction of various new CP-violating terms into the effective Lagrangian (Deser *et al.*, 1980).

We have to emphasize that the approach of Isham *et al.* to the problem is pure topological in its nature, however. By a result of Witt (1986) every oriented, connected three-manifold S appears as a Cauchy surface of a physically reasonable initial data set. It is well-known that the complicated topology of the spacelike submanifold S leads to appearance of singularities in space-time

if it arises as the Cauchy development of S . Indeed, an early result of Gannon (1975) shows that the Cauchy development of a non-simply connected Cauchy surface is geodesically incomplete i.e., singularities occur. If we accept the Cosmic Censorship Hypothesis, these singularities are hidden behind event horizons resulting in a non-trivial causal structure for these space-times, too. A theorem of Galloway (1995) (cf. an earlier version assuming stationarity by Chruściel–Wald (1994)) shows that distant observers can observe only simply connected portions of asymptotically flat space-times: all topological properties are hidden behind event horizons, eventually resulting again in a topologically simple *effective* space-time. Hence one may doubt if Isham’s conclusions remain valid.

In Section 2 we formulate Yang–Mills theories with an arbitrary compact gauge group over general asymptotically flat space-times satisfying the null energy condition with a single globally hyperbolic domain of outer communication. This model provides a good framework for analysing classical Yang–Mills vacua over causally non-trivial space-times. In this setup we simply mimic the above analysis concerning classical Yang–Mills vacua and find that although all vacua are topologically equivalent on the causally connected regime of the space-time, the appearance of a natural boundary condition on the event horizon (also a consequence of the causal structure) introduces non-trivial homotopy classes again.

In Section 3 we calculate explicitly the homotopy classes of vacua for the classical groups. A modification appears compared with Isham and other’s pure topological considerations in the sense that generally the vacuum structure in our case has exactly the same complexity as in the flat Minkowskian case, a surprising result. This demonstrates the “stability” of the θ -problem and justifies concept (i) even in the more general case.

The idea of studying relationship between micro- or virtual black holes, wormholes and θ -vacua is not new. For example, see Hawking (1996) and Preskil *et al.* (1989). An earlier, still incomplete version of this paper appeared in Etesi (2001).

2. ASYMPTOTICALLY FLAT YANG–MILLS THEORY

The general reference for this chapter is Wald (1984). Let (M, g) be a four dimensional, oriented and time oriented smooth Lorentzian manifold i.e., a space-time; choose a complex vector bundle E over M associated to a principal bundle with compact gauge group G via a finite dimensional complex representation. Consider a G -connection ∇ on E and a Yang–Mills theory with action (1) over (M, g) . We will focus on *vacuum solutions on a gravitational background* i.e., pairs (∇, g) where ∇ is a smooth flat G -connection on the bundle E while g is a smooth Lorentzian metric on M . We will suppose that g is a solution of the vacuum or the coupled Einstein’s equation with a matter field given by a

stress-energy tensor T obeying the null energy condition. We will refer the collection (E, ∇, M, g) to as an *Yang–Mills vacuum setup*.

We impose two restrictions. First, we will assume that (M, g) contains a single *asymptotically flat region*. At a first look (for the precise definitions see e.g. Wald (1984)) this means that there is a conformal embedding $i : (M, g) \rightarrow (\tilde{M}, \tilde{g})$ such that the infinitely distant points of M are represented by the connected set $\partial i(\overline{M})$ in the inclusion; furthermore this set is divided naturally into three subsets: the future and past null infinities \mathcal{I}^\pm and the spatial infinity i^0 . We remark that \tilde{g} is not supposed to be smooth in i^0 , even if (M, g) is smooth.

Now consider the *domain of outer communication* $\tilde{N} \subseteq \tilde{M}$ defined as $\tilde{N} := J^-(\mathcal{I}^+) \cap J^+(\mathcal{I}^-)$ and $N := \tilde{N} \cap i(M)$. (Here $J^\pm(X)$ denote the causal future and past of a subset X in a space-time, respectively). Notice that $N = M \setminus (B \cup W)$ where B and W are the black hole and white hole regions of M , respectively. The boundary $\partial(B \cup W)$ is called the *event horizon* of these regions. Our second assumption is that $(N, g|_N)$ is *globally hyperbolic*. Consequently $N \cong S \times \mathbb{R}$ with S being a Cauchy surface for the domain of outer communication N such that the image of the Cauchy surface can be completed to a maximal spacelike submanifold \tilde{S} in \tilde{M} by adding the spacelike infinity $i^0 \in \tilde{M}$ to it: $i(S) \cup \{i^0\} = \tilde{S}$.

Before proceeding further we fix notation. Let V be a smooth, compact, oriented three-manifold (possibly with non-empty boundary), $x^0 \in V \setminus \partial V$ and assume there is a homeomorphism $\varphi : V \setminus \partial V \rightarrow \tilde{S}$ such that $\varphi(x^0) = i^0$. In this case we will say that S is *homeomorphic to the interior of V* . By global hyperbolicity, there is a global time function $T : N \rightarrow \mathbb{R}$. Let $S_t := T^{-1}(t)$ ($t \in \mathbb{R}$) be a Cauchy surface which is the interior of a compact three-manifold V (notice that $S_t \cong S_{t'}$ for all $t, t' \in \mathbb{R}$). Consider a map $\varphi_t : V \setminus (\partial V \cup \{x^0\}) \rightarrow N$ whose image is $\varphi_t(V \setminus (\partial V \cup \{x^0\})) = S_t \subset N$. The points $\varphi_t(x) = (x, t)$ of S_t will be denoted as x_t . Clearly V represents the compactification of a particular Cauchy surface since $V \cong \overline{\varphi_t^{-1}(S_t)} \cong \overline{i(S_t) \cup \{i^0\}}$ for all $t \in \mathbb{R}$. Therefore by abuse of notation we will often think $S_t \subset V$ for all $t \in \mathbb{R}$.

Now we are in a position to address the problem of describing the topology of Yang–Mills vacua *seen by an observer in the domain of outer communication* of the space-time (M, g) . Clearly, at least classically, only this part of the space-time can be relevant for ordinary macroscopic observers. To achieve our goal, we refer to a general result of Galloway (1995) (for an earlier version assuming stationarity cf. Chruściel and Wald (1994)).

Theorem 2.1. (Galloway, 1995). *Let (M, g) be an asymptotically flat space-time containing a single asymptotically flat region whose domain of outer communication $(N, g|_N)$ is globally hyperbolic. Suppose that the null energy condition holds.*

Then N is simply connected i.e., $\pi_1(N) = 1$.

Assume there is a Cauchy surface S_t of N homeomorphic to the interior of a compact three-manifold V . Then if $\partial V \neq \emptyset$, each connected component of ∂V is homeomorphic to S^2 .

This rather surprising observation is a consequence of the so-called Topological Censorship Theorem of Friedman–Schleich–Witt (1993).

We can see that V , the compactification of a Cauchy surface S_t for N , is a simply connected (hence orientable) three-manifold. If M contains black or white hole domains then $\partial V \neq \emptyset$ and all boundary components are homeomorphic to a two-sphere S^2 (“the event horizon of a black or white hole in an asymptotically flat space-time has no handles”).

The following simple lemma ensures us that from a technical viewpoint the vacuum structure at least over the relevant part $(N, g|_N)$ is exactly the same as in the Minkowskian case.

Lemma 2.2. *Let (M, g) be a space-time as in Theorem 2.1 and (E, ∇, M, g) be a Yang–Mills vacuum setup over it. Consider the domain of outer communication with the restricted Yang–Mills data $(E|_N, \nabla|_N, N, g|_N)$. Then*

- (i) *If $\nabla|_N$ is flat and smooth then it can be identified with a \mathfrak{g} -valued 1-form A over N and there is a smooth function $f : N \rightarrow G$ such that $A = f^{-1}df$;*
- (ii) *There is a smooth gauge transformation $g : N \rightarrow G$ transforming $\nabla|_N$ into temporal gauge i.e., there is an $A' = gAg^{-1} + gdg^{-1}$ such that $A'_0 = 0$ where $A'_0 = A'(\text{grad}T)$. If A' is flat then the corresponding f does not depend on t ;*
- (iii) *Fix a $t \in \mathbb{R}$ and consider the restriction $f|_{S_t} =: f_t : S_t \rightarrow G$. Then f_t extends smoothly across the spacelike infinity i^0 i.e., there is a smooth function $\tilde{f}_t : \tilde{S}_t \rightarrow G$, homotopic to f_t on S_t .*

Proof: Concerning (i), the restricted bundle $E|_N$ is trivial hence any G -connection on it can be identified with a \mathfrak{g} -valued 1-form A ; simply connectedness of N implies that any flat connection $\nabla|_N$ must be the trivial connection hence in any gauge it can be represented in the form $A = f^{-1}df$ as claimed.

To see (ii) we can write down the required gauge transformation by solving the ordinary differential equation

$$gA_0g^{-1} + g \frac{\partial g^{-1}}{\partial t} = 0$$

over $N \cong S \times \mathbb{R}$. The solution over a chart $U \subset S$ is

$$g(x, t) = \exp \left(\int_0^t A_0(x, \tau) d\tau \right)$$

with $x \in U$, $t \in \mathbb{R}$ and $\exp: \mathfrak{g} \rightarrow G$ being the exponential map. This solution exists for finite t 's.

The case of part (iii) is also very simple. Notice that there is a neighbourhood $U \subset \tilde{S}_t$ of i^0 such that $U \setminus \{i^0\} \cong S^2 \times [0, 1)$. Consider the restriction $f_t|_{S^2 \times \{0\}}$ and take the function $\text{id}: S^2 \times [1/2, 0) \rightarrow G$ sending all elements to the unit $e \in G$. Then, taking into account that $\pi_2(G) = 0$ for compact Lie groups, there is a smooth homotopy from S^2 to G along $S^2 \times [0, 1/2]$ connecting $f_t|_{S^2 \times \{0\}}$ with $\text{id}|_{S^2 \times [1/2]}$. But this deformed function \tilde{f}_t extends as the identity to the whole \tilde{S}_t and is homotopic to f_t on S_t . \square

A pure Yang–Mills theory being conformally invariant, we may consider our Einstein-matter theory together with a Yang–Mills field over (\tilde{M}, \tilde{g}) instead of the original space-time. The restriction of the extended flat Yang–Mills bundle $\tilde{E}|_{\tilde{N}}$ is trivial even in this case. Certain physical quantities of the extended theory may suffer from singularities on the boundary $\partial i(\tilde{M})$ but classical Yang–Mills vacua in temporal gauge extend smoothly over the whole (\tilde{M}, \tilde{g}) as we have seen by the above lemma. In other words the studying of the vacuum sector of the extended Yang–Mills theory is correct.

Summing up, we can see that dynamically (i.e., homotopically) inequivalent vacua of the Yang–Mills theory are classified by the homotopy classes of smooth maps $f : V \rightarrow G$ satisfying $f(i^0) = e \in G$, usually written as

$$[(V, i^0), (G, e)]. \tag{3}$$

Now suppose that (M, g) contains black and white hole(s). In this case V is a simply connected compact three manifold *with boundary* by the theorem of Galloway. Such manifolds, considered as CW-complexes, have only cells of dimension less than three. Hence by the Cellular Approximation Theorem (Spanier, 1966), every map $f : V \rightarrow G$ descends to a homotopic map with values only on the cells of G having dimension less than three. Being $\pi_2(G) = 0$, G can be approximated by the simple Postnikov-tower $P_2 = K(\pi_1(G), 1)$ where $K(\pi_1(G), 1)$ is an Eilenberg–Mac Lane space yielding

$$[(V, i^0), (G, e)] \cong [V, K(\pi_1(G), 1)] \cong H^1(V, \pi_1(G)) = 0. \tag{4}$$

The result is zero because V is simply connected. For details, see for instance (Spanier, 1966). Consequently all vacuum states are homotopy-equivalent i.e., can be deformed into each other only through vacuum states *over the domain of outer communication N of the space-time (M, g)* . Clearly, classically only this part is relevant for a distant observer.

This result can be explained from a different point of view as well. Since the outer part N of M is globally hyperbolic by assumption, the spacelike submanifold S forms a Cauchy surface for N . Consequently if we know the initial values of two gauge fields, A and A' say, on $S \subset N$, we can determine their values over

the whole *outer* space-time $N \subset M$ by using the field equations. This implies that the values of the fields A and A' “beyond” the event horizon in a moment are irrelevant for an observer outside the black hole. But we just proved that every vacuum fields restricted to $V \supset S$ are homotopic. Roughly speaking, homotopical differences between Yang–Mills vacua “can be swept” into a black hole.

Via (4) for arbitrary smooth functions $f, g : V \rightarrow G$ there is a homotopy

$$F_T : V \times [0, 1] \rightarrow G \quad (5)$$

satisfying $F_T(x, 0) = f(x)$ and $F_T(x, 1) = g(x)$ and $F_T(i^0, t) = e$ for all $(x, t) \in V \times [0, 1]$. Taking two Cauchy surfaces $T^{-1}(t_0) =: S_0$ and $T^{-1}(t_1) =: S_1$ we can regard the two functions as vacua $f|_{S_0} := f_0 : S_0 \rightarrow G$ and $g|_{S_1} := f_1 : S_1 \rightarrow G$. In the homotopy F_T the subscript “ T ” shows that the “time” required for the homotopy is measured by the time function T naturally associated to the globally hyperbolic space-time (M, g) .

But on physical grounds, such a deformation or homotopy is effective only if the vacuum states, corresponding to the initial and final stages of the homotopy, can be compared by an observer in finite proper time. This means the following. Let $k = 0, 1$ and for all $x_k \in S_k$ for which $f_0(x_0) \neq f_1(x_1)$ there must exist an observer $\gamma : \mathbb{R} \rightarrow N$ moving forward in the region N who can measure hence compare $f_0(x_0)$ and $f_1(x_1)$ i.e., there are $\tau_k \in \mathbb{R}$ such that a future directed light beam starting from x_k meets γ in $\gamma(\tau_k)$, and the proper time between $\gamma(\tau_0)$ and $\gamma(\tau_1)$ measured by γ is finite. In other words, there is a $\tau^- \in \mathbb{R}$ such that $C \subset J^-(\gamma(\tau^-))$ where $C \subset S_0 \times S_1$ contains the set of all points where $f_0(x_0) \neq f_1(x_1)$ with $x_k \in S_k$. Because our space-time may contain white hole regions too, we require the existence of another $\tau^+ < \tau^-$ satisfying $C \subset J^+(\gamma(\tau^+))$ as well. The formal definition of such “effective” or “observable” homotopies is the following.

Definition 2.3. Let (M, g) be an asymptotically flat space-time with a single globally hyperbolic domain of outer communication $(N, g|_N)$ and let $T : N \rightarrow \mathbb{R}$ be an associated time-function. Consider a homotopy of the form (5) and let $C \subset S_0 \times S_1$ be such that $f_0(x_0) \neq f_1(x_1)$ with $(x_0, x_1) \in C$.

Then (5) is called an *effective homotopy* or *observable homotopy* if there is a future directed non-spacelike piecewise smooth curve $\gamma : \mathbb{R} \rightarrow N$ and fixed numbers $\tau^\pm \in \mathbb{R}$ such that $C \subset J^\pm(\gamma(\tau^\pm))$.

Remark. We can see that in Minkowski space-time all homotopies of the form (5) are effective homotopies establishing the structure of the Minkowskian Yang–Mills vacuum also from a physical viewpoint.

The following lemma is straightforward.

Lemma 2.4. *Let (M, g) be a space-time as in Definition 2.3. and consider a continuous curve $x : [0, \varepsilon] \rightarrow V$ satisfying $x(0) \in \partial V$. Then we have induced*

spacelike curves $x_k : [0, \varepsilon] \rightarrow S_k$ ($k = 0, 1$) given by $x_k(s) = (x(s), k)$ and satisfy $x_k(0) \in H$.

The (abstract) homotopy (5) is effective if and only if there is a $0 < \delta$ such that

$$F_T(x_0(s), t) = F_T(x_1(s), t)$$

for all $0 \leq s < \delta < \varepsilon$ and $t \in [0, 1]$ that is, the homotopy is trivial in the vicinity of H .

Proof: We restrict our attention first to the case $H^+ = \partial B$, the future event horizon of the black hole regime B . Take a homotopy of the form (5) with $F_T(x_k(s), k) = f_k(x_k(s))$ ($k = 0, 1$) and assume F_T is effective. By construction $x_k(0) \in H^+$ and if $f_0(x_0(0)) \neq f_1(x_1(0))$ then there must exist a future directed non-spacelike curve $\gamma : \mathbb{R} \rightarrow N$ such that $\{x_0(0), x_1(0)\} \subset J^-(\gamma(\tau^-))$ for some $\tau^- \in \mathbb{R}$. However this contradicts the definition of the domain of outer communication N consequently we must have $x_k(0) \notin H^+$. We get the same result for the past white hole horizon $H^- = \partial W$. Therefore $x_k(0) \notin H = H^+ \cup H^-$ as claimed. \square

From here we can see that given an abstract homotopy (5), it gives rise to an effective homotopy if and only if F_T is constant along H . This result can be interpreted as a natural boundary condition on each connected component of ∂V for effectively deformable vacua dictated by the causal structure. Since each boundary component in a “moment” is homeomorphic to the two-sphere S^2 and $\pi_2(G) = 0$ we can extend f_0, f_1 within their homotopy classes in the spirit of part (iii) of Lemma 2.2 to functions $f, g : V \rightarrow G$ obeying $f(\partial V) = g(\partial V) = e \in G$. The same argumentation yields the conditions $f(i^0) = g(i^0) = e$. We just remark that exactly this is the physical reason for keeping the functions as identity in spacelike infinity i^0 when we discuss homotopy classes of vacua over Minkowskian space-time: the spacelike infinity is invisible for an observer in N .

Therefore the classes of effectively deformable vacua are given by the homotopy classes of functions $f : V \rightarrow G$ with the property $f(\partial V) = f(i^0) = e \in G$. The homotopy is also restricted to obey these boundary conditions. This set is denoted by

$$[(V, \partial V, i^0), (G, e)] \tag{6}$$

and replaces (3). To get a more explicit description of this set, we proceed as follows.

3. HOMOTOPIC CLASSIFICATION

First taking into account that a function $f : V \rightarrow G$ we are interested in satisfies that it sends each connected component of ∂V into the unit element $e \in G$,

we can replace the simply connected, compact three-manifold-with-boundary V with a closed, simply connected three-manifold W in the following way. Let us denote by $k > 0$ the number of connected components of ∂V (i.e., the number of black holes and white holes). As we have seen, all such component is an S^2 . Hence we can glue to each such component a three-ball B^3 using the identity function of S^2 to get a three-manifold without boundary

$$W := V \cup_{\partial V} \underbrace{B^3 \cup \dots \cup B^3}_k.$$

Clearly, f extends as the identity to each ball giving rise to the function $f : W \rightarrow G$. Consequently, if we fix a point x^n in each ball ($n \leq k$), then we may equivalently consider functions obeying $f(x^1) = \dots = f(x^k) = f(i^0) = e$. Modifying the allowed homotopies to obey this constraint, we can replace the homotopy set (6) by

$$[(W, x^1, \dots, x^k, i^0), (G, e)]$$

(of course if $k = 0$ then no point except i^0 is distinguished in W). We prove the following proposition:

Proposition 3.1. *Fix a number $k > 0$ and consider the connected, closed, simply connected three-manifold with $k + 1$ distinguished points $(W, x^1, \dots, x^k, i^0)$ constructed above. Denote by (W, i^0) the same space with only one distinguished point. Then there is a natural bijection*

$$[(W, x^1, \dots, x^k, i^0), (G, e)] \cong [(W, i^0), (G, e)]$$

by forgetting the points $x^1, \dots, x^k \in W$ and modifying the allowed homotopies accordingly.

Proof: Fix a number $k \geq 0$. First it is straightforward that if two functions, f_0 and f_1 are homotopic in $[(W, x^1, \dots, x^k, i^0), (G, e)]$ then they represent the same homotopy class in $[(W, i^0), (G, e)]$ i.e., they are homotopic in the later set as well. This is because the allowed homotopies in $[(W, i^0), (G, e)]$ are less restrictive than in $[(W, x^1, \dots, x^k, i^0), (G, e)]$.

Conversely, it is not difficult to see that in each class $[f] \in [(W, i^0), (G, e)]$ there is a representant which belongs to $[(W, x^1, \dots, x^k, i^0), (G, e)]$. Indeed, choose an arbitrary representant $f \in [f] \in [(W, i^0), (G, e)]$ and consider the pre-image $f^{-1}(e) \subset W$. This pre-image contains the point $i^0 \in W$ by construction. Taking into account that W is path connected, we can deform $f^{-1}(e)$ to contain the points x^1, \dots, x^k as well producing a representant which belongs to $[(W, x^1, \dots, x^k, i^0), (G, e)]$.

Now suppose that there are two functions f_0 and f_1 which are homotopic in $[(W, i^0), (G, e)]$ i.e., there is a continuous function $F : (W, i^0) \times [0, 1] \rightarrow (G, e)$

with

$$F(x, 0) = f_0(x), \quad F(x, 1) = f_1(x),$$

$$F(i^0, t) = e \quad \text{for all } t \in [0, 1] \text{ and } x \in (W, i^0).$$

For the sake of simplicity, assume they represent elements in $[(W, x^1, \dots, x^k, i^0), (G, e)]$, too. Then we have to prove that they are also homotopic in $[(W, x^1, \dots, x^k, i^0), (G, e)]$ i.e., there is a function $F' : (W, x^1, \dots, x^k, i^0) \times [0, 1] \rightarrow (G, e)$ with the property

$$F'(x, 0) = f_0(x), \quad F'(x, 1) = f_1(x),$$

$$F'(x^1, t) = \dots = F'(x^k, t) = F'(i^0, t) = e$$

for all $t \in [0, 1]$ and $x \in (W, x^1, \dots, x^k, i^0)$. From here we can see that the orbit of an arbitrary distinguished point x^n is a loop $l^n : [0, 1] \rightarrow G$ under the homotopy F while the constant loop in the case of F' . Hence if these loops are homotopically trivial in G then we can deform F into the homotopy F' by shrinking the loops l^1, \dots, l^k .

Now we prove that this is always possible. First, if $\pi_1(G) = 1$ i.e., the compact Lie group is simply connected then certainly each loop l^n is homotopic to the constant loop. Consequently assume $\pi_1(G) \neq 1$. Consider a distinguished point $x^n \in W$ and two paths $a^n : [0, 1/2] \rightarrow W$ with $a^n(0) = i^0$ and $a^n(1/2) = x^n$ and $b^n : [1/2, 1] \rightarrow W$ with $b^n(1/2) = x^n$ and $b^n(1) = i^0$. These give rise to a continuous loop $b^n * a^n : [0, 1] \rightarrow W$ with $b^n * a^n(0) = b^n * a^n(1) = i^0$. Here $*$ refers to the juxtaposition of curves, loops, etc. The loop $b^n * a^n$ is homotopic to the trivial loop since W is simply connected. Consider the maps $\alpha_0^n := f_0 \circ a^n : [0, 1/2] \rightarrow G$ and $\beta_0^n := f_0 \circ b^n : [1/2, 1] \rightarrow G$. These are loops in G hence so is their product $\beta_0^n * \alpha_0^n$. Construct the same kind of loops $\alpha_1^n := f_1 \circ a^n$ and $\beta_1^n := f_1 \circ b^n$. The product loop $\beta_1^n * \alpha_1^n$ is homotopic in G to $\beta_0^n * \alpha_0^n$ i.e., $[\beta_0^n * \alpha_0^n] = [\beta_1^n * \alpha_1^n]$ because f_0 is homotopic to f_1 . It is clear that

$$\beta_1^n * \alpha_1^n = \beta_0^n * l^n * \alpha_0^n.$$

Consequently we can write for the homotopy classes in question

$$[\beta_1^n * \alpha_1^n] = [\beta_0^n * l^n * \alpha_0^n] = [\beta_0^n][l^n][\alpha_0^n] = [\beta_0^n][\alpha_0^n][l^n]$$

$$= [\beta_0^n * \alpha_0^n][l^n] = [\beta_1^n * \alpha_1^n][l^n].$$

In the third step we have exploited the fact that a topological group always has commutative fundamental group (Postnikov, 1987). This shows that $[l^n] = 1$ that is the loop l^n is contractible in G for all $0 \leq n \leq k$ in other words the homotopy F is deformable into a homotopy F' yielding f_0 and f_1 are homotopic in $[(W, x^1, \dots, x^k, i^0), (G, e)]$ as well. □

The above proposition enables us to give a more explicit description of the set (6).

Theorem 3.2. *Let (M, g) be a space-time obeying the null energy condition. Assume it contains a single asymptotically flat region with globally hyperbolic domain of outer communication. Suppose this region contains a Cauchy surface homeomorphic to the interior of a compact three-manifold V . Let G be a typical compact Lie group i.e., let G be $U(n)$ with $n \geq 2$, or $SO(n)$, $Spin(n)$ with $n \neq 4$, or $SU(n)$, $Sp(n)$ for all n , or G_2, F_4, E_6, E_7, E_8 . Then we have*

$$[(V, \partial V, i^0), (G, e)] \cong \mathbb{Z}.$$

Moreover we have

$$[(V, \partial V, i^0), (U(1), e)] \cong 0,$$

and

$$[(V, \partial V, i^0), (SO(4), e)] \cong [(V, \partial V, i^0), (Spin(4), e)] \cong \mathbb{Z} \oplus \mathbb{Z}$$

for the remaining cases.

Proof: In light of the above considerations and Proposition 3.1, we have

$$[(V, \partial V, i^0), (G, e)] \cong [(W, x^1, \dots, x^k, i^0), (G, e)] \cong [(W, i^0), (G, e)].$$

Hence we can use the results of Isham who classified the set $[(W, i^0), (G, e)]$ and it is summarized in Isham (1983) in Table 1 on p. 207. But in our case W is a connected, closed, simply connected three-manifold hence the above result follows. \square

Remark. We mention that assuming the validity of the three dimensional Poincaré conjecture i.e., if $W \cong S^3$, then our theorem can be derived without using Isham's result since in this case we have simply $[(V, \partial V, i^0), (G, e)] \cong \pi_3(G)$.

We can see by this result that although the homotopy set (6) of effectively deformable vacua is typically non-trivial, it is remarkable more simple than in the original calculations of Isham *et al.* based on topological considerations only. The homotopy sets listed in Theorem 3.2 are exactly the same as for the flat Minkowskian case. Being all these vacua gauge equivalent (since N is simply connected) we have to introduce again linear combinations like (2) in this more general situation. Consequently we can see that approach (i) to the θ -parameter, mentioned in the Introduction, still makes sense in the general case.

4. CONCLUSION AND OUTLOOK

In this paper we have studied the concept of θ -vacua in general Yang–Mills theories. In light of our results, we can see that for outer observers in

asymptotically flat space-times θ -vacua do occur in a Yang–Mills theory and they can be constructed in a rigorous way by referring to geometric quantization as follows.

For simplicity we restrict attention to a simple gauge group G . For the moduli space \mathcal{V} of classical vacuum solutions of a Yang–Mills theory over an asymptotically flat region $(N, g|_N)$ we have the identification $\mathcal{V} \cong [(V, \partial V, i^0), (G, e)]$ as we have seen. Furthermore Theorem 3.2 says that as a set we have a homeomorphism $\mathcal{V} \cong \mathbb{Z}$. This space, regarded as a noncompact zero dimensional manifold is naturally identified with its cotangent bundle $T^*\mathcal{V}$. This setup resembles the situation of a *real polarization* in geometric quantization. Within this framework then the Hilbert space of the vacuum sector of the corresponding quantum Yang–Mills theory is identified with the space of L^2 functions on \mathcal{V} :

$$\mathcal{H}_N = \{f : \mathcal{V} \rightarrow \mathbb{C} \mid \|f\|_{L^2(\mathcal{V})} < \infty\}.$$

This means that an element $f \in \mathcal{H}_N$ is described by complex numbers a_n with $-\infty < n < \infty$ satisfying simply

$$\sum_{n=-\infty}^{\infty} |a_n|^2 < \infty.$$

Assigning to this function f the convergent Fourier series

$$f(\theta) := \sum_{n=-\infty}^{\infty} a_n e^{in\theta}$$

we identify naturally this vacuum Hilbert space with the space of square integrable functions on the circle

$$\mathcal{H}(S^1) := \{f : S^1 \rightarrow \mathbb{C} \mid \|f\|_{L^2(S^1)} < \infty\}.$$

The isomorphism $\mathcal{H}_N \cong \mathcal{H}(S^1)$ provides a very straightforward, natural introduction of a θ parameter into the vacuum sector of Yang–Mills theories as was done heuristically in (2). Moreover we can see that the whole quantum vacuum is just linear combination of θ -vacua. The generalization to non-simple gauge groups is clear. Observe however that in this picture the role of causality is extremely important: without it the classical moduli \mathcal{V} considered over the whole original space-time (M, g) would be complicated topologically in the sense that in general even the connected components of \mathcal{V} would be non-zero dimensional manifolds with non-trivial topology introducing some degeneracy into the vacuum sector.

In summary we can say that despite the possible complicated topology of the underlying Cauchy surface of an asymptotically flat space-time, the vacuum structure is similar to the flat Minkowskian case, due to the causal structure of these space-times which is complicated in the general case, too. Hence the introduction of the various new CP-violating terms studied in Deser *et al.* (1980)

are unnecessary. Taking seriously the causal structure experienced by an observer also fits with the *Heisenberg dictum* that quantum field theory should be formulated in terms of observers.

The suppression of the topology of the underlying Cauchy surface is due to the result of Galloway or Chruściel–Wald which is a consequence of the so-called Topological Censorship Theorem of Friedman–Schleich–Witt (1993). Consequently, the reduction of the problem of the general CP-violation to the flat Minkowskian case is essentially due to this result. However Topological Censorship remains valid in a more general (i.e., not only an asymptotically flat) setting (Galloway *et al.*, 1999); therefore we may expect that our attack on the first approximation of the CP-problem may continue to hold in these more general situations.

Finally, natural questions arise: Are there instanton solutions in the corresponding Wick-rotated theories? Recent results on constructing $SU(2)$ instanton solutions over various gravitational instantons may point towards this possibility (Etesi, 2003; Etesi and Hausel, 2003). What is the physical relevance of these solutions? Do they induce semi-classical tunnelings between vacuum states of different effective winding numbers? If the answer for these questions is yes, beyond (i) we have another, more physical, reason to introduce θ -vacua by concept (ii), also mentioned in the Introduction.

ACKNOWLEDGMENTS

The author expresses his thanks to G.W. Gibbons for calling attention for Isham's papers and the stimulating comments in Japan six years ago. The work was partially supported by OTKA grants No. T43242 and No. 046365 (Hungary).

REFERENCES

- Belavin, A. A., Polyakov, A. M., Schwarz, A. S., and Tyupkin, Yu. S. (1975). Pseudoparticle solutions of the Yang–Mills equations. *Physics Letters B* **59**, 85–87.
- Cheng, L.-P. and Li, L.-F. (1984). *Gauge Theory of Elementary Particle Physics*, Clarendon Press.
- Chruściel, P. T. and Wald, R. M. (1994). On the topology of stationary black holes. *Classical and Quantum Gravity* **11**, L147–L152.
- Deser, S., Duff, M. J., and Isham, C. J. (1980). Gravitationally induced CP-effects. *Physics Letters B* **93**, 419–423.
- Etesi, G. (2001). The structure of Yang–Mills vacua seen by distant observers. In: Sidharth, B. G. and Altaisky, M. V. (Ed.), *Consistent equation of classical gravitation to quantum limit and beyond*. Kluwer Academic/Plenum Press Publishers, New York.
- Etesi, G. (2003). Classification of 't Hooft instantons over multi-centered gravitational instantons. *Nuclear Physics B* **662**, 511–530.
- Etesi, G. and Hausel, T. (2003). On Yang–Mills instantons over multi-centered gravitational instantons. *Communication in Mathematical Physics* **235**, 275–288.
- Friedman, J. L., Schleich, K., and Witt, D. M. (1993). Topological censorship. *Physical Review Letters* **71**, 1486–1489.

- Galloway, G. J. (1995). On the topology of the domain of outer communication. *Classical and Quantum Gravity* **12**, L99–L101.
- Galloway, G. J., Schleich, K., Witt, D. M., and Woolgar, E. (1999). Topological censorship and higher genus black holes. *Physical Review D* **60**, 104039, 11.
- Gannon, D. (1975). Singularities in nonsimply connected space-times. *Journal of Mathematical Physics* **16**, 2364–2367.
- Hawking, S. W. (1996). Virtual black holes. *Physical Review D* **53**, 3099–3107.
- Isham, C. J. (1983). Vacuum tunneling in static space-times. In: Van Der Merwe, A. (Ed.), *Old and new Questions in Physics, Cosmology, Philosophy and Theoretical Biology*, Plenum Press, New York, 189–211.
- Isham, C. J. and Kunstatter, G. (1981). Yang–Mills canonical vacuum structure in a three-space. *Physics Letters B* **102**, 417–420.
- Isham, C. J. and Kunstatter, G. (1982). Spatial topology and Yang–Mills vacua. *Journal of Mathematical Physics* **23**, 1668–1677.
- Kaku, M. (1993). *Quantum Field Theory*, Oxford University Press, Oxford.
- Peccei, R. D. and Quinn, H. R. (1977). Constraints imposed by CP conservation in the presence of pseudoparticles. *Physical Review D* **16**, 1791–1797.
- Postnikov, M. (1987). *Lectures on Geometry V.: Lie groups and Lie algebras*, Mir Publishers, Moscow.
- Preskill, J., Trivedi, S. P., and Wise, M. B. (1989). Wormholes in spacetime and θ_{QCD} . *Physics Letters B* **223**, 26–31.
- Spanier, E. H. (1966). *Algebraic Topology*, Springer–Verlag, Berlin.
- 't Hooft, G. (1976). Symmetry breaking through Bell–Jackiw anomalies. *Physical Review Letters* **37**, 8–11.
- 't Hooft, G. (1986). How instantons solve the U(1) problem. *Physics Reports* **142**, 357–387.
- Wald, R. M. (1984). *General Relativity*, University of Chicago Press, Chicago.
- Witt, D. M. (1986). Vacuum space-times that admit no maximal slice. *Physical Review Letters* **57**, 1386–1389.