

Reichenbach's Common Cause in an Atomless and Complete Orthomodular Lattice

Yuichiro Kitajima

Received: 30 April 2007 / Accepted: 13 June 2007 / Published online: 6 July 2007
© Springer Science+Business Media, LLC 2007

Abstract Hofer-Szabo, Redei and Szabo (Int. J. Theor. Phys. 39:913–919, 2000) defined Reichenbach's common cause of two correlated events in an orthomodular lattice. In the present paper it is shown that if logical independent elements in an atomless and complete orthomodular lattice correlate, a common cause of the correlated elements always exists.

Keywords Orthomodular lattice · Reichenbach · Common cause

1 Introduction

Let $p(A)$ and $p(B)$ be probabilities of the events A and B respectively. When $p(A \wedge B) > p(A)p(B)$, the two events A and B are said to be correlated. Reichenbach [9] regarded the event C as a common cause of the correlated events A and B when the following conditions were satisfied.

- $0 < p(C) < 1$
- $p(A \wedge B|C) = p(A|C)p(B|C)$
- $p(A \wedge B|\neg C) = p(A|\neg C)p(B|\neg C)$
- $p(A|C) > p(A|\neg C)$
- $p(B|C) > p(B|\neg C)$

When a common cause of the events A and B exists, A and B are correlated ([9], pp. 159–160). When the events A and B are correlated, does a common cause of A and B exist? Such a problem are investigated by Gyenis and Redei [1] and Redei and Summers [8].

Gyenis and Redei [1] examined whether Reichenbach's common cause of two logically independent and correlated elements in a Boolean algebra exists or not. They showed that uncountably infinite common causes of two logically independent and correlated elements in an atomless Boolean algebra exist ([1], Proposition 7).

Y. Kitajima (✉)
Department of Philosophy, Kyoto University, Kyoto, Japan
e-mail: yuichiro.kitajima@at7.ecs.kyoto-u.ac.jp

Algebraic quantum field theory predicts correlations between two events which belong to spacelike separated spacetime regions in many normal states (e.g. [2]). Redei [7] defined Reichenbach's common cause in algebraic quantum field theory, and Summers and Redei [8] showed that if two events which belong to spacelike separated regions V_1 and V_2 correlate in a faithful normal state, a common cause exists, which locates in the union of the backward light cones of V_1 and V_2 (see also [10]).

Hofer-Szabo, Redei and Szabo [3] defined Reichenbach's common cause in an orthomodular lattice. In the present paper we examine whether Reichenbach's common cause of two correlated elements in an orthomodular lattice exists or not. We show that if logical independent elements in an atomless and complete orthomodular lattice correlate in a completely additive probability measure, a nontrivial common cause of these elements always exists (Proposition 3.9), and that if logically independent elements in an atomless and complete orthomodular lattice correlate in a faithful completely additive probability measure, uncountably infinite common causes of these elements exist (Proposition 3.10).

2 Mathematical Preliminaries

Definition 2.1 A lattice \mathcal{L} is a partially ordered set any two of whose elements a and b have a least upper bound $a \vee b$ and a greatest lower bound $a \wedge b$, which are respectively called the join and the meet of a and b . The least element and the greatest element, if they exist, are denoted by 0 and 1 respectively.

Definition 2.2 A lattice \mathcal{L} with 0 and 1 is called orthocomplemented when there is a mapping $a \rightarrow a^\perp$ of \mathcal{L} into itself satisfying the following three conditions:

1. $a \vee a^\perp = 1$ and $a \wedge a^\perp = 0$.
2. $a \leq b$ implies $a^\perp \geq b^\perp$.
3. $(a^\perp)^\perp = a$ for every a .

When $a \leq b^\perp$, we say that a and b are orthogonal.

Definition 2.3 An orthocomplemented lattice \mathcal{L} is called orthomodular when $a \leq b$ implies $b = a \vee (b \wedge a^\perp)$.

Definition 2.4 Let a and b be elements of an orthomodular lattice \mathcal{L} . We say that a commutes with b and we write aCb when $a = (a \wedge b) \vee (a \wedge b^\perp)$.

Definition 2.5 A complete lattice is a lattice \mathcal{L} in which for every subset S of \mathcal{L} , the join $\bigvee\{a|a \in S\}$ and the meet $\bigwedge\{a|a \in S\}$ exists.

For example, the set of all projections of a von Neumann algebra is a complete orthomodular lattice.

Definition 2.6 An orthomodular lattice is called atomless when for any nonzero element $a \in \mathcal{L}$ there exists $b \in \mathcal{L}$ such that $0 < b < a$.

For example, the set of all projections of a von Neumann algebra without direct summand of type I is an atomless orthomodular lattice.

In the present paper, \mathbb{N} denotes $\{1, 2, 3, \dots\}$. For mutually orthogonal elements x_1, \dots, x_n in an orthomodular lattice, $x_1 + \dots + x_n$ denotes $x_1 \vee \dots \vee x_n$.

Definition 2.7 The mapping μ of an orthomodular lattice \mathcal{L} to $[0, 1]$ is called a completely additive probability measure on \mathcal{L} when μ satisfies the following conditions.

1. $\mu(1) = 1$
2. For any mutually orthogonal elements $\{x_a\}_{a \in A}$, if $\sum_{a \in A} x_a$ exists,

$$\mu\left(\sum_{a \in A} x_a\right) = \sum_{a \in A} \mu(x_a)$$

For example, a normal state on a von Neumann algebra is a completely additive probability measure on the set of all projections of a von Neumann algebra.

Definition 2.8 Let μ be a completely additive probability measure on an orthomodular lattice \mathcal{L} . μ is called faithful when $\mu(x) \neq 0$ for any nonzero element $x \in \mathcal{L}$.

Lemma 2.9 Let μ be a completely additive probability measure on an orthomodular lattice \mathcal{L} and let x and y be elements in \mathcal{L} .

1. $\mu(x^\perp) = 1 - \mu(x)$.
2. If x commutes with y , $\mu(x) = \mu(x \wedge y) + \mu(x \wedge y^\perp)$.
3. If $x \leq y$, $\mu(x) \leq \mu(y)$.
4. If μ is faithful and $x \leq y$, $x = y$ is equivalent to $\mu(x) = \mu(y)$.

Proof

1. Since $1 = \mu(1) = \mu(x + x^\perp) = \mu(x) + \mu(x^\perp)$, $\mu(x^\perp) = 1 - \mu(x)$.
2. Since $x \mathcal{C} y$, $x = (x \wedge y) + (x \wedge y^\perp)$. Therefore $\mu(x) = \mu(x \wedge y) + \mu(x \wedge y^\perp)$.
3. Since $y = x + (x^\perp \wedge y)$, $\mu(y) = \mu(x) + \mu(x^\perp \wedge y) \geq \mu(x)$.
4. When $\mu(x) = \mu(y)$, $\mu(x^\perp \wedge y) = 0$ because $\mu(y) = \mu(x) + \mu(x^\perp \wedge y)$. Since μ is faithful, $x^\perp \wedge y = 0$, which implies $x = y$. \square

3 Existence of Reichenbach's Common Cause in an Atomless and Complete Orthomodular Lattice

Definition 3.1 ([3], p. 914) Let \mathcal{L} be an orthomodular lattice, let μ be a completely additive probability measure on \mathcal{L} and let a and b be elements of \mathcal{L} . $c \in \mathcal{L}$ is called a common cause of a and b in μ when $cCa, cCb, 0 < \mu(c) < 1$ and c satisfies the following equations.

$$\frac{\mu(a \wedge b \wedge c)}{\mu(c)} = \frac{\mu(a \wedge c)}{\mu(c)} \frac{\mu(b \wedge c)}{\mu(c)}, \quad (1)$$

$$\frac{\mu(a \wedge b \wedge c^\perp)}{\mu(c^\perp)} = \frac{\mu(a \wedge c^\perp)}{\mu(c^\perp)} \frac{\mu(b \wedge c^\perp)}{\mu(c^\perp)}, \quad (2)$$

$$\frac{\mu(a \wedge c)}{\mu(c)} > \frac{\mu(a \wedge c^\perp)}{\mu(c^\perp)}, \quad (3)$$

$$\frac{\mu(b \wedge c)}{\mu(c)} > \frac{\mu(b \wedge c^\perp)}{\mu(c^\perp)}. \quad (4)$$

If a commutes with b , $\mu(a) = \mu(a \wedge b) + \mu(a \wedge b^\perp)$ and $\mu(b) = \mu(a \wedge b) + \mu(a^\perp \wedge b)$ by Lemma 2.9. So for any elements a and b such that aCb and $\mu(a \wedge b) > \mu(a)\mu(b)$,

$$\frac{\mu(a \wedge b)}{\mu(b)} > \frac{\mu(a \wedge b^\perp)}{\mu(b^\perp)} \quad \text{and} \quad \frac{\mu(b \wedge a)}{\mu(a)} > \frac{\mu(b \wedge a^\perp)}{\mu(a^\perp)}$$

since $\mu(a \wedge b) > \mu(a)\mu(b)$ implies

$$\frac{\mu(a \wedge b)}{\mu(b)} > \frac{\mu(a) - \mu(a \wedge b)}{1 - \mu(b)} \quad \text{and} \quad \frac{\mu(b \wedge a)}{\mu(a)} > \frac{\mu(b) - \mu(a \wedge b)}{1 - \mu(a)}.$$

Therefore if a commutes with b and $\mu(a \wedge b) > \mu(a)\mu(b)$, both a and b are always common causes of a and b in μ .

Definition 3.2 Let a , b and c be elements in an orthomodular lattice \mathcal{L} . If c is a common cause of a and b , and differs from both a and b , we call the element c a nontrivial common cause.

Definition 3.3 ([5, 6]) Let a and b be elements in an orthomodular lattice \mathcal{L} . a and b are called logically independent if the following conditions hold.

$$a \wedge b \neq 0, \quad a^\perp \wedge b \neq 0, \quad a \wedge b^\perp \neq 0, \quad a^\perp \wedge b^\perp \neq 0.$$

We examine whether a nontrivial common cause of two correlated elements in an atomless and complete orthomodular lattice exists or not (Proposition 3.9 and Proposition 3.10).

We prove the following lemma in reference to Lemma 3 in [8].

Lemma 3.4 Let μ be a completely additive probability measure on an orthomodular lattice \mathcal{L} and let a and b be elements in \mathcal{L} such that $\mu(a \wedge b) > \mu(a)\mu(b)$. Then $1 - \mu(a) - \mu(b) + \mu(a \wedge b) > 0$ and the following facts hold.

1. If $\mu(a) > \mu(a \wedge b)$, then

$$\frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} < \mu(a \wedge b).$$

2. If $\mu(a) = \mu(a \wedge b)$, then

$$\frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} = \mu(a \wedge b).$$

Proof Since $\mu(a \wedge b) > \mu(a)\mu(b)$, $\mu(a) < 1$ and $\mu(b) < 1$. Therefore

$$1 - \mu(a) - \mu(b) + \mu(a \wedge b) > 1 - \mu(a) - \mu(b) + \mu(a)\mu(b) = (1 - \mu(a))(1 - \mu(b)) > 0.$$

1. If $\mu(a) > \mu(a \wedge b)$, then

$$\begin{aligned} \mu(a)\mu(b) &= \mu(a)((\mu(b) - \mu(a \wedge b)) + \mu(a \wedge b)) \\ &= \mu(a)(\mu(b) - \mu(a \wedge b)) + \mu(a)\mu(a \wedge b) \\ &> \mu(a \wedge b)(\mu(b) - \mu(a \wedge b)) + \mu(a)\mu(a \wedge b) \\ &= (\mu(a) + \mu(b) - \mu(a \wedge b))\mu(a \wedge b). \end{aligned}$$

So $\mu(a)\mu(b)/\mu(a \wedge b) > \mu(a) + \mu(b) - \mu(a \wedge b)$, which implies

$$1 - \frac{\mu(a)\mu(b)}{\mu(a \wedge b)} < 1 - \mu(a) - \mu(b) + \mu(a \wedge b).$$

Because $1 - \mu(a) - \mu(b) + \mu(a \wedge b) > 0$,

$$\frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} < \mu(a \wedge b).$$

□

2. It can be proved in a similar way to the proof of 1.

We prove the following lemma in reference to Lemma 3 in [8].

Lemma 3.5 *Let μ be a completely additive probability measure on an orthomodular lattice \mathcal{L} and let a, b and c be elements in \mathcal{L} such that $\mu(a \wedge b) > \mu(a)\mu(b)$, $c \leq a \wedge b$ and*

$$\mu(c) = \frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)}.$$

Then c is a common cause of a and b in μ .

Proof Since $c \leq a$ and $c \leq b$, cCa and cCb ([4], Lemma (36.3)). Elementary algebraic calculation shows that

$$\mu(c) = \frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)}$$

is equivalent to

$$(1 - \mu(c))(\mu(a \wedge b) - \mu(c)) = (\mu(a) - \mu(c))(\mu(b) - \mu(c)).$$

Because $a \wedge b \geq c$ and $\mu(a \wedge b) \neq 1$, $\mu(c) \neq 1$. So,

$$\frac{\mu(a \wedge b) - \mu(c)}{1 - \mu(c)} = \frac{\mu(a) - \mu(c)}{1 - \mu(c)} \frac{\mu(b) - \mu(c)}{1 - \mu(c)}$$

Since $a \wedge b \geq c$, one can write

$$x = c + (x \wedge c^\perp) \quad (x = a \wedge b, a, b),$$

which implies

$$\mu(x) = \mu(c) + \mu(x \wedge c^\perp) \quad (x = a \wedge b, a, b).$$

Therefore (2) holds. Since $c \leq a \wedge b$, (1) holds trivially.

Since $\mu(a \wedge b) > \mu(a)\mu(b)$, $\mu(a) < 1$ and $\mu(b) < 1$, which implies $\mu(a) - \mu(c) < 1 - \mu(c)$ and $\mu(b) - \mu(c) < 1 - \mu(c)$. So

$$\frac{\mu(a \wedge c)}{\mu(c)} = \frac{\mu(c)}{\mu(c)} = 1 > \frac{\mu(a) - \mu(c)}{1 - \mu(c)} = \frac{\mu(a \wedge c^\perp)}{\mu(c^\perp)},$$

$$\frac{\mu(b \wedge c)}{\mu(c)} = \frac{\mu(c)}{\mu(c)} = 1 > \frac{\mu(b) - \mu(c)}{1 - \mu(c)} = \frac{\mu(b \wedge c^\perp)}{\mu(c^\perp)}.$$

Therefore (3) and (4) hold. □

Lemma 3.6 Let μ be a completely additive probability measure on an atomless and complete orthomodular lattice \mathcal{L} and let r be an element in \mathcal{L} such that $\mu(r) \neq 0$. For any real number α such that $0 < \alpha < \mu(r)$ there exists $c \in \mathcal{L}$ such that $c < r$ and $\mu(c) = \alpha$.

Proof Let \mathbb{S} be the set of all subsets of \mathcal{L} and define \mathbb{T} as

$$\left\{ T \in \mathbb{S} \mid \begin{array}{l} \text{For any finite elements } t_1, \dots, t_k \in T, \text{ they are mutually orthogonal,} \\ \sum_{i=1}^k \mu(t_i) \leq \alpha \text{ and } \sum_{i=1}^k t_i < r \end{array} \right\}.$$

\mathbb{T} is partially ordered by set inclusion. Since $\{0\} \in \mathbb{T}$, \mathbb{T} is not empty.

Let \mathbb{T}' be any linearly ordered subset of \mathbb{T} . For any finite elements t'_1, \dots, t'_l in $\bigcup \mathbb{T}'$, there exists $T' \in \mathbb{T}'$ such that $t'_1, \dots, t'_l \in T'$ since \mathbb{T}' is a linear ordered subset of \mathbb{T} . So t'_1, \dots, t'_l are mutually orthogonal elements such that $\sum_{i=1}^l \mu(t'_i) \leq \alpha$ and $\sum_{i=1}^l t'_i < r$. Therefore $\bigcup \mathbb{T}' \in \mathbb{T}$. By Zorn's lemma \mathbb{T} has a maximal element U . Since \mathcal{L} is complete, $\bigvee \{u | u \in U\}$ exists. Define c as $\bigvee \{u | u \in U\}$. Since $u < r$ for any element $u \in U$, $c \leq r$.

Let $\{u_a | a \in A\}$ denote U . Then

$$\mu(c) = \mu\left(\sum_{a \in A} u_a\right) = \sum_{a \in A} \mu(u_a).$$

Since $\sum_{a \in A'} \mu(u_a) \leq \alpha$ for any finite subset A' of A , $\mu(c) \leq \alpha$.

Suppose that $\mu(c) < \alpha$. Because $\mu(c) < \alpha < \mu(r)$ and $c \leq r$, $c < r$. So $c^\perp \wedge r \neq 0$. Let v denote $c^\perp \wedge r$. Define the countably infinite set $\{v_i\}_{i \in \mathbb{N}}$ of mutually orthogonal nonzero elements in \mathcal{L} such that $v > \sum_{i=1}^n v_i$ for any $n \in \mathbb{N}$ as follows. Since \mathcal{L} is atomless, there exists an element $v' \in \mathcal{L}$ such that $0 < v' < v$. Define v_1 as v' . If mutually orthogonal nonzero elements v_1, \dots, v_k such that $v > \sum_{i=1}^k v_i$ are defined, then

$$v = \sum_{i=1}^k v_i + \left(\left(\sum_{i=1}^k v_i \right)^\perp \wedge v \right)$$

and $(\sum_{i=1}^k v_i)^\perp \wedge v \neq 0$. There exists $v'' \in \mathcal{L}$ such that $(\sum_{i=1}^k v_i)^\perp \wedge v > v'' > 0$ since \mathcal{L} is atomless. Define v_{k+1} as v'' . Then

$$v > \sum_{i=1}^k v_i + v_{k+1} = \sum_{i=1}^{k+1} v_i.$$

Since $v_1^\perp \wedge \dots \wedge v_k^\perp = (\sum_{i=1}^k v_i)^\perp > v_{k+1}$, any elements in $\{v_1, \dots, v_{k+1}\}$ are mutually orthogonal. Because \mathcal{L} is complete, $\sum_{i=1}^\infty v_i$ exists and $v \geq \sum_{i=1}^\infty v_i$.

If $\mu(v_i) \geq \alpha - \mu(c)$ for any $i \in \mathbb{N}$, $\lim_{n \rightarrow \infty} \sum_{i=1}^n \mu(v_i) \rightarrow \infty$. Because $\mu(v) \geq \sum_{i=1}^\infty \mu(v_i)$, this is a contradiction. So there is $v_k \in \{v_i\}_{i \in \mathbb{N}}$ such that $\mu(v_k) < \alpha - \mu(c)$, which implies $\mu(c + v_k) < \alpha$. Therefore for any finite elements u'_1, \dots, u'_m in $U \cup \{v_k\}$, they are mutually orthogonal, $\sum_{i=1}^m \mu(u'_i) \leq \alpha$ and $\sum_{i=1}^m u'_i < r$. Therefore $U \cup \{v_k\} \in \mathbb{T}$. This is a contradiction because $v_k \notin U$ and U is a maximal element of \mathbb{T} . Therefore $\mu(c) = \alpha$. \square

We will prove the following lemma in reference to the proof of Proposition 7 in [1].

Lemma 3.7 Let μ be a faithful completely additive probability measure on an atomless and complete orthomodular lattice \mathcal{L} , let r be a nonzero element in \mathcal{L} and let α be a real number such that $0 < \alpha < \mu(r)$. Then the set $\{c \in \mathcal{L} | c < r \text{ and } \mu(c) = \alpha\}$ is an uncountably infinite set.

Proof By Lemma 3.6, there exists $c \in \mathcal{L}$ such that $\mu(c) = \alpha$ and $c < r$. Since $c < r$, $r \wedge c^\perp \neq 0$, that is, $\mu(r \wedge c^\perp) \neq 0$. Let β be any real number such that $0 < \beta < \min\{\mu(r \wedge c^\perp), \alpha\}$. By Lemma 3.6, there exist $c_{\beta 1}, c_{\beta 2} \in \mathcal{L}$ such that $c_{\beta 1} < c$, $c_{\beta 2} < r \wedge c^\perp$ and $\mu(c_{\beta 1}) = \mu(c_{\beta 2}) = \beta$.

Define c_β as $(c \wedge c_{\beta 1}^\perp) + c_{\beta 2}$. Then $c_\beta < c \vee (r \wedge c^\perp) < r$ and

$$\mu(c_\beta) = \mu(c \wedge c_{\beta 1}^\perp) + \mu(c_{\beta 2}) = \mu(c) - \mu(c_{\beta 1}) + \mu(c_{\beta 2}) = \alpha.$$

For any real numbers β and γ such that $0 < \beta < \gamma < \min\{\mu(r \wedge c^\perp), \alpha\}$, $c_{\beta 2} \neq c_{\gamma 2}$ since μ is a faithful completely additive probability measure. Therefore $c_\beta \neq c_\gamma$. \square

Reichenbach [9] proved the following lemma (pp. 159–160).

Lemma 3.8 Let μ be a completely additive probability measure on an orthomodular lattice \mathcal{L} and let a, b and c be elements in \mathcal{L} . If c is a common cause of a and b in μ , then $\mu(a \wedge b) > \mu(a)\mu(b)$.

Proof Since aCc and bCc , $a \wedge bCc$ ([4], Lemma (36.4)). So $\mu(a) = \mu(a \wedge c) + \mu(a \wedge c^\perp)$, $\mu(b) = \mu(b \wedge c) + \mu(b \wedge c^\perp)$ and $\mu(a \wedge b) = \mu(a \wedge b \wedge c) + \mu(a \wedge b \wedge c^\perp)$ by Lemma 2.9. Then

$$\mu(a \wedge b) - \mu(a)\mu(b) = \mu(c)(1 - \mu(c)) \left(\frac{\mu(a \wedge c)}{\mu(c)} - \frac{\mu(a \wedge c^\perp)}{\mu(c^\perp)} \right) \left(\frac{\mu(b \wedge c)}{\mu(c)} - \frac{\mu(b \wedge c^\perp)}{\mu(c^\perp)} \right)$$

by (1) and (2). By (3) and (4), $\mu(a \wedge b) > \mu(a)\mu(b)$. \square

Proposition 3.9 Let μ be a completely additive probability measure on an atomless and complete orthomodular lattice \mathcal{L} and let a and b be elements in \mathcal{L} such that $a \wedge b^\perp \neq 0$ and $a^\perp \wedge b \neq 0$. Then the following conditions are equivalent.

1. $\mu(a \wedge b) > \mu(a)\mu(b)$
2. There exists a nontrivial common cause of a and b in μ .

Proof $2 \Rightarrow 1$ It holds by Lemma 3.8.

$1 \Rightarrow 2$ If $\mu(a \wedge b) = \mu(a)$,

$$\frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} = \mu(a \wedge b)$$

by Lemma 3.4. By Lemma 3.5 $a \wedge b$ is a common cause of a and b in μ . If $a \wedge b = a$ or $a \wedge b = b$, $a \wedge b^\perp = 0$ or $a^\perp \wedge b = 0$. This contradicts the assumption. So $a \wedge b$ is a nontrivial common cause of a and b in μ .

If $\mu(a \wedge b) \neq \mu(a)$,

$$\frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} < \mu(a \wedge b)$$

by Lemma 3.4. By Lemma 3.6 there exists $c \in \mathcal{L}$ such that

$$c < a \wedge b \text{ and } \mu(c) = \frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)}.$$

By Lemma 3.5 c is a nontrivial common cause of a and b in μ . \square

Proposition 3.10 *Let μ be a faithful completely additive probability measure on an atomless and complete orthomodular lattice \mathcal{L} and let a and b be elements in \mathcal{L} such that $a \wedge b^\perp \neq 0$. Then the following conditions are equivalent.*

1. $\mu(a \wedge b) > \mu(a)\mu(b)$
2. *There exist uncountably infinite nontrivial common causes of a and b in μ .*

Proof 2 \Rightarrow 1 It holds by Lemma 3.8.

1 \Rightarrow 2 If $\mu(a \wedge b) = \mu(a)$, $a \wedge b = a$ by Lemma 2.9. Then $a \wedge b^\perp = a \wedge b \wedge b^\perp = 0$. This contradicts the assumption. So $\mu(a \wedge b) < \mu(a)$. By Lemma 3.4

$$\frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} < \mu(a \wedge b).$$

By Lemma 3.7 the set

$$\left\{ c \in \mathcal{L} \mid c < a \wedge b \text{ and } \mu(c) = \frac{\mu(a \wedge b) - \mu(a)\mu(b)}{1 - \mu(a) - \mu(b) + \mu(a \wedge b)} \right\}$$

is an uncountably infinite set. By Lemma 3.5 the element in the set is a nontrivial common cause of a and b in μ . \square

4 Concluding Remarks

The common cause of a and b in Proposition 3.9 is a lower bound of $a \wedge b$. Redei [7] called such a common cause a strong common cause (p. 1312). There is another common cause c such that $c \not\leq a$ and $c \not\leq b$. Redei [7] called such a common cause a genuinely probabilistic common cause.

If $\mu(a \wedge b) > \mu(a)\mu(b)$ implies $1 - \mu(a)\mu(b)/\mu(a \wedge b) < \mu(a^\perp \wedge b^\perp)$ in Lemma 3.4, it can be proved that there exists a genuinely probabilistic common cause of two correlated elements in an atomless and complete orthomodular lattice as well as the proof of Proposition 3.9 (cf. [1], Proposition 7). When a commutes with b , $\mu(a \wedge b) > \mu(a)\mu(b)$ implies $1 - \mu(a)\mu(b)/\mu(a \wedge b) < \mu(a^\perp \wedge b^\perp)$. But it is not obvious whether the inequality holds in the case where a does not commute with b .

Acknowledgement The author wishes to thank Masanao Ozawa for helpful comments on an earlier draft.

References

1. Gyenis, B., Redei, M.: When can statistical theories be causally closed? Found. Phys. **34**, 1285–1303 (2004)
2. Halvorson, H., Clifton, R.: Generic bell correlation between arbitrary local algebras in quantum field theory. J. Math. Phys. **41**, 1711–1717 (2001)

3. Hofer-Szabo, G., Redei, M., Szabo, L.E.: Common cause completness of classical and quantum probability spaces. *Int. J. Theor. Phys.* **39**, 913–919 (2000)
4. Maeda, M., Maeda, S.: Theory of Symmetric Lattices. Springer, Berlin (1970)
5. Redei, M.: Logical independence in quantum logic. *Found. Phys.* **25**, 411–422 (1995)
6. Redei, M.: Logically independent von Neumann lattices. *Int. J. Theor. Phys.* **34**, 1711–1718 (1995)
7. Redei, M.: Reichenbach's common cause principle and quantum field theory. *Found. Phys.* **27**, 1309–1321 (1997)
8. Redei, M., Summers, S.J.: Local primitive causality and the common cause principle in quantum field theory. *Found. Phys.* **32**, 335–355 (2002)
9. Reichenbach, H.: The Direction of Time. Dover, New York (1956)
10. Svozil, K., Redei, M., Summers, S.J.: Remarks on causality in relativistic quantum field theory. *Int. J. Theor. Phys.* **44**, 1029–1039 (2005)