

# Kaluza-Klein Type Robertson Walker Cosmological Model with Dynamical Cosmological Term $\Lambda$

G.S. Khadekar · Vrushali Patki

Received: 23 September 2007 / Accepted: 9 November 2007 / Published online: 27 November 2007  
© Springer Science+Business Media, LLC 2007

**Abstract** In this paper we have analyzed the Kaluza-Klein type Robertson Walker (RW) cosmological models by considering three different forms of variable  $\Lambda$ :  $\Lambda \sim (\frac{\dot{a}}{a})^2$ ,  $\Lambda \sim (\frac{\ddot{a}}{a})$  and  $\Lambda \sim \rho$ . It is found that, the connecting free parameters of the models with cosmic matter and vacuum energy density parameters are equivalent, in the context of higher dimensional space time. The expression for the look back time, luminosity distance and angular diameter distance are also derived. This work has thus generalized to higher dimensions the well-known results in four dimensional space time. It is found that there may be significant difference in principle at least, from the analogous situation in four dimensional space time.

**Keywords** Higher dimensional space time · Cosmology · Cosmological parameters · Cosmological tests

## 1 Introduction

Higher dimensional theories of Kaluza-Klein (KK)-type have been considered to study some aspects of early universe [1–4]. In such KK-theory it has been assumed that the extra dimension form a compact manifold of very small size undetectable at present day energies. Thus in such higher dimensional theories one would expect that at the grand unification scale the word manifold has more than one dimension. It is widely believed that a consistent unification of all fundamental forces in nature would be possible within the space-time with an extra dimensions beyond those four observed so far. The absences of any signature of extra dimension in current experiments is usually explained the compactness of extra dimensions. The idea of dimensional reduction or self compactification fits in particularly well in cosmology because if we believe in the big bang, our universe was much smaller at the early

---

G.S. Khadekar (✉) · V. Patki  
Department of Mathematics, R.T.M. Nagpur University, Mahatma Jyotiba Phule Educational Campus,  
Amravati Road, Nagpur 440 033, India  
e-mail: [gkhadekar@yahoo.com](mailto:gkhadekar@yahoo.com)

V. Patki  
e-mail: [patki\\_vrushali@rediffmail.com](mailto:patki_vrushali@rediffmail.com)

stage and the present four dimensional stage could have been preceded by a higher dimensional one [1]. In this work we consider five dimensional Robertson Walker (RW) model as a test case. In RW type of homogeneous cosmological model, the dimensionality has a marked effect on the time temperature relation of the universe and our universe appears to cool more slowly in higher dimensional space time [5].

In recent year, models with a relic cosmological constant  $\Lambda$  have received considerable attention among researchers for various reasons ([6–11] and references therein). We should realize that the existence of a nonzero cosmological constant in Einstein's equations is a feature of deep and profound consequence. The recent observations indicate that  $\Lambda \sim 10^{-55} \text{ cm}^{-2}$  while particle physics prediction for  $\Lambda$  is greater than this value by a factor of order  $10^{120}$ . This discrepancy is known as cosmological constant problem. Some of the recent discussions on the cosmological constant “problem” and consequence on cosmology with a time-varying cosmological constant are investigated by Ratra and Peebles [12], Dolgov [13, 14], Dolgov et al. [15], Sahni and Starobinsky [16], Padmanabhan [17] and Peebles [18]. For earlier reviews on this topic, the reader is referred to Zeldovich [19], Weinberg [20] and Carrol et al. [21].

Recent observations of type Ia supernovae (SNe Ia) at redshift  $z < 1$  provide startling and puzzling evidence that the expansion of the universe at the present time appears to be *accelerating*, behaviour attributed to “dark energy” with negative pressure. These observations [22–29] strongly favour a significant and positive value of  $\Lambda$ . The main conclusion of these observations is that the expansion of the universe is accelerating.

A number of authors have argued in favour of the dependence  $\Lambda \sim t^{-2}$  first expressed by Bertolami [30, 31] and later on by several authors Berman [32–34], Berman and Som [35], Berman et al. [36], Berman and Gomide [37], Beesham [38, 39], Singh et al. [40], Pradhan and Kumar [41], Pradhan and Yadav [42], Gasperini [43, 44], Peebles and Ratra [45] in different context. Recently, motivated by dimensional grounds in keeping with quantum cosmology, Chen and Wu [46], Abdel-Rahaman [47, 48] considered a  $\Lambda$  varying as  $a^{-2}$ , where  $a$  is a scale factor. Carvalho et al. [49], Waga [50], Silveira and Waga [51], Vishwakarma [52] have also considered/modified the same kind of variation. Such a dependence alleviates some problems in reconciling observational data with the inflationary universe scenario. Al-Rawaf and Taha [53] and Al-Rawaf [54] and Overdin and Cooperstock [55] proposed a cosmological model with a cosmological constant of the form  $\Lambda = \beta \frac{\ddot{a}}{a}$ , where  $\beta$  is a constant. Following the same decay law recently Arbab [56, 57] have investigated cosmic acceleration with positive cosmological constant and also analyze the implication of a model built-in cosmological constant for four-dimensional space time. The cosmological consequences of this decay law are very attractive. This law provides reasonable solutions to the cosmological puzzles presently known. One of the motivations for introducing  $\Lambda$  term is to reconcile the age parameter and the density parameter of the universe with recent observational data.

Vishwakarma [58] has studied the magnitude-redshift relation for the type Ia supernovae data and the angular size-redshift relation for the updated compact radio sources data [59] by considering four variable  $\Lambda$ -models:  $\Lambda \sim a^{-2}$ ,  $\Lambda \sim H^{-2}$ ,  $\Lambda \sim \rho$  and  $\Lambda \sim t^{-2}$ . Recently Ray and Mukhopadhyay [60] have solved Einstein's equations for specific dynamical models of the cosmological term  $\Lambda$  in the form:  $\Lambda \sim (\frac{\dot{a}}{a})^2$ ,  $\Lambda \sim (\frac{\ddot{a}}{a})$  and  $\Lambda \sim \rho$  and shown that the models are equivalent in the framework of flat RW space time.

In this context, the aim of the present work is based on recent available observational information. In this paper the implication of cosmological model with cosmological term of three different forms:  $\Lambda = \alpha (\frac{\dot{a}}{a})^2$ ,  $\Lambda = \beta (\frac{\ddot{a}}{a})$  and  $\Lambda = 8\pi\gamma\rho$ , where  $\alpha$ ,  $\beta$  and  $\gamma$  are free parameters, are analyzed within the framework of higher dimensional space time. The paper is

organized as follows. The Einstein’s field equations and their general solutions for different  $\Lambda$ -dependent models are presented in Sects. 2 and 3 respectively. The equivalence of different forms of dynamical cosmological term  $\Lambda$  are discussed in Sect. 4. The cosmological tests pertaining neoclassical test, luminosity distance, angular diameter distance and look back time for the model are discussed in Sects. 5–8. As a concluding part some discussions are made in Sect. 9.

## 2 The Metric and Field Equations

We consider the Kaluza-Klein type Robertson Walker (RW) space time

$$ds^2 = dt^2 - a^2(t) \left[ \frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) + (1 - kr^2)d\psi^2 \right], \tag{1}$$

where  $k = 0, \pm 1$  is the curvature parameter.

The usual energy-momentum tensor is modified by addition of a term

$$T_{ij}^{vac} = -\Lambda(t)g_{ij}, \tag{2}$$

where  $\Lambda(t)$  is the cosmological term and  $g_{ij}$  is the metric tensor.

Einstein’s field equations (in gravitational units  $c = 1, G = 1$ ) read as

$$R_{ij} - \frac{1}{2}Rg_{ij} = -8\pi T_{ij} - \Lambda(t)g_{ij}. \tag{3}$$

The energy-momentum tensor  $T_{ij}$  in the presence of a perfect fluid has the form

$$T_{ij} = (p + \rho)u_i u_j - pg_{ij}, \tag{4}$$

where  $p$  and  $\rho$  are, respectively, the energy and pressure of the cosmic fluid, and  $u_i$  is the fluid five-velocity such that  $u^i u_i = 1$ .

The Einstein field equations (3) with (4) for the metric (1) take the form

$$6 \left( \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) = 8\pi\rho + \Lambda(t), \tag{5}$$

$$\frac{3\ddot{a}}{a} + 3 \left( \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) = -8\pi p + \Lambda(t). \tag{6}$$

An over dot indicates a derivative with respect to time  $t$ . The energy conservation law can be written as

$$\dot{\rho} + 4(\rho + p)H = -\frac{\dot{\Lambda}}{8\pi}, \tag{7}$$

where  $H = \frac{\dot{a}}{a}$ , is the Hubble parameter.

For complete determinacy of the system, we consider the barotropic equation of state

$$p = \omega\rho, \tag{8}$$

where equation of state parameter  $\omega$  can take the constant values 0, 1/3,  $-1$  and  $+1$  respectively for the dust, radiation, vacuum fluid and stiff fluid.

By using (8) and (5), (6) reduces to

$$\frac{\ddot{a}}{a} + \frac{4\pi\rho}{3}(2\omega + 1) = \frac{\Lambda}{6}. \tag{9}$$

By using (5), (6) and (8) for eliminating  $\rho$ , we obtain

$$\frac{3\ddot{a}}{a} + 3(1 + 2\omega) \left( \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) = (1 + \omega)\Lambda(t). \tag{10}$$

This is the dynamical equation related to the cosmic scale factor  $a$  for a known value of the dynamical cosmological term  $\Lambda$ . It can be readily be observed from the above equations (9) and (10) that  $\Lambda$  is dependent on the factor  $\frac{\ddot{a}}{a}$ ,  $\rho$ ,  $(\frac{\dot{a}}{a})^2$  and  $a^{-2}$  in specific way. However, inflationary theory of universe predicts and CMB detectors such as BOOMERANG [61–63], MAXIMA [64–66], DASI [67], CBI [68] and WMAP [69, 70] confirm that the universe is spatially flat. Therefore,  $\Lambda \sim a^{-2}$  case does not sustain for  $k = 0$  and therefore omitted here. Hence we shall consider the cosmological term related to the cases  $(\frac{\dot{a}}{a})^2$ ,  $\frac{\ddot{a}}{a}$  and  $\rho$  and obtained the solutions in the following sections.

### 3 Three Different Forms of $\Lambda$

#### 3.1 Model for $\Lambda \sim (\frac{\dot{a}}{a})^2$

If we use  $\Lambda = \alpha(\frac{\dot{a}}{a})^2 = \alpha H^2$ , where  $\alpha$  is a constant, then for flat universe  $k = 0$ , (10) reduces to

$$a\ddot{a} + \frac{1}{3} [\omega(6 - \alpha) + (3 - \alpha)] \dot{a}^2 = 0. \tag{11}$$

Integrating (11), we get general solution as

$$a(t) = \left[ \frac{1}{3} (6 - \alpha)(1 + \omega)k_1 t \right]^{\frac{3}{(6-\alpha)(1+\omega)}}. \tag{12}$$

Then from (5)

$$\rho(t) = \frac{9}{8\pi(6 - \alpha)(1 + \omega)^2} \frac{1}{t^2}, \tag{13}$$

$$\Lambda(t) = \frac{9\alpha}{(6 - \alpha)^2(1 + \omega)^2} \frac{1}{t^2}, \tag{14}$$

where  $k_1$  is an integrating constant.

From the above set of equations (12–14) it is to be noted that  $\alpha \neq 6$  for physical validity. From (13), it is observed that  $\rho > 0$  implies  $0 < \alpha < 6$ . The case  $\alpha \geq 6$  is either unphysical or not compatible with time dependent  $\Lambda$ . As  $\Lambda \neq 0$ , then  $\dot{a} \neq 0$ , which suggests that when  $\Lambda \sim (\frac{\dot{a}}{a})^2$ , then as long as  $\Lambda \neq 0$ , the expansion of the universe stops.

### 3.2 Model for $\Lambda \sim \frac{\ddot{a}}{a}$

If we consider  $\Lambda = \beta \frac{\ddot{a}}{a}$ , where  $\beta$  is a constant, then for flat universe  $k = 0$ , (10) reduces to

$$a\ddot{a} + \frac{3(1 + 2\omega)}{(3 - \omega\beta - \beta)}\dot{a}^2 = 0. \tag{15}$$

Equation (15) gives the general solution as

$$a(t) = \left[ \frac{(1 + \omega)(6 - \beta)}{(3 - \omega\beta - \beta)} k_2 t \right]^{\frac{(3 - \omega\beta - \beta)}{(1 + \omega)(6 - \beta)}}, \tag{16}$$

$$\Lambda(t) = \left[ \frac{3\beta(1 + 2\omega)(\beta + \omega\beta - 3)}{(1 + \omega)^2(\beta - 6)^2} \right] \frac{1}{t^2}, \tag{17}$$

$$\rho(t) = \frac{3(\beta + \omega\beta - 3)}{8\pi(\beta - 6)(1 + \omega)^2} \frac{1}{t^2}, \tag{18}$$

where  $k_2$  is an integrating constant.

It is remarkable to mention here from (16–18) that for physical valid solutions  $\beta \neq 0$  and also  $\beta \neq 6$ . It is also observed from above equations that for  $\beta > 3$ ,  $\Lambda > 0$  and for  $\beta > 6$ ,  $\rho > 0$  for dust model. Thus, either  $\beta < 0$  or  $\beta > 6$  for our present model.

### 3.3 Model for $\Lambda \sim \rho$

If we set  $\Lambda = 8\pi\gamma\rho$ , where  $\gamma$  is a free parameter, then using (5), (10) reduces to

$$a\ddot{a} - \left( \frac{\gamma - 2\omega - 1}{1 + \gamma} \right) \dot{a}^2 = 0 \tag{19}$$

for a flat universe ( $k = 0$ ).

Solving (19) we get the set of general solution as

$$a(t) = \left[ \frac{2(1 + \omega)k_3}{(1 + \gamma)} t \right]^{\frac{(1 + \gamma)}{2(1 + \omega)}}, \tag{20}$$

$$\rho(t) = \frac{3(1 + \gamma)}{16\pi(1 + \omega)^2} \frac{1}{t^2}, \tag{21}$$

$$\Lambda(t) = \frac{3\gamma(1 + \gamma)}{2(1 + \omega)^2} \frac{1}{t^2}, \tag{22}$$

where  $k_3$  is an integrating constant. In this case for physical valid solution  $\gamma > 0$ .

## 4 Equivalence of Three Forms of $\Lambda$

Now, let us find out the interrelations between  $\alpha$ ,  $\beta$  and  $\gamma$  and hence the equivalence of the different forms of the dynamical cosmological terms  $\Lambda \sim (\frac{\ddot{a}}{a})^2$ ,  $\Lambda \sim (\frac{\ddot{a}}{a})$  and  $\Lambda \sim \rho$ .

From (12), after differentiating it and the dividing by  $a$ , we get

$$t = \frac{3}{(6 - \alpha)(1 + \omega)H}, \tag{23}$$

where  $H$  is the Hubble parameter as mentioned earlier. Therefore for specific values of  $\alpha$  and  $\omega$ , (23) shows that  $H \sim t^{-1}$ .

Using (23) in (13) and also from the definition of the cosmic matter-density parameter  $\Omega (= 8\pi\rho/6H^2)$ , we get

$$\Omega_{m\alpha} = 1 - \frac{\alpha}{6}, \tag{24}$$

where  $\Omega_{m\alpha}$  is the cosmic matter energy density parameter for the  $\alpha$ -related dynamic  $\Lambda$ -model.

Using (23) in (14) and also from the definition of the cosmic vacuum-energy density parameter  $\Omega_\Lambda = \Lambda/6H^2$ , we have

$$\Omega_{\Lambda\alpha} = \frac{\alpha}{6}, \tag{25}$$

where, in the similar fashion,  $\Omega_{\Lambda\alpha}$  is the cosmic vacuum-energy density parameter for the  $\alpha$ -related dynamic  $\Lambda$ -model.

From (24) and (25), we obtain

$$\Omega_{m\alpha} + \Omega_{\Lambda\alpha} = 1, \tag{26}$$

which is the relation between the cosmic matter- and vacuum-energy density parameters for a flat ( $k = 0$ ) universe.

Similarly from (16–18) we obtain

$$\Omega_{m\beta} = \frac{(\beta - 6)}{2(\beta + \omega\beta - 3)}, \tag{27}$$

$$\Omega_{\Lambda\beta} = \frac{\beta(1 + 2\omega)}{2(\beta + \omega\beta - 3)}, \tag{28}$$

where  $\Omega_{m\beta}$  and  $\Omega_{\Lambda\beta}$  are respectively the cosmic matter and vacuum energy density parameters for the  $\beta$ -related dynamic  $\Lambda$ -model.

Adding (27) and (28), we get

$$\Omega_{m\beta} + \Omega_{\Lambda\beta} = 1, \tag{29}$$

which is again the relation between the cosmic matter- and vacuum-energy density parameters for a flat ( $k = 0$ ) universe.

In the same manner from (20–22) the cosmic matter and vacuum energy density parameter  $\Omega_{m\gamma}$  and  $\Omega_{\Lambda\gamma}$  respectively for the model  $\Lambda \sim \rho$  also satisfy the relation

$$\Omega_{m\gamma} + \Omega_{\Lambda\gamma} = 1, \tag{30}$$

where

$$\gamma = \frac{\Omega_{\Lambda\gamma}}{\Omega_{m\gamma}}. \tag{31}$$

Thus, from the relations (26), (29) and (30) without any loss of generality we can set

$$\Omega_{m\alpha} = \Omega_{m\beta} = \Omega_{m\gamma} = \Omega_m, \tag{32}$$

$$\Omega_{\Lambda\alpha} = \Omega_{\Lambda\beta} = \Omega_{\Lambda\gamma} = \Omega_\Lambda, \tag{33}$$

where  $\Omega_m$  and  $\Omega_\Lambda$  are respectively the cosmic matter and vacuum energy density parameters which in absence of any curvature satisfy the general relation

$$\Omega = \Omega_m + \Omega_\Lambda = 1. \tag{34}$$

Now from (25) and (33) we obtain

$$\alpha = 6\Omega_\Lambda. \tag{35}$$

Also dividing (27) by (28) and using (32) and (33)  $\beta$  is obtained as

$$\beta = \frac{6\Omega_\Lambda}{\Omega_\Lambda - (1 + 2\omega)\Omega_m}. \tag{36}$$

Similarly from (31) after using (32) and (33)

$$\gamma = \frac{\Omega_\Lambda}{\Omega_m}. \tag{37}$$

Hence from (34) we get

$$\Omega = (1 + \gamma)\Omega_m = 1, \tag{38}$$

which is an other relation of total cosmic energy density in the case of flat universe.

All the above general relations of  $\alpha$ ,  $\beta$  and  $\gamma$  in terms of  $\Omega_m$  and  $\Omega_\Lambda$  also hold for the particular cases of dust ( $\omega = 0$ ) and radiation ( $\omega = 1/4$ ). It is interesting to note that while relation of  $\alpha$  and  $\gamma$  with the cosmic matter and the vacuum-energy density parameters are independent of  $\omega$ , relation of  $\beta$  with  $\Omega_m$  and  $\Omega_\Lambda$  depends on  $\omega$ .

It can easily be shown that the particular solution of  $\Lambda \sim (\frac{\dot{a}}{a})^2$  model for dust and radiation cases become identical with their corresponding counterparts for the other models in term of the dependency of  $a$ ,  $\rho$  and  $\Lambda$  on time  $t$  when expressed in terms of  $\Omega_m$  and  $\Omega_\Lambda$ . Therefore, these results imply that in view of  $\Omega_m$  and  $\Omega_\Lambda$  there are no distinctive features so that one can differentiate between the different forms of dynamic cosmological model viz.  $\Lambda \sim (\frac{\dot{a}}{a})^2$ ,  $\Lambda \sim (\frac{\ddot{a}}{a})$  and  $\Lambda \sim \rho$ . Thus, starting from any one of our  $\Lambda$ -models, since they are equivalent, we can arrive at the other relations.

Now from (35), after using (28) and (33), one can obtain

$$\alpha = \frac{3\beta(1 + 2\omega)}{\beta + \omega\beta - 3}. \tag{39}$$

Also from (35), after using (37) and (38), we get

$$\alpha = \frac{6\gamma}{1 + \gamma}. \tag{40}$$

Thus from (39) and (40), we find that the parameters involve in three dynamical relations are connected by

$$\alpha = \frac{3\beta(1 + 2\omega)}{\beta + \omega\beta - 3} = \frac{6\gamma}{1 + \gamma}. \tag{41}$$

It is observed from above equation that the three forms  $\Lambda = \alpha(\frac{\dot{a}}{a})^2$ ,  $\Lambda = \beta\frac{\ddot{a}}{a}$  and  $\Lambda = 8\pi\gamma\rho$  are equivalent and the three parameters  $\alpha$ ,  $\beta$  and  $\gamma$  are connected by the relation (41), which indicates that it is possible to find out the identical physical features of others if any one of the phenomenological  $\Lambda$  relation is known and hence these relations are not independent of each other. Also we can see that for the dust case (41), for the relation between  $\alpha$  and  $\beta$ , gives

$$\alpha = \frac{3\beta}{(\beta - 3)}. \tag{42}$$

In cosmology there are many ways to specify the distance between two points, because in the expanding universe, the distance between comoving objects are constantly changing and earth-bound observers look back in time as they look out in distance. In the following sections we discussed some cosmological distances for the case  $\Lambda \sim (\frac{\dot{a}}{a})^2$ .

**5 Neoclassical Tests (Proper Distance  $d(z)$ )**

A photon emitted by a source with coordinate  $r = r_1$  and  $t = t_1$  and received at a time  $t_0$  by an observer located at  $r = 0$ . The emitted radiation will follow null geodesics on which  $(\theta, \phi, \psi)$  are constant.

The proper distance between the source and observer is given by

$$d(z) = a_0 \int_a^{a_0} \frac{da}{a\dot{a}}, \tag{43}$$

$$r_1 = \int_{t_1}^{t_0} \frac{dt}{a} = \frac{a_0^{-1}H_0^{-1}A_0}{(1 - A_0)} [1 - (1 + z)^{\frac{A_0-1}{A_0}}].$$

Hence

$$d(z) = r_1 a_0 = H_0^{-1} \left( \frac{A_0}{1 - A_0} \right) [1 - (1 + z)^{\frac{A_0-1}{A_0}}], \tag{44}$$

where  $(1 + z) = \frac{a_0}{a}$  = redshift and  $a_0$  is the present scale factor of the universe and

$$A_0 = \frac{3}{(6 - \alpha)(1 + \omega)}.$$

For small value  $z$ , (44) reduces to

$$H_0 d(z) = z - \frac{1}{2A_0} z^2 + \dots \tag{45}$$

By using (12) the deceleration parameter can be written as

$$q = -\frac{a\ddot{a}}{\dot{a}^2} = -\left( \frac{A_0 - 1}{A_0} \right).$$

With the use of above equation, (45) can be written as

$$H_0 d(z) = z - \frac{1}{2}(1 + q)z^2 + \dots \tag{46}$$



From (44), it is observed that the distance  $d$  is maximum at  $z = \infty$ . Hence

$$d(z = \infty) = H_0^{-1} \left( \frac{A_0}{1 - A_0} \right). \quad (47)$$

## 6 Luminosity Distance

Luminosity distance is the another important concept of theoretical cosmology of a light source. The luminosity distance is a way of expanding the amount of light received from a distant object. It is the distance that the object appears to have, assuming the inverse square law for the reduction of light intensity with distance holds. The luminosity distance is *not* the actual distance to the object, because in the real universe the inverse square law does not hold. It is broken both because the geometry of the universe need not be flat, and because the universe is expanding. In other words, it is defined in such a way as generalizes the inverse-square law of the brightness in the static Euclidean space to an expanding curved space [50].

If  $d_L$  is the luminosity distance to the object, then

$$d_L = \left( \frac{L}{4\pi l} \right)^{\frac{1}{2}}, \quad (48)$$

where  $L$  is the total energy emitted by the source per unit time,  $l$  is the apparent luminosity of the object. Therefore one can write

$$d_L = d(1 + z). \quad (49)$$

Using (44), (49) reduces to

$$H_0 d_L = (1 + z) \left( \frac{A_0}{1 - A_0} \right) \left[ 1 - (1 + z)^{\frac{A_0 - 1}{A_0}} \right]. \quad (50)$$

For small  $z$ , (50) gives

$$H_0 d_L = z + \frac{1}{2}(1 - q)z^2 + \dots \quad (51)$$

or by using (24) and (32), (51) becomes

$$H_0 d_L = z + [1 - (1 + \omega)\Omega_m]z^2 + \dots \quad (52)$$

The luminosity distance depends on the cosmological model we have under discussion, and hence can be used to tell us which cosmological model describe our universe. Unfortunately, however, the observable quantity is the radiation flux density received from an object, and this can only be translated into a luminosity distance if the absolute luminosity of the object is known. This problem can however be circumvented if there are a population of objects at different distances which are believed to have the same luminosity; even if that luminosity is not known, it will appear merely as an overall scaling factor.

### 7 Angular Diameter Distance

The angular diameter distance  $d_A$  is defined as the ratio of an object’s physical transverse size to its angular size. It is used to convert angular separations in telescope images into proper separations at the sources.

The angular diameter  $d_A$  of a light source of proper distance  $d$  is given by

$$d_A = d(z)(1 + z)^{-1} = d_L(1 + z)^{-2}. \tag{53}$$

Applying (50) we obtain

$$H_0 d_A = \frac{A_0}{1 - A_0} \left[ \frac{1 - (1 + z)^{\frac{A_0 - 1}{A_0}}}{(1 + z)} \right]. \tag{54}$$

Usually  $d_A$  has a minimum (or maximum) for some  $Z = Z_m$ .

The angular diameter and luminosity distances have similar forms, but have a different dependence on redshift. As with the luminosity distance, for nearby objects the angular diameter distance closely matches the physical distance, so that objects appear smaller as they are put further away. However the angular diameter distance has a much more striking behaviour for distant objects. The luminosity distance effect dims the radiation and the angular diameter distance effect means the light is spread over a large angular area. This is so-called surface brightness dimming is therefore a particularly strong function of redshift.

### 8 Look Back Time

The time in the past at which the light we now receive from a distant object was emitted is called the look back time. How long ago the light was emitted (the look back time) depends on the dynamics of the universe.

The radiation travel time (or lookback time)  $(t - t_0)$  for photon emitted by a source at instant  $t$  and received at  $t_0$  is given by

$$t - t_0 = \int_a^{a_0} \frac{da}{\dot{a}}. \tag{55}$$

Equation (12) can be rewritten as

$$a = B_0 t^{A_0}, \quad B_0 = \text{constant}. \tag{56}$$

This follows that

$$\frac{a_0}{a} = 1 + z = \left( \frac{t_0}{t} \right)^{A_0}. \tag{57}$$

The above equation gives

$$t = t_0(1 + z)^{-\frac{1}{A_0}}. \tag{58}$$

This equation can also be expressed as

$$H_0(t_0 - t) = A_0[1 - (1 + z)^{-\frac{1}{A_0}}]. \tag{59}$$

For small  $z$  one obtain

$$H_0(t_0 - t) = z - \left(1 + \frac{q}{2}\right)z^2 + \dots \tag{60}$$

From (58) and (59), we observe that at  $z \rightarrow \infty$ ,  $H_0t_0 = A_0$  (constant).

### 9 Discussion

In this paper we have analyzed KK type RW cosmological models by considering three different forms of variable  $\Lambda$ . The equivalence of those forms is shown in terms of solutions thus obtained. Since  $\frac{\ddot{a}}{a} = \dot{H} + H^2$  then  $\Lambda \sim \frac{\ddot{a}}{a}$  models can be thought of as a combination of two models viz.  $\Lambda \sim \dot{H}$ ,  $\Lambda \sim H^2$ , we observe that  $\Lambda \sim \frac{\ddot{a}}{a}$  and  $\Lambda \sim \frac{\dot{a}^2}{a^2}$  models becomes identical when  $\dot{H} = 0$ . Now  $\dot{H} = 0$  implies  $H$  is constant. In this case we get exponential expansion and hence an inflationary scenario. Thus the idea of inflation is inherent in phenomenological model  $\Lambda \sim \frac{\ddot{a}}{a}$ . Moreover  $\Lambda \sim \frac{\dot{a}^2}{a^2}$  and  $\Lambda \sim \frac{\ddot{a}}{a}$  models can not exist as separate entity during inflation.

We have also established the relation between  $\alpha$ ,  $\beta$  and  $\gamma$ , the three parameters of the three forms of  $\Lambda$  which ultimately yields  $\Omega_m + \Omega_\Lambda = 1$ , the relation between cosmic matter and the vacuum energy density parameters for flat universe. Since  $\Omega_\Lambda = \frac{\rho_\Lambda}{\rho_c}$  and  $\Omega_m = \frac{\rho_m}{\rho_c}$  then it is clear from (35), (36) and (37) that  $\beta$  and  $\gamma$  are independent of  $\rho_c$ , the critical density of the universe whereas  $\alpha$  depends upon the critical density. It can also be shown that while  $\beta$  and  $\gamma$  decreases with the age of universe,  $\alpha$  increases with the passage of time.

Using (12) one can obtained the expression for the deceleration parameter  $q$  as

$$q = -\frac{a\ddot{a}}{\dot{a}^2} = \frac{(6 - \alpha)(1 + \omega)}{3} - 1.$$

Thus for an accelerating universe

$$\alpha > \frac{3(1 + 2\omega)}{(1 + \omega)}.$$

From above equation, it is evident that for dust case, an accelerating universe demands  $\alpha > 3$ . Now  $\alpha$  being the cosmic vacuum density parameter by the virtue of relation (35), we find that our model is fit for an accelerating universe since modern accepted value of  $\Omega_\Lambda$  is about 0.6666 [71–74]. Thus (34) implies that the value of  $\Omega_m$  is 0.3334, which provides  $\Omega_m + \Omega_\Lambda = 1$  and  $q = -0.3332$ , for the dust case.

It has already been shown that the three phenomenological  $\Lambda$ -models presented here are equivalent, then causal connection of the universe as indicated by the above model equally implies that the other models of the universe are also causally connected in the framework of higher dimensional space-time. For the present model age of the universe is inversely proportional to the Hubble parameter.

We also attempted to discuss the well known astrophysical phenomena, namely the neo-classical test, the luminosity distance-redshift, the angular diameter distance-redshift and look back time-redshift for the model in higher dimensional space time. It is has been observed that such models are compatible with the results of recent observations and the cosmological term  $\Lambda$  gradually reduces as the universe expands.

**Acknowledgements** The authors wish to thanks the IUCAA, Pune, India, for providing warm hospitality and excellent facilities where this work was done.

## References

1. Chodos, A., Detweiler, S.: Where has the fifth dimension gone? *Phys. Rev. D* **21**, 2167 (1980)
2. Freund, P.G.O.: Kaluza-Klein cosmologies. *Nucl. Phys. B* **209**, 146 (1982)
3. Sahdev, D.: Towards a realistic Kaluza-Klein cosmology. *Phys. Lett. B* **137**, 155 (1984)
4. Shafi, Q., Wetterich, C.: Cosmology from higher dimensional gravity. *Phys. Lett. B* **129**, 387 (1983)
5. Chatterjee, S.: Some remarks on luminosity distance and nucleosynthesis in higher dimensional cosmologies. *Astrophys. J.* **397**, 1 (1992)
6. Frieman, J.A., Waga, I.: Constraints from high redshift supernovae upon scalar field cosmologies. *Phys. Rev. D* **57**, 4642 (1998)
7. Carlberg, R., et al.: Galaxy cluster virial masses and Omega. *Astrophys. J.* **462**, 32 (1996)
8. Özer, M., Taha, M.O.: *Nucl. Phys. B* **287**, 776 (1987)
9. Freese, K., Adams, F.C., Frieman, J.A., Motta, E.: *Nucl. Phys. B* **287**, 797 (1987)
10. Carvalho, J.C., Lima, J.A.S., Waga, I.: Cosmological consequences of a time dependent Lambda term. *Phys. Rev. D* **46**, 2404 (1992)
11. Silveira, V., Waga, I.: *Phys. Rev. D* **50**, 4890 (1994)
12. Ratra, B., Peebles, P.J.E.: Cosmological consequences of a rolling homogeneous scalar field. *Phys. Rev. D* **37**, 3406 (1988)
13. Dolgov, A.D.: In: Gibbons, G.W., Hawking, S.W., Siklos, S.T.C. (eds.) *The Very Early Universe*. Cambridge University Press, Cambridge (1983)
14. Dolgov, A.D.: Higher spin fields and the problem of cosmological constant. *Phys. Rev. D* **55**, 5881 (1997)
15. Dolgov, A.D., Sazhin, M.V., Zeldovich, Y.B.: *Basics of Modern Cosmology*. Editions Frontiers (1990)
16. Sahni, V., Starobinsky, A.: The case for a positive cosmological Lambda-term (2000). *Int. J. Mod. Phys. D* **9**, 373 (2000), astro-ph/9904398
17. Padmanabhan, T.: Cosmological constant—the weight of the vacuum. *Phys. Rep.* **380**, 235 (2003), hep-th/0212290
18. Peebles, P.J.E.: The cosmological constant and dark energy. *Rev. Mod. Phys.* **75**, 599 (2003), astro-ph/0207347
19. Zeldovich, Y.B.: *Phys. Usp.* **11**, 381 (1968)
20. Weinberg, S.: The cosmological constant problem. *Rev. Mod. Phys.* **61**, 1 (1989)
21. Carrol, S.M., Press, W.H., Turner, E.L.: *Ann. Rev. Astron. Astrophys.* **30**, 499 (1992)
22. Perlmutter, S., et al.: Measurements of the cosmological parameters omega and lambda from the first supernova at  $z \geq 0.35$ . *Astrophys. J.* **483**, 565 (1997), astro-ph/9608192
23. Perlmutter, S., et al.: Discovery of a supernova explosion at half the age of the universe and its cosmological implication. *Nature* **391**, 51 (1998), astro-ph/9712212
24. Perlmutter, S., et al.: The supernova cosmological project. *Astrophys. J.* **517**, 565 (1999), astro-ph/9608192
25. Riess, A.G., et al.: Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astron. J.* **116**, 1009 (1998), astro-ph/9805201
26. Riess, A.G.: *PASP* **114**, 1284 (2000)
27. Garnavich, P.M., et al.: The constraints on cosmological models from Hubble spacetelescope observations of high z-supernova. *Astrophys. J.* **493**, L53 (1998), astro-ph/9710123
28. Garnavich, P.M., et al.: Supernova limits on the cosmic equation of state. *Astrophys. J.* **509**, 74 (1998), astro-ph/9806396
29. Schmidt, B.P., et al.: The high-Z supernova search: Measuring cosmic deceleration and global curvature of the universe using type Ia supernova. *Astrophys. J.* **507**, 46 (1998), astro-ph/9805200
30. Bertolami, O.: *Nuovo Cimento* **93**, 36 (1986)
31. Bertolami, O.: *Fortschr. Phys.* **34**, 829 (1986)
32. Berman, M.S.: *Int. J. Theor. Phys.* **29**, 567 (1990)
33. Berman, M.S.: *Int. J. Theor. Phys.* **29**, 1419 (1990)
34. Berman, M.S.: *Phys. Rev. D* **43**, 75 (1991)
35. Berman, M.S., Som, M.M.: *Int. J. Theor. Phys.* **29**, 1411 (1990)
36. Berman, M.S., Som, M.M., Gomide, F.M.: *Gen. Relativ. Gravit.* **21**, 287 (1989)
37. Berman, M.S., Gomide, F.M.: *Gen. Relativ. Gravit.* **22**, 625 (1990)
38. Beesham, A.: *Int. J. Theor. Phys.* **25**, 1295 (1986)
39. Beesham, A.: *Phys. Rev. D* **48**, 3539 (1993)
40. Singh, T., Beesham, A., Mbokazi, W.S.: *Gen. Relativ. Gravit.* **30**, 537 (1998)
41. Pradhan, A., Kumar, A.: *Int. J. Mod. Phys. D* **10**, 291 (2001)
42. Pradhan, A., Yadav, V.K.: *Int. J. Mod. Phys. D* **11**, 983 (2002)
43. Gasperini, M.: *Phys. Lett. B* **194**, 347 (1987)

44. Gasperini, M.: *Class. Quantum Gravity* **5**, 521 (1988)
45. Peebles, P.J.E., Ratra, B.: *Astrophys. J.* **325**, L17 (1988)
46. Chen, W., Wu, Y.S.: *Phys. Rev. D* **41**, 695 (1990)
47. Abdel-Rahaman, A.-M.M.: *Gen. Relativ. Gravit.* **22**, 655 (1990)
48. Abdel-Rahaman, A.-M.M.: *Phys. Rev. D* **45**, 3492 (1992)
49. Carvalho, J.C., Lima, J.A.C., Waga, I.: *Phys. Rev. D* **46**, 2404 (1992)
50. Waga, I.: *Astrophys. J.* **414**, 436 (1993)
51. Silveira, V., Waga, I.: *Phys. Rev. D* **50**, 4890 (1994)
52. Vishwakarma, R.G.: *Class. Quantum Gravity* **17**, 3833 (2000)
53. Al-Rawaf, A.S., Taha, M.O.: *Gen. Relativ. Gravit.* **28**, 935 (1996)
54. Al-Rawaf, A.S.: *Mod. Phys. Lett. A* **13**, 429 (1998)
55. Overdin, J.M., Cooperstock, F.I.: *Phys. Rev. D* **58**, 043506 (1998)
56. Arbab, A.I.: *J. Cosmol. Astropart. Phys.* **5**, 008 (2003), astro-ph/0212565
57. Arbab, A.I.: *Class. Quantum Gravity* **20**, 93 (2003), gr-qc/9905066
58. Vishwakarma, R.G.: *Class. Quantum Gravity* **18**, 1159 (2001)
59. Gurvits, L.I., Kellermann, K.I., Frey, S.: *Astron. Astrophys.* **342**, 378 (1999)
60. Ray, S., Mukhopadhyay, U.: astro-ph/0407295 (2004)
61. der Bernardis, P., et al.: *Nature* **404**, 955 (2000)
62. der Bernardis, P., et al.: *Astrophys. J.* **564**, 559 (2002)
63. Netterfield, C.B., et al.: *Astrophys. J.* **571**, 604 (2002)
64. Hanany, S., et al.: *Astrophys. J.* **545**, L5 (2000)
65. Lee, A.T., et al.: *Astrophys. J.* **561**, L1 (2001)
66. Balbi, A., et al.: *Astrophys. J.* **545**, L1 (2001)
67. Halverson, N.W., et al.: *Astrophys. J.* **568**, 38 (2002)
68. Sievers, J.L., et al.: *Astrophys. J.* **591**, 599 (2003)
69. Bennet, C.L., et al.: *Astrophys. J. Suppl.* **148**, 1 (2003)
70. Spergel, D.N., et al.: *Astrophys. J. Suppl.* **148**, 175 (2003)
71. Turner, M.S.: Preprint astro-ph/0207297 (2002)
72. Frampton, P.H., Takahashi, T.: *Phys. Lett. B* **557**, 135 (2003)
73. Perlmutter, S.: *Phys. Today*, 53 (2003)
74. Cardone, V.F., Troisi, A., Capozziello, S.: *Phys. Rev. D* **69**, 083517 (2004)