

Geometric Phase in a Time-Dependent System with Higgs Algebra Structure

Zhao-Xian Yu · Zhi-Yong Jiao · Xiang-Gui Li

Received: 4 May 2009 / Accepted: 2 July 2009 / Published online: 14 July 2009
© Springer Science+Business Media, LLC 2009

Abstract By using of the invariant theory, we have studied the geometric phase in a time-dependent system with Higgs algebra structure, the dynamical and geometric phases are given, respectively. The Aharonov-Anandan phase is also obtained under the cyclical evolution. The disappearing condition of the geometric phase is given.

Keywords Geometric phase · Higgs algebra · Invariant theory

1 Introduction

It is known that the concept of geometric phase was first introduced by Pancharatnam [1] in studying the interference of classical light in distinct states of polarization. Berry [2] found the quantal counterpart of Pancharatnam's phase in the case of cyclic adiabatic evolution. The extension to non-adiabatic cyclic evolution was developed by Aharonov and Anandan [3]. Samuel et al. [4] generalized the pure state geometric phase by extending it to non-cyclic evolution and sequential projection measurements. The geometric phase is a consequence of quantum kinematics and is thus independent of the detailed nature of the dynamical origin of the path in state space. Mukunda and Simon [5] gave a kinematic approach by taking the path traversed in state space as the primary concept for the geometric phase. Further generalizations and refinements, by relaxing the conditions of adiabaticity, unitarity, and cyclicity of the evolution, have since been carried out [6]. Recently, the geometric phase of the mixed states has also been studied [7–9].

Z.-X. Yu (✉)

Department of Physics, Beijing Information Science and Technology University, Beijing 100192, China
e-mail: zyu1965@163.com

Z.-Y. Jiao

Department of Physics, China University of Petroleum (East China), Dongying 257061, China

X.-G. Li

Department of Mathematics, Beijing Information Science and Technology University, Beijing 100192, China

As we know that the quantum invariant theory proposed by Lewis and Riesenfeld [10] is a powerful tool for treating systems with time-dependent Hamiltonians. It was generalized by introducing the concept of basic invariants and used to study the geometric phases [11–14] in connection with the exact solutions of the corresponding time-dependent Schrödinger equations. The discovery of Berry’s phase is not only a breakthrough in the older theory of quantum adiabatic approximations, but also provides us with new insights in many physical phenomena. The concept of Berry’s phase has been developed in some different directions [15–27]. In this paper, by using of the invariant theory, we shall study the geometric phase in a time-dependent system with Higgs algebra structure.

2 Model

The Hamiltonian of the generalized time-dependent system with Higgs algebra structure can be written by

$$\hat{H} = \omega(t)\hat{L}_z + g(t)\hat{R}_+ + g^*(t)\hat{R}_-, \tag{1}$$

here \hat{L}_z and \hat{R}_\pm are the generators of the Higgs algebra, and they satisfy the commutation relations

$$[\hat{L}_z, \hat{R}_\pm] = \pm\hat{R}_\pm, \quad [\hat{R}_+, \hat{R}_-] = f\hat{L}_z + h\hat{L}_z^3 \tag{2}$$

where $\hat{L}_z = \hat{x}\hat{p}_y - \hat{y}\hat{p}_x$.

3 Dynamical and Geometric Phases

For self-consistent, we first illustrate the Lewis-Riesenfeld (L-R) invariant theory [10]. For a one-dimensional system whose Hamiltonian $\hat{H}(t)$ is time-dependent, then there exists an operator $\hat{I}(t)$ called invariant if it satisfies the equation

$$i\frac{\partial\hat{I}(t)}{\partial t} + [\hat{I}(t), \hat{H}(t)] = 0. \tag{3}$$

The eigenvalue equation of the time-dependent invariant $|\lambda_n, t\rangle$ is given

$$\hat{I}(t)|\lambda_n, t\rangle = \lambda_n|\lambda_n, t\rangle, \tag{4}$$

where $\frac{\partial\lambda_n}{\partial t} = 0$. The time-dependent Schrödinger equation for this system is

$$i\frac{\partial|\psi(t)\rangle_s}{\partial t} = \hat{H}(t)|\psi(t)\rangle_s. \tag{5}$$

According to the L-R invariant theory, the particular solution $|\lambda_n, t\rangle_s$ of (5) is different from the eigenfunction $|\lambda_n, t\rangle$ of $\hat{I}(t)$ only by a phase factor $\exp[i\delta_n(t)]$ for the non-degenerate state, i.e.,

$$|\lambda_n, t\rangle_s = \exp[i\delta_n(t)]|\lambda_n, t\rangle, \tag{6}$$

which shows that $|\lambda_n, t\rangle_s$ ($n = 1, 2, \dots$) forms a complete set of the solutions of (5). Then the general solution of the Schrödinger equation (5) can be written by

$$|\psi(t)\rangle_s = \sum_n C_n \exp[i\delta_n(t)]|\lambda_n, t\rangle, \tag{7}$$

where

$$\delta_n(t) = \int_0^t dt' \left\langle \lambda_n, t' \left| i \frac{\partial}{\partial t'} - \hat{H}(t') \right| \lambda_n, t' \right\rangle, \tag{8}$$

and $C_n = \langle \lambda_n, 0 | \psi(0) \rangle_s$.

We can introduce the L-R invariant as follows

$$\hat{I} = \alpha(t) \hat{L}_z + \beta(t) \hat{R}_+ + \beta^*(t) \hat{R}_-, \tag{9}$$

one has the auxiliary relations from (3) and (9)

$$i\dot{\alpha} + (f - k)(\beta g^* - \beta^* g) = 0, \quad i\dot{\beta} + \alpha g - \beta \omega = 0, \tag{10}$$

where dot denotes the time derivative. Here we have used the large quantum numbers approximation, i.e., we let $[\hat{R}_+, \hat{R}_-] = f \hat{L}_z + h \hat{L}_z^3 = f \hat{L}_z + k$, k is the constant.

We now construct the unitary transformation

$$\hat{V}(t) = \exp[\xi \hat{R}_+ - \xi^* \hat{R}_-]. \tag{11}$$

It is easy to find that when satisfying the following relations

$$\frac{\alpha k}{f} [\cos(\sqrt{2f}|\xi|) - 1] - \frac{\beta k(\xi + \xi^*)}{\sqrt{2f}|\xi|} \sin(\sqrt{2f}|\xi|) = 0, \tag{12}$$

$$\frac{\alpha \xi}{\sqrt{2f}|\xi|} \sin(\sqrt{2f}|\xi|) + \frac{\beta}{2} [\cos(\sqrt{2f}|\xi|) + 1] + \frac{\beta^* \xi^2}{2|\xi|^2} [\cos(\sqrt{2f}|\xi|) - 1] = 0, \tag{13}$$

$$\alpha = \cos(\sqrt{2f}|\xi|), \quad \theta = \sqrt{2f}|\xi|, \quad \xi = \frac{-\theta}{\sqrt{2f}} e^{-i\gamma}, \quad \beta = \frac{\sin \theta}{\sqrt{2f}} e^{-i\gamma}, \tag{14}$$

then a time-independent invariant \hat{I}_V appears

$$\hat{I}_V = \hat{V}^\dagger(t) \hat{I} \hat{V}(t) = \hat{L}_z. \tag{15}$$

In terms of the unitary transformation $\hat{V}(t)$ and the Baker-Campbell-Hausdorff formula [28]

$$\hat{V}^\dagger(t) \frac{\partial \hat{V}(t)}{\partial t} = \frac{\partial \hat{\phi}}{\partial t} + \frac{1}{2!} \left[\frac{\partial \hat{\phi}}{\partial t}, \hat{\phi} \right] + \frac{1}{3!} \left[\left[\frac{\partial \hat{\phi}}{\partial t}, \hat{\phi} \right], \hat{\phi} \right] + \frac{1}{4!} \left[\left[\left[\frac{\partial \hat{\phi}}{\partial t}, \hat{\phi} \right], \hat{\phi} \right], \hat{\phi} \right] + \dots, \tag{16}$$

where $\hat{V}(t) = \exp[\hat{\phi}(t)]$. When the following relation holds

$$\omega + \sqrt{\frac{f}{2}} [g(1 + \cos \theta) e^{i\gamma} - g^*(1 - \cos \theta) e^{-i\gamma}] - i \dot{\xi} \sqrt{2f} e^{i\gamma} + \dot{\gamma} (1 - \sin \theta) = 0, \tag{17}$$

one has

$$\begin{aligned} \hat{H}_V(t) &= \hat{V}^\dagger(t) \hat{H}(t) \hat{V}(t) - i \hat{V}^\dagger(t) \frac{\partial \hat{V}(t)}{\partial t} \\ &= \frac{k}{f} \sin^2 \theta - \frac{k}{f} \dot{\gamma} (1 - \cos \theta) \\ &\quad + \left[\omega \cos \theta + \sqrt{\frac{f}{2}} (g e^{i\gamma} + g^* e^{-i\gamma}) \sin \theta + \dot{\gamma} (1 - \cos \theta) \right] \hat{L}_z, \end{aligned} \tag{18}$$

it is apparent that $\hat{H}(t)$ differs from \hat{I}_V only by a time-dependent c-number factor.

The phase $\delta(t) = \delta^d(t) + \delta^g(t)$ includes the dynamical phase

$$\delta^d(t) = - \int_{t_0}^t \left\{ \frac{k}{f} \sin^2 \theta + \left[\omega \cos \theta + \sqrt{\frac{f}{2}} (g e^{i\gamma} + g^* e^{-i\gamma}) \sin \theta \right] \langle \hat{L}_z \rangle \right\} dt', \quad (19)$$

and the geometric phase

$$\delta^g(t) = - \int_{t_0}^t \left\{ -\frac{k}{f} \dot{\gamma} (1 - \cos \theta) + \dot{\gamma} (1 - \cos \theta) \langle \hat{L}_z \rangle \right\} dt'. \quad (20)$$

Particularly, the geometric phase becomes under the cyclical evolution

$$\delta^g(t) = \oint \left[\frac{k}{f} - \langle \hat{L}_z \rangle \right] (1 - \cos \theta) d\gamma, \quad (21)$$

which is the known geometric Aharonov-Anandan phase. It is pointed out that when satisfying $\frac{k}{f} = \langle \hat{L}_z \rangle$, the geometric phase disappears.

Acknowledgements This work was supported by the NSF of China under Grant No.10671023, and the Beijing NSF under Grant No. 1072010, and supported by Scientific Creative Platform Foundation of Beijing Municipal Commission of Education under Grant No. PXM2008014224067420.

References

- Pancharatnam, S.: Proc. Indian Acad. Sci., Sect. A **44**, 247 (1956)
- Berry, M.V.: Proc. R. Soc. Lond., Ser. A **392**, 45 (1984)
- Aharonov, Y., Anandan, J.: Phys. Rev. Lett. **58**, 1593 (1987)
- Samuel, J., Bhandari, R.: Phys. Rev. Lett. **60**, 2339 (1988)
- Mukunda, N., Simon, R.: Ann. Phys. (N.Y.) **228**, 205 (1993)
- Pati, A.K.: Phys. Rev. A **52**, 2576 (1995)
- Uhlmann, A.: Rep. Math. Phys. **24**, 229 (1986)
- Sjöqvist, E.: Phys. Rev. Lett. **85**, 2845 (2000)
- Tong, D.M., et al.: Phys. Rev. Lett. **93**, 080405 (2004)
- Lewis, H.R., Riesenfeld, W.B.: J. Math. Phys. **10**, 1458 (1969)
- Gao, X.C., Xu, J.B., Qian, T.Z.: Phys. Rev. A **44**, 7016 (1991)
- Gao, X.C., Fu, J., Shen, J.Q.: Eur. Phys. J. C **13**, 527 (2000)
- Gao, X.C., Gao, J., Qian, T.Z., Xu, J.B.: Phys. Rev. D **53**, 4374 (1996)
- Shen, J.Q., Zhu, H.Y.: [arXiv:quant-ph/0305057v2](https://arxiv.org/abs/quant-ph/0305057v2)
- Richardson, D.J., et al.: Phys. Rev. Lett. **61**, 2030 (1988)
- Wilczek, F., Zee, A.: Phys. Rev. Lett. **25**, 2111 (1984)
- Moody, J., et al.: Phys. Rev. Lett. **56**, 893 (1986)
- Sun, C.P.: Phys. Rev. D **41**, 1349 (1990)
- Sun, C.P.: Phys. Rev. A **48**, 393 (1993)
- Sun, C.P.: Phys. Rev. D **38**, 298 (1988)
- Sun, C.P., et al.: J. Phys. A **21**, 1595 (1988)
- Sun, C.P., et al.: Phys. Rev. A **63**, 012111 (2001)
- Chen, G., Li, J.Q., Liang, J.Q.: Phys. Rev. A **74**, 054101 (2006)
- Chen, Z.D., Liang, J.Q., Shen, S.Q., Xie, W.F.: Phys. Rev. A **69**, 023611 (2004)
- He, P.B., Sun, Q., Li, P., Shen, S.Q., Liu, W.M.: Phys. Rev. A **76**, 043618 (2007)
- Li, Z.D., Li, Q.Y., Li, L., Liu, W.M.: Phys. Rev. E **76**, 026605 (2007)
- Niu, Q., Wang, X.D., Kleinman, L., Liu, W.M., Nicholson, D.M.C., Stocks, G.M.: Phys. Rev. Lett. **83**, 207 (1999)
- Wei, J., Norman, E.: J. Math. Phys. **4**, 575 (1963)