

# ERRATA TO: FIRST ORDER THEORY OF PERMUTATION GROUPS<sup>†</sup>

BY  
SAHARON SHELAH

Lemmas 4.1, 4.4 (of [1]) are incorrect, hence 0.1 and 4.2, 4.4, 4.5 fall. We give here a correct version.

DEFINITION 1. (A) For any  $\alpha$  there is a unique representation

$$\alpha = \Omega^\omega \alpha_\omega + \dots + \Omega^n \alpha_{[n]} + \dots + \Omega \alpha_{[1]} + \alpha_{[0]}; \alpha_{[n]} < \Omega.$$

Let

$$\alpha[n] = \Omega^\omega \alpha_\omega + \dots + \Omega^{n+1} \alpha_{[n+1]}; \alpha^{[n]} = \begin{cases} 1 + \text{cf} \alpha[n]; \text{cf} \alpha[n] < \Omega \\ 0 & \text{otherwise.} \end{cases}$$

(B) Define  $K^{15}$  by

$$M_\alpha^{15} = \langle U_\alpha, \alpha_{[0]}, \dots, \alpha_{[n]}, \dots, \alpha^{[0]}, \dots, \alpha^{[n]}, \dots, R_n(A_\alpha), \dots; \langle \rangle \rangle$$

where  $A_\alpha = U_\alpha \cup \bigcup_{n < \omega} \alpha_{[n]} \cup \bigcup_{n < \omega} \alpha^{[n]}$  in abuse of notation; this is a disjoint union.

Note that  $|A_\alpha| \leq 2^{\aleph_0}$ .

(C) Define  $K^{16}$  by

$$M_\alpha^{16} = \langle \alpha + 1, U_\alpha, \dots, R_n^\Omega((\alpha + 1) \cup U_\alpha), \dots; \langle \rangle \rangle.$$

REMARK. In  $M_\alpha^{15}$ , instead of one order sign  $\langle$ , we should have many: one for each  $\alpha_{[n]}, \alpha^{[n]}$ . Also we should have separate each  $R_n(A_\alpha)$  according to which place (in the relation) is designated for which domain.

Our main result is:

THEOREM 1.  $K^1, K^{15}, K^{16}$  are bi-interpretable.

---

<sup>†</sup> Volume 14, Number 2, pages 142-162  
Received May 23, 1973

Hence instead of conclusion 0.1 we have:

**COROLLARY 2.**  $\langle P_\alpha; \circ \rangle \equiv \langle P_\beta; \circ \rangle$  iff

$$\langle U_\alpha, \alpha_{[0]}, \alpha_{[1]}, \dots, \alpha^{[0]}, \alpha^{[1]}, \dots; < \rangle \equiv_{L_2} \langle U_\beta, \beta_{[0]}, \beta_{[1]}, \dots, \beta^{[0]}, \beta^{[1]}, \dots; < \rangle.$$

**REMARK.** A natural question is: in what is  $K^{15}$  better than  $K^7$  or even  $K^1$ ? A possible answer is

(1) Comparing the cardinals of the union of the domains, we get for  $K^1$ ,  $2^{\aleph_\alpha}$ , for  $K^7 \leq |\alpha| + 2^{\aleph_0}$ , and for  $K^{15} \leq 2^{\aleph_0}$ .

(2) In Corollary 2 the second part of the equivalence speaks on a well-known logic —  $L_2$ .

(3) In  $M_\alpha^{15}$  much irrelevant information on  $\alpha$  is thrown.

So for many  $\alpha$ 's we get isomorphic  $M_\alpha^{15}$ 's hence it is clear that they have the same first-order theory of  $\langle P_\alpha; \circ \rangle$ .

**LEMMA 3.**  $K^1, K^{16}$  are bi-interpretable.

**PROOF.** Trivially  $K^{16}, K^7$  are explicitly bi-interpretable. So by [1] 3.3  $K^1, K^{16}$  are bi-interpretable.

**LEMMA 4.**  $K^{15}$  is explicitly interpretable in  $K^{16}$ .

**PROOF.** Clearly there are formulas in  $L(K^{16})$  which define for  $\beta \in \alpha + 1$ ; the following (in  $M_\alpha^{16}$ ):

$\beta = \alpha, |\beta| \geq \Omega, \beta$  is divisible by  $\Omega, \beta$  is divisible by  $\Omega^2$  (i.e.  $\{\gamma: \gamma < \beta, \gamma$  divisible by  $\Omega\}$  has an order-type divisible by  $\Omega$ ),  $\beta$  is divisible by  $\Omega^n$ ;

$\gamma < \Omega$  is the cofinality of  $\beta$ ;  $\beta$  is the maximal  $\gamma \leq \alpha$  which is divisible by  $\Omega^n$  (for any fixed  $n$ ).

From this the lemma is clear.

**MAIN LEMMA 5.**  $K^{16}$  is interpretable in  $K^{15}$ .

**DEFINITION 2.** For any ordinal  $i$  and set of ordinals  $I$  let  $\gamma(i, I) =$  order type of  $\{j \leq i: (\forall \alpha \in I)(\alpha < i \rightarrow \alpha < j)\}$ .

**DEFINITION 3.** A  $k$ -representation of  $\langle \bar{a}, \bar{b}, \bar{r} \rangle = \langle a_1, \dots, b_1, \dots, r_1, \dots \rangle$ , where  $a_i \in (\alpha + 1), b_i \in U_\alpha, r_i \in R_n^\Omega((\alpha + 1) \cup U_\alpha)$  is a sequence,

$$\langle A, B, \bar{g}, < *, \bar{a}', \bar{b}', \bar{r}' \rangle = \langle A, B, g^0, \dots, g^k, g_0, \dots, g_k, < *, a'_1, \dots, b'_1, \dots, r'_1, \dots \rangle$$

such that, for some function  $F$ :

(1)  $A, B$  are disjoint subsets of  $U_\alpha$  [more exactly  $A, B \in R_1(A_\alpha)$ ].

(2)  $a'_i \in A, b'_i \in B, r'_i \in R_{n_i}, (A \cup B) \subseteq R_{n_i}((\alpha + 1) \cup U_\alpha), <^*$  a well ordering of  $U_\alpha$ , the  $g_n, g^n$ 's are one-place functions from  $A$  into  $U_\alpha$ .

(3)  $F$  is one-to-one, with domain  $A \cup B$  and range  $\subseteq (\alpha + 1) \cup U_\alpha$ ,

(4) the  $a_i, b_i$ 's and  $\alpha$  belong to the range of  $F$ , and  $U_\alpha$  and the domains of the  $r_i$ 's are included in it,

(5)  $F(a'_i) = a_i, F(b'_i) = b_i, F$  maps  $r'_i$  onto  $r_i, F$  maps  $A$  into  $(\alpha + 1), B$  onto  $U_\alpha$ ; and for  $a, b \in A, a <^* b$  if  $F(a) < F(b)$ ,

(6) for any  $a \in A$ ,

$$\text{order type of } \{c \in U_\alpha : c <^* g_i(a)\} = \gamma(F(a), F(A))_{[U]}$$

$$\text{order type of } \{c \in U_\alpha : c <^* g^l(a)\} = \gamma(F(a), F(a))^{[l]}.$$

REMARK. The definition depends on  $\alpha$ .

DEFINITION 4.  $\alpha \sim_k \beta$  if  $\alpha_{[l]} = \beta_{[l]}, \alpha^{[l]} = \beta^{[l]}$  for  $l \leq k$ .

LEMMA 6. A)  $\sim_k$  is an equivalence relation between ordinals; for each  $\alpha$  there is  $\beta < \Omega^{k+2}$  such that  $\alpha \sim_k \beta$ ; and if  $\alpha \sim_k \beta$ , then  $\alpha < \Omega^{k+1} \Leftrightarrow \beta < \Omega^{k+1} \Rightarrow \alpha = \beta$ .

B) If  $\alpha_i \sim_k \beta_i$  for  $i < \gamma$ , then  $\alpha = \sum_{i < \gamma} \alpha_i \sim_k \sum_{i < \gamma} \beta_i = \sum_{i < \gamma} \beta_i$ .

C) If  $\alpha \sim_{k+1} \beta, A \subset \alpha, |A| < \Omega$  then there is an order-preserving  $F: A \rightarrow \beta$  such that for every  $a \in A \cup \{\alpha\} \gamma(a, A) \sim_k \gamma(F(a), F(A))$  (where we stipulate  $F(\alpha) = \beta$ ).

REMARK. This lemma is not new, in fact, see e.g. Kino [2].

PROOF. A) Trivial.

B) We prove by induction on  $\gamma$ .

(I) For  $\gamma = 0, 1$  there is nothing to prove.

(II) For  $\gamma + 1$ , if  $\alpha_\gamma \geq \Omega^{k+1}$  then  $\beta_\gamma \geq \Omega^{k+1}$  (and vice versa) and then  $\alpha \sim_k \alpha_\gamma, \beta \sim_k \beta_\gamma$  hence  $\alpha \sim_k \beta$ . So assume  $\alpha_\gamma < \Omega^{k+1}$  so  $\alpha_\gamma = \beta_\gamma$  and it is easy to check that  $\alpha \sim_k \beta$ .

(III)  $\gamma$  a limit ordinal.

We can assume each  $\alpha_i, \beta_i$  is  $\neq 0$ , hence  $cf \alpha = cf \beta$ . If  $\{i < \gamma : \alpha_i \geq \Omega^{k+1}\}$  is unbounded, so is  $\{i < \gamma : \beta_i \geq \Omega^{k+1}\}$  hence  $\alpha$  and  $\beta$  are divisible by  $\Omega^{k+1}$ , together this implies  $\alpha \sim_k \beta$ . So we can assume that there is  $i_0 < \gamma$  such that  $i_0 \leq i < \gamma \Rightarrow \alpha_i < \Omega^{k+1}$  hence  $\alpha_i = \beta_i$ . So  $\sum_{\gamma > i \geq i_0} \beta_i = \sum_{\gamma > i \geq i_0} \alpha_i$  hence  $\alpha = \sum_{i \leq i_0} \alpha_i + (\sum_{\gamma > i > i_0} \alpha_i) \sim_k \sum_{i \leq i_0} \beta_i + (\sum_{\gamma > i > i_0} \beta_i) = \beta$ .

C) As  $\alpha \sim_{k+1} \beta, \alpha = \alpha^1 + \xi, \beta = \beta^1 + \xi, \alpha_1 \beta_1$  are divisible by  $\Omega^{k+2}$  and have equal cofinalities or cofinalities  $\geq \Omega$ . It suffices to prove for the case  $\xi = 0$ ,

because we can define for  $\alpha^1 \leq i < \alpha$ ,  $F(i) = \beta^1 + (i - \alpha^1)$ . If  $\lambda = \text{cf } \alpha = \text{cf } \beta < \Omega$ , then  $\alpha = \sum_{i < \lambda} \alpha_i$ ,  $\beta = \sum_{i < \lambda} \beta_i$ , each  $\alpha_i \beta_i$  is divisible by  $\Omega^{k+2}$  and has cofinality  $\geq \Omega$ . We can now, for each  $i < \lambda$ , define  $F$  on

$$A \cap \{\xi: \sum_{j < i} \alpha_j \leq \xi < \sum_{j \leq i} \alpha_j\} \text{ into } \{\xi: \sum_{j < i} \beta_j \leq \xi < \sum_{j \leq i} \beta_j\}.$$

So we reduce the problem to the case  $\text{cf } \alpha, \text{cf } \beta \geq \Omega$ . In this case define  $F$  inductively, so that for each  $a \in A$ ,  $\gamma(a, A) \sim_k \gamma(F(a), F(A))$  and  $\gamma(F(a), F(A)) < \Omega^{k+2}$ . As  $\Omega$  is regular  $|A| < \Omega$ , this implies, by induction, that  $F(a) < \Omega^{k+2} \leq \beta$ .

CLAIM 7. In  $L(K^{15})$  there are formulas  $\phi_k$  such that  $M_\alpha^{15} \models \phi_k[A, <^*, \bar{g}, b]$  iff  $<^*$  is a well-ordering of  $U_\alpha$ ,  $A \subseteq U_\alpha$ ,

$$\bar{g} = \langle g_0, \dots, g_k, g^0, \dots, g^k \rangle,$$

$g_l, g^l$  are one-place functions from  $A$  into  $U_\alpha$

and if  $A = \{a_i: i < i_0\}$ ,  $U_\alpha = \{b(j): j < j_0\}$ ,  $b = b(j_1)$ ,  $i < j \Rightarrow a_i <^* a_j$ ,  $b_i <^* b_j$  and  $\alpha_i$  is an ordinal and  $b((\alpha_i)_{[1]}) = g_l(a_i)$ ,  $b((\alpha_i)^{[l]}) = g^l(a_i)$  for each  $i < i_0$ ,  $l \leq k$  then

$$\sum_{i < i_0} \alpha_i = j_1.$$

REMARK. Remember 6B.

PROOF. Just formalize what was said in the proof of Lemma 6B. As we have second order quantifiers, in fact, on  $U_\alpha$  (in  $M_\alpha^{15}$ ) this is easy.

CLAIM 8. For every kind of sequence  $\langle a_1, \dots, b_1, \dots, r_1, \dots \rangle$  (i.e. the number of  $a_i$ 's,  $b_i$ 's and  $r_i$ 's; and the number of places of each  $r_i$ ) for every  $k$ ;

(A) there is a formula  $\psi_k \in L(K^{15})$  such that for any sequence  $\langle A, B, \bar{g}, <^*, \bar{a}', \bar{b}', \bar{r}' \rangle$  of the right kind (for being a suitable presentation)

$M_\alpha^{15} \models \psi_k[A, B, \bar{g}, <^*, \bar{a}', \bar{b}', \bar{r}']$  iff  $\langle A, B, \bar{g}, <^*, \bar{a}', \bar{b}', \bar{r} \rangle$  is a  $k$ -representation (in  $M_\alpha^{15}$ ).

(B) Similarly there is a formula  $\theta_k \in L(K^5)$  saying that two  $k$ -representations have a common source.

PROOF. (A) Just go through Definition 3 and see that it can be done with the help of Claim 7.

(B) Go through the representation with minimal  $A$ : that is let  $A' = \{a'_1, \dots\} \cup \{\text{last element of } A\} \cup \{\text{the domain of the } r_i\text{'s intersection with } A\}$ .

LEMMA 9. For each formula  $\phi(\bar{x}, \bar{y}, \bar{z})$  in  $L(K^{16})$  we can effectively find

$k(\psi)$  and  $\phi^* \in L(K^{15})$  such that for any  $\alpha$ , and any  $a_1, \dots, \in \alpha + 1$ ,  $b_1, \dots, \in U_\alpha$ ,  $r_i \in R_{n_i}^\Omega((\alpha + 1) \cup U_\alpha)$  and any  $k(\psi)$ -representation  $\langle A, B, \bar{g}, <, \bar{a}', \bar{b}', \bar{r} \rangle$ ,  $M_\alpha^{15} \models \phi^*[A, B, \bar{g}, <, \bar{a}', \bar{b}', \bar{r}']$  iff  $M_\alpha^{16} \models \phi[a_1, \dots, b_1, \dots, r_1, \dots]$ .

REMARK. The proof is similar to 3.2, except the added parameter  $k$ .

PROOF. We prove it by induction of  $\phi$ . For atomic formulas conjunction and negation there is no problem (remember that from  $k_1$ -representation we can get a  $k_2$ -representation, if  $k_1 > k_2$ , by omitting some  $g$ 's). So we are left with the case of existential quantifiers. Now note that two  $k$ -representations may have a common source, but nevertheless not all their sources are common. But by Lemma 6(C) if two  $(k + 1)$ -representations have a common source, then any source of one is  $k$ -represented by the other after omitting the suitable  $g$ 's. Now the proof should be clear.

PROOF OF LEMMA 5. By Lemma 9 it is immediate.

#### REFERENCES

1. S. Shelah, *First order theory of permutation groups*, Israel J. Math. 14 (1973), 149–162.
2. A. Kino, *On definability of ordinals with infinitely long expressions*, J. Symbolic Logic 31 (1966), 365–375 (correction in 32 (1967), 343).

INSTITUTE OF MATHEMATICS  
THE HEBREW UNIVERSITY OF JERUSALEM  
JERUSALEM, ISRAEL