

# DIFFERENTIALLY CLOSED FIELDS

BY

SAHARON SHELAH

## ABSTRACT

We prove that even the prime, differentially closed field of characteristic zero, is not minimal; that over every differential radical field of characteristic  $p$ , there is a closed prime one, and that the theory of closed differential radical fields is stable.

## Introduction

Let  $T_d$  be the theory of differential fields, that is, the axioms of fields in addition to the following axioms on the (abstract) differentiation operator:

$$D(x + y) = Dx + Dy$$

$$D(xy) = (Dx)y + xDy.$$

Let an upper index indicate the characteristic of the field.

$T_d$  is a natural generalization of the theory of fields which Ritt [4] invented. It is natural to look for an analog to the algebraic closure of a field. Seidenberg [7] has done algebraic work along these lines. Using his work, Robinson [5] showed that  $T_d^0$  has a model completion  $T_{dc}^0$  (that is, the theory of differentially closed fields of characteristic zero). Thus every  $T_d^0$ -field can be extended to a  $T_{dc}^0$ -field, however Robinson does not give an explicit set of axioms for  $T_{dc}^0$ . Blum [1] showed that the following axioms suffice:

(1)  $T_d^0$

(2) For differential polynomials  $P_1, P_2$  (in single variable  $y$ ) of order  $m_1, m_2$ , for  $m_1 > m_2$ , there is a solution of  $P_1 = 0$  which is not a solution of  $P_2 = 0$ , and there is a solution of  $P_2 = 0$ , provided that  $P_2$  has degree greater than zero.

Blum also showed that  $T_{dc}^0$  is totally transcendental, and the maximal Morley rank is  $\omega$  hence over every  $T_d^0$ -field there is a prime  $T_{dc}^0$ -field. (See Morely [3], or [6] for example.) By a general result of [10] (or [7], for example) this prime  $T_{dc}^0$ -field is unique. However here we answer a question of Blum (which appears in [6]) by showing that the prime  $T_{dc}^0$ -field is not necessarily minimal. This shows that the analogy with algebraically closed fields fails. The proof indicates to me (in contrast to Sacks [6, p. 307]) the following conjecture.

CONJECTURE 1.

(i) For every  $m < \omega$  there is a  $T_d^0$ -field  $F$ , and a differential polynomial  $P(y)$  of order  $m$  with coefficients in  $F$  such that the Morley rank of  $P(y) = 0$  is 1, (or at least less than  $m$ ) and that even  $P$  has integer coefficients.

(ii) Moreover, there is a differential polynomial of order 0 (less than  $m$ ),  $P_1(y) \neq 0$ , such that if  $y_1, \dots, y_n$  are solutions of  $P(y) = 0 \wedge P_1(y) \neq 0$  and  $P_2(y_1, \dots, D^i y_j, \dots)_{i < m} = 0$ , where  $P_2$  is a polynomial with coefficients in  $F$ , then  $P_2$  is the zero polynomial. (ii) implies (i).

Let us try to generalize to partial differentiation. Then we have a field with  $n$  differential operators,  $D_1, \dots, D_n$ , satisfying in addition, that  $D_i D_j y = D_j D_i y$ . But nothing new results. When the characteristic of the field is zero, we obtain a model completion with elimination of quantifiers, which is totally transcendental and has maximal Morley rank  $\omega n$ . I am quite sure that for characteristic  $p$  as well, this does not make any essential difference. If we add  $D_n$ ,  $n < \omega$ , we arrive at a stable but not superstable theory.

We also show that although  $T_{dc}^0$  is trivial in some aspects, when we allow cardinality quantifiers, it becomes complex. Hence  $T_{dc}^0$  has  $2^\lambda$  non-isomorphic models in every  $\lambda > \aleph_0$ .

CONJECTURE 2.  $T_{dc}^0$  has  $2^{\aleph_0}$  non-isomorphic models of power  $\aleph_0$ .

Wood [13], again using Seidenberg [7] deals with  $T_d^p$  for  $p > 0$ . (Notice that here if an element  $a$  has a  $p$ -th root, then it is constant, that is,  $Da = 0$ .) Wood showed that  $T_d^p$  does not have the amalgamation property. However, if we add the axiom

$$[Dx = 0 \rightarrow (r(x)^p = y)] \wedge [Dx \neq 0 \rightarrow r(x) = 0]$$

and obtain  $T_{rd}^p$  (that is, the theory of radical differential fields of characteristic  $p$ ), then it has the amalgamation property, and has a model completion  $T_{rdc}^p$ , which has elimination of quantifiers. (A  $T_d^p$ -field  $F$  can be expanded to a  $T_{rd}^p$ -field if

$Da = 0 \rightarrow (\exists x)(x^p = a)$  for  $a \in F$ , and the expansion is unique; thus we do not differentiate strictly between the field and its expansion.) Wood showed that, unlike  $F_{dc}^0$ ,  $T_{rdc}^p$  is not totally transcendental, hence the existence of a prime  $T_{rdc}^p$ -field remains an open question. Wood and the author independently solved the question (the author proved it after [9] but before [14] were submitted, see [15]). We do not know however, whether Conjecture 3 holds.

NOTE. Some of the results of this paper were previously announced in [9].

CONJECTURE 3. The prime  $T_{rdc}^p$ -field over any  $T_{rdc}^p$  is unique.

By small changes in [13] it follows that  $T_{rdc}^p$  is not superstable (see [12]).

### 1. The non-minimality of the prime differentially closed field

Now we state the main lemma of this section.

LEMMA 4. Let  $F$  be a differential field of characteristic zero,  $y_1, \dots, y_n$  distinct nonzero solution (in  $F$ ) of

$$Dy = \frac{y}{1+y}.$$

If  $P(x_1, \dots, x_n)$  is a polynomial with rational coefficients and, in  $F$ ,  $P(y_1, \dots, y_n) = 0$ , then  $P$  is identically zero.

PROOF.

Stage (i). Without loss of generality, assume  $F$  includes the field  $F_0$  of algebraic numbers. We suppose  $y_1, \dots, y_{n+1} \in F$  are distinct and not zero,  $P_0(y_1, \dots, y_{n+1}) = 0$ ,  $P_0(x_1, \dots, x_{n+1})$  is a nontrivial polynomial with algebraic coefficients and we shall arrive at a contradiction. Without loss of generality,  $n$  is minimal; for this  $n$  the degree of  $P_0$  in  $x_{n+1}$  is minimal and then the degree of  $P_0$  is minimal. Hence  $P_0(y_1, \dots, y_n, x)$  is indecomposable over  $F_0(y_1, \dots, y_n)$ , and  $y_1, \dots, y_n$  are transcendently independent over  $F_0$ .

Stage (ii). Let us look at the function  $x = y + \ln y$ . Clearly, for real  $y > 0$ ,  $y + \ln y$  is an increasing function whose range is the set of real numbers; let its inverse be  $y = f(x)$ . Thus  $f(x)$  is defined for every real  $x$ ; it increases with  $x$  and as  $x \rightarrow -\infty$ ,  $f(x) \rightarrow 0$ . As  $f(x) + \ln f(x) = x$ ,  $e^{f(x)} f(x) = e^x$ , hence for  $x \rightarrow -\infty$   $e^{o(1)} f(x) = e^x$  or  $f(x) = e^x(1 + o(1))$ . The function  $f(x)$  is also defined for complex arguments, and then it is holomorphic but it is not single valued.

Stage (iii). Look at the differential equation  $dy/dx = y/(1+y)$  (for  $y \neq 0$ ) or equivalently  $(1+y)/y dy = dx$  or  $dy/y + dy = dx$  or  $\ln y + y = x + c$  or  $y = f(x + c)$ . Thus if  $y = g(t)$  is a function with complex values defined for all

negative real numbers  $t < t_0$ , and if it is a solution of the equation, then for some complex  $c$  and branch of  $f$ ,  $g(t) = f(t + c)$  for every such  $t$ . Because, if  $t_1 < t_0$ ,  $g(t_1) \neq 0, -1$ , choose  $c_1 = g(t_1) + \ln g(t_1) - t_1$ ; then for a proper branch of  $f$ ,  $g(t_1) = f(t_1 + c_1)$ . Hence by the uniqueness theorem  $g(t) = f(t + c)$  for a neighborhood of  $t_1$ , and hence for all  $t < t_0$ . (We choose  $g(t_1) \neq 0$  to make  $f(g(t_1))$  well defined, and  $g(t_1) \neq -1$  to avoid the branching point when  $d/dy(y + \ln g(y)) = 0$ ). As  $g(t)$  for  $t < t_1$  cannot take always the values  $0, -1$ , we are through (remember that we assume  $y \neq 0$ ).

Looking at  $y + \ln y = x + c$ , we know that

(3) if  $x \rightarrow -\infty$  then either  $y \rightarrow -\infty$  or  $y \rightarrow 0$ .

If  $y \rightarrow -\infty$ ,  $y = x + O(\ln x)$ .

If  $y \rightarrow 0$ , then as before,  $y = e^{x+c}(1 + o(1))$ .

Stage (iv). Choose real negative numbers  $a_1, \dots, a_n$  such that  $f(a_1), \dots, f(a_n)$  will be algebraic numbers which are linearly independent over the rationals. By Lindemann's theorem (see [8])

$$e^{f(a_1)}, \dots, e^{f(a_n)}$$

are transcendently independent. Since  $f(a_i)$  are algebraic  $\neq 0$ , and  $e^{f(x)} f(x) = e^x$ ; also  $e^{a_1}, \dots, e^{a_n}$  are transcendently independent.

Stage (v). If  $P(x_1, \dots, x_n)$  is nontrivial, polynomial with algebraic coefficients, then for some  $t_1 < 0$   $P(f(t + a_1), \dots, f(t + a_n)) \neq 0$  for all  $t < t_1$ . Suppose not. Let us see what will be the dominant term when  $t \rightarrow -\infty$ . If  $P$  has a free constant as  $t \rightarrow -\infty$ ,  $f(t + a_i) \rightarrow 0$  (see Stage (ii)). This is a contradiction. Now let  $P(x_1, \dots, x_n) = \sum_{\eta \in I} c_\eta \prod_{i=1}^n x_i^{\eta(i)}$ , for  $c_\eta$  algebraic,  $\eta(i)$  natural numbers.

Then  $f(t + a_i) = \exp(t + a_i)(1 + o(1))$ ,

$$f(t + a_i)^n = \exp(nt + na_i)(1 + o(1))$$

$$\prod_{i=1}^n f(t + a_i)^{\eta(i)} = \exp([\sum \eta(i)]t) \exp(\sum \eta(i)a_i) (1 + o(1)).$$

As  $t \rightarrow -\infty$ , clearly the dominant terms will be those with minimal  $\sum \eta(i)$ , say  $m$ . Let  $J = \{\eta \in I : \sum_{i=1}^n \eta(i) = m\}$  so for some  $\eta' \in J$   $c_{\eta'} \neq 0$ .

$$\begin{aligned} P(\dots, f(t + a_i), \dots) &= \sum_{\eta \in J} c_\eta \exp(mt) \exp(\sum \eta(i)a_i) (1 + o(1)) + O(\exp((m + 1)t)) \\ &= \sum_{\eta \in J} c_\eta \prod_{i=1}^n \exp(\sum \eta(i)a_i) \exp(mt) (1 + o(1)) \\ &\quad + O(\exp(t(m + 1))). \end{aligned}$$

For this to be zero for arbitrarily small  $t < 0$ , necessarily

$$0 = \sum_{\eta \in J} c_\eta \exp(\sum \eta(i)a_i) = \sum_{\eta \in J} c_\eta \prod_{i=1}^n (\exp(a_i))^{\eta(i)} .$$

As  $\eta' \in J$ ,  $c_{\eta'} \neq 0$ , this contradicts the transcendental independence of the  $e^{a_i}$  (see Stage (iv)).

Stage (vi). By (v),  $P_0(f(t + a_1), \dots, f(t + a_n), y) = 0$  as an equation in  $y$ , has a solution  $y = g(t)$  for each  $t < t_0$ , for some  $t_0$  (make the leading coefficient nonzero). Also we can assume that for  $t < t_0$ , the resultant of this polynomial is not zero. (If it is identically zero as a polynomial in  $f(t + a_i)$ ,  $P_0(x_1, \dots, x_{n+1})$  will be decomposable over  $F_0(x_1, \dots, x_n)$ , contradicting the minimality of the degree of  $P_0$  in  $x_{n+1}$ .)

Thus we can choose one branch of the solution  $y = g(t)$  hence, clearly,  $g$  is an analytic function.

Stage (vii).  $g$  is a solution of  $Dy = y/(1 + y)$ .

Let  $P_0 = \sum_{\eta \in I} c_\eta \prod_{i=1}^{n+1} x_i^{\eta(i)}$  ( $c_\eta$  algebraic,  $\eta(i)$  natural numbers). Note that if  $h(t)$  solves  $Dy = y/(1 + y)$ , then  $(d/dt) h(t)^m = h(t)^m [m/(1 + h(t))]$ . Then

$$\begin{aligned} 0 &= \frac{d}{dt} P_0(f(t + a_1), \dots, f(t + a_n), g(t)) \\ &= \sum_{\eta \in I} \left[ c_\eta \prod_{i=1}^n f(t + a_i)^{\eta(i)} g(t)^{\eta(n+1)} \left( \sum_{i=1}^n \frac{\eta(i)}{1 + f(t + a_i)} \right) \right] \\ &\quad + \sum_{\eta \in I} \left[ c_\eta \prod_{i=1}^n f(t + a_i)^{\eta(i)} g(t)^{\eta(n+1)-1} \eta(n+1) \right] \frac{dg(t)}{dt} . \end{aligned}$$

The coefficient of  $dg(t)/dt$  is  $(d/dy) P_0(x_1, \dots, x_n, y)$ . As for all  $t < t_0$  the resultant of  $P_0(f(t + a_1), \dots, f(t + a_n), y)$  is not zero, it has no common root with its derivative. So from the above-mentioned equality we can solve  $dg/dt$  (since the  $a_i$ 's are real  $f(t + a_i) > 0$ , hence  $1 + f(t + a_i) \neq 0$ ). Thus,  $dg/dt = P_1(\dots, f(t + a_i), \dots, g(t))/P_2(\dots, f(t + a_i), \dots, g(t))$ . In the same way, in the differential field  $F$ ,

$$Dy_{n+1} = P_1(\dots, y_i, \dots, y_{n+1})/P_2(\dots, y_i, \dots, y_{n+1}).$$

On the other hand  $Dy_{n+1} = y_{n+1}/(1 + y_{n+1})$ , so define

$$P_3(y_1, \dots, y_{n+1}) \equiv P_1(y_1, \dots, y_{n+1})(1 + y_{n+1}) - P_2(y_1, \dots, y_{n+1})y_{n+1} = 0.$$

As  $n$  was minimal,  $y_1, \dots, y_n$  were transcendentially independent. Hence the

polynomial  $P_3(y_1, \dots, y_n, x)$  is divisible by  $P_0(y_1, \dots, y_n, x)$ . (The quotient has coefficients in  $F_0(y_1, \dots, y_n)$  and we can assume no denominator becomes zero when we replace  $y_i$  by  $f(t + a_i)$   $t < t_0$ .) So  $P_3(f(t + a_1), \dots, f(t + a_n), g(t)) = 0$  or equivalently  $dg(t)/dt = g(t)/(1 + g(t))$ .

Stage (viii). For some  $b$  and proper branch of  $f$ ,  $g(t) = f(t + b)$  for every  $t < t_0$ ; and

(a)  $f(t + b) = t + O(\ln |t|)$  for  $t \rightarrow -\infty$

or

(b)  $f(t + b) = e^{t+b}(1 + o(1))$  for  $t \rightarrow -\infty$ .

We obtain this result by combining stages (iii) and (vii) and (3).

We shall now contradict possibility (a). What will be the dominant part of  $P_0(f(t + a_1), \dots, f(t + a_n), f(t + b))$  (which is identically zero)?

If  $P_0(x_1, \dots, x_{n+1})$  has a term  $c_1 x_{n+1}^m$ ,  $m \geq 0$ ,  $c_1 \neq 0$ , letting  $m$  be the maximal one, we obtain

$$P_0(f(t + a_1), \dots, f(t + a_n), f(t + b)) = c_1 t^m + O(t^{m-1} \ln |t|).$$

(Remember  $f(t) = e^t(1 + o(1))$  for  $t \rightarrow -\infty$ ). This goes to infinity when  $t \rightarrow -\infty$  a contradiction, so there is no such term. Let

$$P_0(x_1, \dots, x_{n+1}) = \sum_{\eta \in I} c_\eta \prod_{i=1}^{n+1} x_i^{\eta(i)}$$

where  $c_\eta$  are algebraic. Then this equals

$$(4) \quad \sum_{\eta \in I} c_\eta \prod_{i=1}^n \exp((t + a_i)\eta(i)) t^{\eta(n+1)}(1 + o(1))$$

so the dominant terms are those with  $\sum_{i=1}^n \eta(i)$  minimal, say  $m$ , and among them, those with maximal  $\eta(n + 1)$ , say  $k$ . So letting  $J = \{\eta \in I : \sum \eta(i) = m, \eta(i + 1) = k\}$ , (4) equals

$$\left( \sum_{\eta \in J} c_\eta \prod_{i=1}^n \exp(a_i \eta(i)) \right) e^{m t} \cdot t^k (1 + o(1)).$$

Hence necessarily  $\sum_{\eta \in J} c_\eta \prod_{i=1}^n (\exp(a_i))^{\eta(i)} = 0$ , contradicting Stage (iv).

So, necessarily, (b) holds.

Stage (ix). Let  $a_{n+1} = b$ ; by the last stage  $f(t + a_i) = \exp(t + a_i)(1 + o(1))$  for  $1 \leq i \leq n + 1$ .

As  $P_0(\dots, f(t + a_i), \dots) = 0$  and the dominant part of it for  $t \rightarrow -\infty$  is

$$\left( \sum_{\eta \in J} c_\eta \prod_{i=1}^n (\exp(a_i)^{\eta(i)}) \right) \exp(t \sum \eta(i))$$

( $J$  is the set of  $\eta \in I$  with minimal  $\sum \eta(i)$ ) so

$$P_4(x_1, \dots, x_{n+1}) = \sum_{\eta \in J} c_\eta \prod_{i=1}^{n+1} x_i^{\eta(i)}$$

is homogeneous) then necessarily  $P_4(\dots, e^{a_i}, \dots) = 0$ , that is,  $e^{a_1}, \dots, e^{a_{n+1}}$  are transcendently dependent. As  $P_4$  is homogeneous, for every  $t$ ,

$$P_4(\dots, e^{t+a_i}, \dots) = 0 \text{ or}$$

- (a)  $P_4(\dots, f(t + a_i) \exp(f(t + a_i)), \dots) = 0$ ; but also
- (b)  $P_0(\dots, f(t + a_i), \dots) = 0$ .

*Stage (x).* We choose  $a_1, \dots, a_n$ , only so that  $P^i(\dots, a_i, \dots, \dots, e^{a_i}, \dots) \neq 0$  for a specific finite set of polynomials  $P^i$  with algebraic coefficients. Thus there is an  $\varepsilon > 0$  and  $t'_0$  so that every  $a_i \in (a_i - \varepsilon, a_i + \varepsilon)$  will satisfy the same demands for  $t < t'_0$ , hence all our conclusions, in particular the existence of  $a'_{n+1}$ . Hence for  $t < t'_0$  (by (a), (b) from stage (ix))

- (a)  $P_4(\dots, f(t + a'_i) \exp(f(t + a'_i)), \dots) = 0$  and
- (b)  $P_0(\dots, f(t + a'_i), \dots) = 0$ .

Let  $k_1$  be the degree of  $P_0(x_1, \dots, x_{n+1})$ , and  $k_2$  be the dimension of the field  $F_1$  generated by the coefficients of  $P_0$  over the rationals.

Now choose  $t^* < t'_0$  so that  $t^* + a_i + \varepsilon < t'_0$ ; and choose  $a'_i$  in  $(a'_i - \varepsilon, a_i + \varepsilon)$  so that  $f(t^* + a'_i)$ ,  $i = 1, n$  are algebraic but not linearly dependent over the rationals and moreover  $f(t^* + a'_i) = q_1^i + q_2^i a^i$ ,  $q_1^i, q_2^i$  rationals,  $q_2^i \neq 0$  and  $a^i$  is the  $p^{(i)}$ -root of 2 where  $p^{(1)} > k_1 k_2$ ,  $p^{(i+1)} > \prod_{j \leq i} p^{(j)} k_1$ ,  $p^{(i)}$  natural numbers.

By (b)  $f(t^* + a'_{n+1})$  is algebraic over  $f(t^* + a'_i)$ ,  $i = 1, n$ ; hence algebraic, and  $\exp(f(t^* + a'_i))$ ,  $i = 1, n$  are transcendently independent by Lindman theorem, but  $\exp(f(t + a'_{n+1}))$  depends on them, by (a).

By (a) and Lindman's theorem (see [8]),  $f(t^* + a'_{n+1})$  is linearly dependent on  $a'_1, \dots, a'_n$  over the rationals, hence for rationals  $q_i$ ,  $f(t^* + a'_{n+1}) = \sum_{i=1}^n q_i f(t^* + a'_i)$ . We can substitute this in  $P_0(\dots, f(t^* + a'_i), \dots) = 0$  and obtain  $P_5(f(t^* + a'_1), \dots, f(t^* + a'_n)) = 0$ , where  $P_5$  is a polynomial over  $F_1$ , and the degree of  $P_5$  is  $\leq k_1$ . This implies that  $P_5$  is identically zero by dimensional consideration, and the condition on the set of  $p^{(i)}$ .

If we substitute in  $P_0(x_1, \dots, x_{n+1})$   $x_{n+1} = \sum q_i x_i$ , we obtain the zero polynomial. By the minimality of the degree of  $P_0$  in  $x_{n+1}$ , and in general, we can assume  $P_0(x_1, \dots, x_{n+1}) = x_{n+1} - \sum q_i x_i$ .

Stage (xi). Now

$$y_{n+1} = \sum_{i=1}^n q_i y_i \text{ for } q_i \text{ complex rationals.}$$

Hence

$$\begin{aligned} Dy_{n+1} &= \sum_{i=1}^n q_i Dy_i = \sum_{i=1}^n q_i \frac{y_i}{1 + y_i} = Dy_{n+1} = \frac{y_{n+1}}{1 + y_{n+1}} \\ &= \sum_{i=1}^n q_i \frac{y_i}{1 + y_i} = \sum_{i=1}^n q_i y_i \Big/ \left( 1 + \sum_{i=1}^n q_i y_i \right). \end{aligned}$$

As  $y_1, \dots, y_n$  are transcendently independent, this is an identity so it holds if we substitute for the set of  $y_i$  complex numbers. If  $i \neq j$ ,  $q_i \neq 0$ ,  $q_j \neq 0$  set  $y_i = -1 + \varepsilon$ ,  $y_j \neq -1$ ,  $-(1 + q_i y_i) 1/q_j$  and  $y_k = 0$  for  $k \neq i, j$ . Then we obtain a contradiction as  $\varepsilon \rightarrow 0$ . Thus  $n = 1$ ,  $y_2 = y_{n+1} = q_1 y_1$ , and

$$q_1 \frac{y_1}{1 + y_1} = \frac{q_1 y_1}{1 + q_1 y_1}.$$

For  $y_1 \neq 0$  we obtain  $q_1 = 0$  or  $q_1 = 1$ . If  $q_1 = 0$ ,  $y_2 = 0$ ; if  $q_1 = 1$ ,  $y_2 = y_1$ , a contradiction in any case.

**THEOREM 5.** *The prime differentially closed field is not minimal. (It is the prime  $T_{dc}^0$ -field over the field of rational numbers.)*

**PROOF.** Let  $F$  be that field. The equation  $Dy = y/(1 + y)$  is not an algebraic formula since in some  $T_{dc}^0$ -field (of functions) it has infinitely many solutions. Hence it has infinitely many nonzero solutions  $y_i \in F$ ,  $i < \omega$ . Since the theory  $T_{dc}^0$  has elimination of quantifiers, clearly the  $\{y_i; i < \omega\}$  is an indiscernible set, hence by [10] (or see for example [6]),  $F$  is not minimal. (The elaboration for this particular case is easy: there is a field  $F' \subseteq F$  prime over the field generated by  $\{y_{2i}; i < \omega\}$ , and  $F' \neq F$  as  $y_{2i+1} \notin F'$ ).

**LEMMA 6.** *Let  $F$  be a differential field;  $\{f, g\}$  differentially independent elements of  $F$ . Let  $y_1, \dots, y_n$  be distinct nonzero solutions of  $Dy = yf/(1 + y)$ ;  $y^1, \dots, y^m$  be distinct nonzero solutions of  $Dy = (y/(1 + y))g$ . Then for no non-trivial polynomial  $P$  with rational coefficients,  $P(y_1, \dots, y_n, y^1, \dots, y^m) = 0$ .*



PROOF. Similar to that of Theorem 5.

REMARK. No doubt the restrictions on  $f, g$  can be weakened.

THEOREM 7. For every  $\lambda > \aleph_0$ ,  $T_{dc}^0$  has  $2^\lambda$  non-isomorphic fields of power  $\lambda$ .

PROOF. Let  $F$  be a differentially closed field of power  $\lambda$ ,  $f_i, g_i \in F$ , and  $\{f_i: i < \lambda\} \cup \{g_i: i < \lambda\}$  a differentially independent set with  $F$  prime over it.

Let  $\phi(x_1, x_2) = [Dx_1 = (x_1/(1 + x_1))x_2]$ . By Lemma 6, if  $y$  is a new element satisfying  $\phi(y, f_i g_j)$ ,  $F'$  the prime differentially closed field over  $F(y)$ , and  $\langle h, l \rangle \neq \langle i, j \rangle$  then no  $y' \in F' - F$  satisfies  $\phi(y', f_h g_l)$ . By repeating, we can obtain for any binary relation  $R$  over  $\lambda$  a field  $F_R$  such that

$$|\{y \in F_R: \phi(y, f_i g_j)\}| = \aleph_1 \text{ iff } \langle i, j \rangle \in R \text{ iff}$$

$$|\{y \in F_R: \phi(y, f_i g_j)\}| \neq \aleph_0.$$

Then by [11] the result follows easily.

**2. On the existence of  $T_{rdc}^p$ -prime field over  $T_{rd}^p$ -field**

THEOREM 8. Over every differential radical field of characteristic  $p$  ( $= T_{rd}^p$ -field) there is a prime differentially closed radical field ( $= T_{rdc}^p$ -field).

PROOF.

Stage (i). By Morley [3] (or see [6]) it suffices to prove the following. (Remember that by Wood [13],  $T_{rdc}^p$  has elimination of quantifiers.)

Let  $F$  be a  $T_{rd}^p$ -field and let  $\phi(x)$  be a consistent formula with parameters from  $F$ . Then there is a consistent formula  $\psi(x)$  with parameters from  $F$  such that  $\psi(x) \rightarrow \phi(x)$  and  $\psi(x)$  defines an isolated type, that is, if  $y$  satisfies  $\psi$ , then the structure of  $F_{rd}(y)$  (the  $T_{rd}^p$ -field generated by  $F, y$ ) is uniquely defined. Without loss of generality,  $\phi$  is a quantifier-free formula and moreover it is a conjunction of atomic formulas and negation of action formulas.

We can also assume without loss of generality that  $F$  is separately closed.

Stage (ii). Let  $F' \cong F$  be a  $T_{rd}^p$ -field in which  $y$  satisfies  $\phi(x)$ . Let  $\tau_0 = \tau_0(y) = y$  and  $\tau_1 = \tau_1(y), \dots, \tau_n = \tau_n(y)$  be the terms appearing (maybe as subterms) in  $\phi(y)$  which are of the form  $r(\dots)$ . (Remember  $r$  is the  $p$ th root.) Let  $n(i)$  be the highest  $n$  such that  $D^n \tau_i$  appear in  $\phi$ . We can assume without loss of generality that in  $\phi$  there appears no term of the form  $D(\sigma_1 + \sigma_2)$  or  $D(\sigma_1 \sigma_2)$  (since then we could simplify it); and that if  $r(\sigma)$  appears in it, then one of the conjuncts of  $\phi$  is  $D\sigma = 0$

Thus if  $F \subseteq F'' \subseteq F'$ ,  $F''$  is a  $T^p$ -field and  $D^j\tau_i(y) \in F''$  for  $j \leq n(i)$ , then  $\phi(y)$  is meaningful in  $F''$ .

*Stage (iii).* We derive  $\phi'$  from  $\phi$  by adding to it for each  $i \leq n$  a conjunct as follows:

(a) If there is an  $m = m(i)$  such that  $D^m\tau_i(y)$  is in the separable closure of  $F'_i = F(\dots, D^j\tau_k(y), \dots, D^l\tau_i(y), \dots)_{k < i, l < m}$  then let  $P_i(x) = \sum_i \sigma_i^l x^l$  be an indecomposable polynomial over  $F'_i$  of which  $D^m\tau_i(y)$  is a root. Then the conjunct will be  $\sum_i \sigma_i^l [D^m\tau_i(y)]^l = 0 \wedge \sigma \neq 0$  where  $\sigma$  is the resultant of  $P_i(x)$ .

This guarantees that  $D^l\tau_i(y)$ ,  $l \geq m$  is in  $F_i(D^m\tau_i(y))$  and that  $D^m\tau_i(y)$  is separably algebraic over  $F'_i$ .

(b) If there is not such an  $m$ , we add nothing.

*Stage (iv).* Let  $F'' \subseteq F'$  be the  $T^p$ -field generated by  $F$  and  $D^j\tau_i(y)$  for  $j \leq n(i)$  (that is, generated only by the field operations). Supplement it by defining  $D(D^{n(i)}\tau_i(y)) = 0$ , if  $i$  satisfies (b) above; we obtain a  $T_d^p$ -field  $F^*$  and by [7] there is a  $T_{r_d}^p$ -field  $F^{**} \cong F^*$ . Add to  $\phi'$ , for each such  $i$ , the conjunct  $D^{n(i)+1}\tau_i(x) = 0$  to obtain  $\phi''$ .

*Stage (v).* Now case (a) of Stage (iii) always occurs, hence we can express each  $D^j\tau_i(y)$  ( $j > n(i)$ ) by a polynomial in  $\{D_k\tau_l(y) : k \leq n(l), l \leq n(i)\}$  with coefficients in  $F$ . Add to  $\phi''$  conjuncts so that the transcendence rank of  $F(\dots, D^k\tau_e(y), \dots) = F_d(\tau_0(y), \dots, \tau_n(y))$  is minimal. For each  $j \leq n(i)$ , if  $D^j\tau_i(y)$  is algebraically dependent on  $\{D^l\tau_k(y) : k < i \text{ or } k = i, l < j\}$ , then we obtain  $\phi'''$  by adding conjuncts to  $\phi$  to make the degree of the polynomial it solves as small as possible.

Without loss of generality let  $y$  in  $F'$  satisfy  $\psi(y) \equiv \phi'''(y)$ .

Now  $\psi$  completely determines the structure of

$$F'' = {}^{d^f}F(\dots, D^j\tau_i(y), \dots)_{j \leq n(i), i \leq n} = F_d(\tau_0(y), \dots, \tau_n(y)).$$

If  $F''$  is a  $T_{r_d}^p$ -field, then we are through. This is equivalent to saying that  $c \in F'' - F$ ,  $Dc = 0$  implies  $c$  has a  $p$  root in  $F''$ .

*Stage (vi).* Suppose  $c \in F'' - F$ ,  $Dc = 0$  but  $c$  has no  $p$  root in  $F''$ . We arrive at a contradiction.

Let  $c = P_0(\dots, D^j\tau_i(y), \dots)$ , where  $P_0$  is a polynomial over  $F$ . Now if in Stage (v) we had also added  $P_0(\dots, D^j\tau_i(x), \dots) = b$  for any  $b \in F$  to  $\phi''(x)$ , the transcendence rank of  $F_d(\tau_0(y), \dots)$  would have become smaller. We have not done it because it is impossible. In other words, letting

$$\theta_0(x_1) = (\exists x)(x_1 = P_0(\dots, D^j\tau_i(x), \dots) \wedge \psi(x))$$

and  $F^c \cong F''$  be a  $T_{rdc}^p$ -field, then for no  $b \in F$ ,  $F^c \models \theta_0(b)$ . As  $T_{rdc}^p$  has elimination of quantifiers for some quantifier-free  $\theta_1(x_1)$ ,  $T_{rdc}^p \vdash (\forall x_1) [\theta_1(x_1) \equiv \theta_0(x_1)]$ . Without loss of generality,  $F^c$  is  $|F|^+$ -saturated.

*Stage (vii).* Let  $F^0 \subseteq F$  be the prime field (that is, the one generated by 1) and let  $a_n \in F$ , for  $n < \omega$ , be distinct elements which are in the separable closure of  $F^0$  in  $F$ . Clearly  $F \models \neg \theta_1(a_n) \wedge Da_n = 0$ . By the compactness theorem there is an element  $a \in F^c - F$ ,  $F^c \models \neg \theta_1(a) \wedge D(a) = 0$ . Let  $F^1$  be the separable closure of  $F_{rd}^0(a)$  in  $F^c$  and let  $F^2$  be the separable closure of  $F^0(c)$  in  $F''$ . Clearly for  $b \in F^2$   $Db = 0$ , and there is an embedding  $f: F^2 \rightarrow F^1$ ,  $f(c) = a$  which is the identity on  $F^*$  (see below). Let  $F^3$  be the closure of  $F^1$  to a  $T_{rd}$ -subfield of  $F^c$ . Notice that  $F^* = \{b \in F^c: b \text{ is separably algebraic over } F^0\}$  is a  $T_{rd}$ -field; hence  $F^*$  is algebraically closed. The diagram is shown in Fig. 1 (arrows denote inclusion).




Fig. 1

Notice that:

(a) ([7]) although the amalgamation property does not hold for  $T_d^p$ -fields in general, if

1.  $g_1: F^\delta \rightarrow F_\alpha, g_2: F^\delta \rightarrow F_\beta$  are embeddings of  $T_d^p$ -fields, and
2.  $b \in F^\delta, Db = 0$  but has no  $p$ -th root in  $F^\delta$  implies  $g_1(b)$  has no  $p$ -th root in  $F_\alpha$ ,
3. no  $b \in F_\alpha - g_1(F^\delta)$  is the root of a separable polynomial over  $g_1(F^\delta)$ , then there is a  $T_d^1$ -field  $F_\gamma$ , and embeddings  $f_1: F_\alpha \rightarrow F_\gamma, f_2 = F_\beta \rightarrow F_\gamma$  such that  $f_1 g_1 = f_2 g_2$ , and without loss of generality for example  $f_1$  is the identity.

(b) If  $b \in F^2$ , and  $b$  has no  $p$ -th root in  $F^2$  then  $b$  has no  $p$ -th root in  $F''$ . Because, without loss of generality,  $b \notin F^*$ . Suppose  $b$  has a  $p$ -th root in  $F''$ . Then  $\sum_{j < n} (\sum_{i < n(j)} t_{ij} c^i) b^j = 0, t_{ij} \in F^0$  where  $\sum_{j < n} (\sum_{i < n(j)} t_{ij} c^i) x^j$  is indecomposable, and for some  $j \neq 0 \pmod p$   $\sum_i t_{ij} c^i \neq 0$  (because  $b \in F^2$ ). We can assume  $n, n(j)$  are minimal. As  $c, b \notin F^*$ , they are transcendental over  $F_0$ , hence  $\sum_{ij} t_{ij} x^i y^j$  is indecomposable over  $F^0$ , and  $\sum_i (\sum_j t_{ij} b^j) x^i$  is indecomposable over  $F^0$ . Then in

$F^c$ ,  $\sum_i t_{ij} r(c)^i r(b)^j = 0$  (remember  $r(t_{ij}) = t_{ij}$  as  $t_{ij} \in F^0$ ). Since  $r(b) \in F''$ ,  $\sum_j t_{ij} r(b)^j \in F''$  but  $r(c)$  cannot be separably algebraic over  $F''$ . Now  $i \neq 0 \pmod p$  implies  $\sum_j t_{ij} r(b)^j = 0$ , hence  $\sum_j t_{ij} b^j = 0$  and  $t_{ij} = 0$  (as  $b$  is not algebraic over  $F^0$ ). Thus  $\sum_{ij} t_{pij} c^{pi} b^j = 0$ , and in  $F''$ ,  $\sum_{ij} t_{pij} c^i r(b)^j = 0$ , so  $r(b)$  is separably algebraic over  $F^2$  and  $r(b) \in F'' - F^2$ , and we have finished.

(c) No  $b \in F'' - F^2$  is the root of a separable polynomial over  $F^2$ , because  $F^2$  is the separable closure of  $F(c)$  in  $F''$ .

Stage (viii). Combine (a), (b), (c), and  $f: F^2 \rightarrow F^1$  from Stage (v).

Let  $F^\delta = F^2$ ,  $F_\alpha = F''$ ,  $F_\beta = F^3$ ,  $g_1 =$  the identity,  $g_2 = f$ . Then by (b), (c), (2) and (3) of (a) hold. Hence there are a  $T_{rdc}^p$ -field  $F_\gamma \supseteq F''$  and an embedding  $g: F^3 \rightarrow F_\gamma$  such that  $gf =$  identity, hence  $g(a) = c$ . Now  $F^3 \models \neg \theta_1(a)$  (we chose  $a$  in this way) hence  $F_\gamma \models \neg \theta_1(c)$ , hence  $F_\gamma \models \neg \theta_0(c)$ . But  $F_\gamma \supseteq F''$ , so  $F_\gamma \models \theta_0(c)$ , a contradiction.

Q.E.D.

### 3. Stability of $T_{rdc}^p$

THEOREM 9.  $T_{rdc}^p$  is stable.

PROOF.

Stage (i). Suppose  $F^1 \subseteq F^2$  are  $T_{rdc}^p$ -fields,  $|F^1| \leq \lambda$ . We should prove that the set of types elements of  $F^2$  realized over  $F^1$  is  $\leq \lambda^{k_0}$ . For each  $y \in F^2$  choose a countable field  $F_y \subseteq F^2$  such that

- (a)  $F_y$  is a countable  $T_{rdc}^p$ -field,  $y \in F_y$ ,
- (b)  $F_y \cap F^1$  is a  $T_{rdc}^p$ -field,
- (c) if  $a_1, \dots, a_n \in F_y$  are linearly dependent over  $F_1$ , then they are linearly dependent over  $F_y \cap F^1$ .

Let  $F^y$  be the field (=  $T_d^p$ -field) generated by  $F_y \cup F^1$ .

Stage (ii). Now define an equivalence relation over  $F^2$ :

$$y_1 \sim y_2 \text{ iff } F_{y_1} \cap F^1 = F_{y_2} \cap F^1,$$

and there is an isomorphism  $f$  from  $F_{y_1}$  onto  $F_{y_2}$ ,  $f(y_1) = y_2$ ,  $f$  restricted to  $(F_{y_1} \cap F^1) =$  identity.

Clearly  $\sim$  has  $\leq \lambda^{k_0}$  equivalence classes; if  $y_1 \sim y_2$  then we can extend the corresponding  $f$  to an isomorphism from  $F^{y_1}$  onto  $F^{y_2}$  which is the identity over  $F^1$ . If  $F^{y_1}$  is a  $T_{rdc}^p$ -field this implies (as  $T_{rdc}^p$  has elimination of quantifiers) that  $y_1, y_2$  realize the same type over  $F^1$ . Hence it suffices to prove

(5)  $F^y$  is  $T_d^p$ -field.

Let  $F = F_y \cap F^1, F_1 = F_y, F_2 = F^1$ .

REMARK. In fact we have more than the needed information to prove that the  $T_d^p$ -field  $F^y$ , generated by  $F_1, F_2$ , is a  $T_d^p$ -field.

Stage (iii). Suppose  $c^* \in F^y, Dc = 0$  but  $c$  has no  $p$ -th root in  $F^y$ . Thus  $c^* = \sum a_i b_i / \sum a^i b^i, a_i, a^i \in F_1, b_i, b^i \in F_2$ . Then  $c = \sum a_i'' b_i'' / (\sum^n a_i' b_i')^p$ , and clearly  $D(\sum a_i'' b_i'') = 0$ . So without loss of generality  $c = \sum_{i=1}^n a_i b_i, a_i \in F_1, b_i \in F_2$ . Choose the sets  $a_i, b_i$  so that  $n$  is minimal. This implies that

- (a)  $\{a_1, \dots, a_n\}$  are linearly independent over  $F$ ,
- (b)  $b_1, \dots, b_n$  are linearly independent over  $F$ . Hence
- (c)  $\{a_i b_j; i, j \leq n\}$  are linearly independent over  $F$ .

PROOF OF (c). If  $\sum_{i,j} t_{i,j} a_i b_j = 0, t_{i,j} \in F$  then  $\sum_i a_i (\sum_j t_{i,j} b_j) = 0$ . Since the  $a_i \in F_1$  are linearly independent over  $F$  they are also linearly independent over  $F_2$  (by Stages (a)-(c)); thus  $\sum_j t_{i,j} b_j = 0$  and hence  $t_{i,j} = 0$ .

Stage (iv).

- (a)  $Da_i$  is linearly dependent on  $\{a_1, \dots, a_n\}$  over  $F$ ;
- (b)  $Db_i$  is linearly dependent on  $\{b_1, \dots, b_n\}$  over  $F$ .

PROOF. Choose  $1 \leq i_1 < \dots < i_l \leq n$  such that  $\{a_1, \dots, a_n, Da_{i_1}, \dots, Da_{i_l}\}$  is linearly independent over  $F$ , and each  $Da_{i_l}$  depends on it over  $F$ . Choose similarly  $1 \leq j_1 < \dots < j_k \leq n$  such that  $\{b_1, \dots, b_n, Db_{j_1}, \dots, Db_{j_k}\}$  is linearly independent over  $F$  but each  $Db_{j_l}$  depends on it over  $F$ .

$$0 = Dc = \sum_i a_i Db_i + \sum_i (Da_i) b_i.$$

Substitute the expressions of  $Da_i, i \notin \{i_1, \dots, i_l\}$ , and for  $Db_j, j \notin \{j_1, \dots, j_k\}$ , and collect the terms. Then as in (iii) the coefficient of each  $a_i b_j, a_i Db_j, (Da_{i_m}) b_j$  is zero. If  $l > 0$  the coefficient of  $(Da_{i_1}) b_1$  is 1, a contradiction. Thus  $l = 0$ , and similarly  $k = 0$ . Hence for some  $t_j^i \in F, s_j^i \in F, Da_i = \sum_j t_j^i a_j, Db_i = \sum_j s_j^i b_j$ .

Stage (v). Let

$$(6) \quad Dx_i = \sum_{j=1}^n u_j^i x_j, \quad i < n; \quad u_j^i \in F$$

or, in short,  $D\bar{x} = U\bar{x}, \bar{x}$  is a vector of length  $n, U$  an  $n \times n$  matrix. Then there are solutions  $\bar{a}_0, \dots, \bar{a}_m$  (for  $m < n$ ) in  $F$  such that for any other solution  $\bar{a}$  from  $F^2$  there are  $d_0, \dots, d_m \in F^2, Dd_i = 0$  such that  $\bar{a} = \sum_{1 \leq m} d_i \bar{a}_i$ . Let  $\bar{a}_0, \dots, \bar{a}_m$  be a

maximal set of solutions of (6) which are linearly independent over  $F$  (as vectors). Let  $\bar{a}$  be any other solution from  $F$ . (This is sufficient as  $F$  is an elementary submodel of  $F^2$ .) Then

$$\begin{aligned} \bar{a} &= \sum_{i \leq m} d_i \bar{a}_i \text{ for some } d_i \in F^2 \text{ and} \\ U\bar{a} &= D\bar{a} = D \left( \sum_i d_i \bar{a}_i \right) = \sum_i D(d_i) \bar{a}_i + \sum_i d_i D\bar{a}_i \\ &= \sum_i (Dd_i) \bar{a}_i + \sum_i d_i (U\bar{a}_i) = \sum_i (Dd_i) \bar{a}_i + U \left( \sum_i d_i \bar{a}_i \right) \\ &= \sum (Dd_i) \bar{a}_i + U\bar{a}. \end{aligned}$$

Thus  $\sum_i (Dd_i) \bar{a}_i = 0, Dd_i \in F$ . Since the set of  $\bar{a}_i$  was linearly independent in  $F$ , and  $F$  is an elementary submodel of  $F^2$ , and since  $T_{rdc}^p$  is model-complete, the set of  $\bar{a}_i$  is linearly independent in  $F^2$ . Hence  $Dd_i = 0$ . The same holds for  $F_1, F_2$  instead of  $F^2$ .

Stage (vi). Combining the conclusions of (iv), (v), we arrive at the following representations:

$$\begin{aligned} a_i &= \sum_j \alpha_j^i d_j \text{ for } \alpha_j^i \in F, d_j \in F_1, Dd_j = 0, j < n_1. \\ b_i &= \sum_j \beta_j^i e_j \text{ for } \beta_j^i \in F, e_j \in F_2, De_j = 0, j < n_2. \end{aligned}$$

Hence

$$c = \sum \gamma_j^i d_i e_j \text{ for } \gamma_j^i \in F, d_i \in F_1, e_j \in F_2, Dd_i = De_j = 0.$$

Choose such representation with minimal  $n_1$ ; among those with minimal  $n_1$ , choose a representation with a minimal  $n_2$ . Hence the set of  $d_i$  is linearly independent over  $F$  and also the  $e_j$  are linearly independent over  $F$ .

Hence, as in Stage (iii),  $\{d_i e_j : i < n_1, j < n_2\}$  is linearly independent over  $F$ . Now since

$$0 = Dc = \sum_{i,j} (D\gamma_j^i) d_i e_j \text{ (as } Dd_i = De_j = 0)$$

and  $D\gamma_j^i \in F$ , clearly  $D\gamma_j^i = 0$ . Thus  $\gamma_j^i$  have a  $p$ -th root in  $F$ ,  $d_i$  has a  $p$ -th root in  $F_1$ , and  $e_j$  has a  $p$ -th root in  $F_2$ . Thus

$$r(c) = \sum_{i,j} r(\gamma_j^i) r(d_i) r(e_j) \in (\text{the field generated by } F_1, F_2).$$

Q.E.D.

## ACKNOWLEDGEMENT

I would like to thank Y. Kannai for suggesting that I try, for Section 1, differential equations with transcendental first-integral (when I presented him with the problem in "differential equations theory" terms); and to thank B. Weiss and Y. Hirshfeld for helpful discussions.

## ADDED IN PROOF

1. The non-minimality of the prime  $T_{dc}^0$ -field was also proved, independently by Rosenlicht [5a].
2. Wood [14], [15] also gives a nice set of axioms of  $T_{rdc}^p$ .
3. The answer to Conjecture 3 is positive.

## REFERENCES

1. L. Blum, *Generalized Algebraic Structures: Model Theoretic Approach*, Ph. D. thesis, M. I. T., 1968.
2. N. Jacobson, *Lectures in Abstract Algebra III*, Van Nostrand, Princeton, N. J., 1964.
- 2a. E. R. Kolchin, *Constrained extension of differential fields* (to appear).
3. M. D. Morley, *Categoricity in power*, Trans. Amer. Math. Soc. **114** (1965), 514–538.
4. J. F. Ritt, *Differential algebra*, Amer. Math. Soc. Colloquium Publ. **33** (1950).
5. A. Robinson, *On the concept of a differentially closed field*, Bull. Res. Council Israel, Sect. F. **8** (1959), 113–128.
- 5a. M. Rosenlicht, *The non-minimality of the differential closure*, Pacific J. Math. (to appear).
6. G. Sacks, *Saturated Model Theory*, Benjamin, Reading, Mass., 1972.
7. A. Seidenberg, *An elimination theory for differential algebra*, Univ. California Publ. Math. (new Series) **3** (1964), 31–66.
8. C. L. Siegel, *Transcendental numbers*, Annals of Math. Studies, **16** Princeton Univ. Press, 1949.
9. S. Shelah, *Differentially closed fields*, Notices Amer. Math. Soc. **20** (1973), A-444.
10. S. Shelah, *Uniqueness and characterization of prime models over sets for totally transcendental first-order theories*, J. Symbolic Logic **37** (1972), 107–113.
11. S. Shelah, *The number of non-isomorphic models of an unstable first-order theory*, Israel J. Math **9** (1971), 473–487.
12. S. Shelah, *Stability, superstability and the f. c. p. model-theoretic properties of formulas in first order theory*, Annals of Math. Logic **3** (1971), 271–362.
13. C. Wood, *The model theory of differential fields of characteristic  $p \neq 0$*  Proc. Amer. Math. Soc. **40** (1973), 577–584.
14. C. Wood, *Prime model extensions of differential fields, characteristic  $p \neq 0$* . Notices Amer. Math. Soc. **20** (1973), p. A-445.
15. C. Wood, *Prime model extensions for differential fields of characteristic  $p \neq 0$*  (to appear).