

# NEIGHBORLY POLYTOPES

BY  
IDO SHEMER

## ABSTRACT

A  $2m$ -polytope  $Q$  is *neighborly* if each  $m$  vertices of  $Q$  determine a face. It is shown that the combinatorial structure of a neighborly  $2m$ -polytope determines the combinatorial structure of every subpolytope. We develop a construction of "sewing a vertex onto a polytope", which, when applied to a neighborly  $2m$ -polytope, yields a neighborly  $2m$ -polytope with one more vertex. Using this construction, we show that the number  $g(2m + \beta, 2m)$  of combinatorial types of neighborly  $2m$ -polytopes with  $2m + \beta$  vertices grows superexponentially as  $\beta \rightarrow \infty$  ( $m \geq 2$  fixed) and as  $m \rightarrow \infty$  ( $\beta \geq 4$  fixed).

## 1. Introduction

In this paper we deal with simplicial  $k$ -neighborly  $d$ -polytopes in general, and in particular with neighborly (i.e.  $m$ -neighborly)  $2m$ -polytopes.

In sections 2 and 3 we establish general combinatorial properties of such polytopes.

In sections 4 and 5 we present and investigate a construction of "sewing" an additional vertex to a neighborly  $2m$ -polytope.

By repeated use of this construction we obtain in section 6 lower bounds for the number  $g(v, 2m)$  of combinatorial types of neighborly  $2m$ -polytopes with  $v$  vertices.  $g(2m + 4, 2m)$  increases superexponentially with  $m$ , and  $g(v, m)$  tends to infinity superexponentially with  $v$  for each fixed  $m \geq 2$ .

The techniques developed in sections 2–5 are used in [11] to prove that a non-cyclic neighborly  $2m$ -polytope with  $v \geq 2m + 5$  vertices has at most  $2m$  cyclic subpolytopes with  $v - 1$  vertices.

The notation and conventions in this paper follow [6]. In addition, we denote by  $[A_1, A_2, \dots]$  the set  $\text{conv}(A_1 \cup A_2 \cup \dots)$ , where  $A_1, A_2, \dots \subset R^d$ . If  $a \in R^d$ , then  $[\dots, a, \dots]$  stands for  $[\dots, \{a\}, \dots]$ .

All the polytopes in this paper are simplicial polytopes but not a simplex, unless otherwise specified.

Throughout the paper, the letter  $P$  denotes a  $d$ -polytope, and  $Q$  denotes a neighborly  $2m$ -polytope.

We shall use the following characterization of faces [6, sec. 2.1, th. 11]: If  $P$  is a polytope and  $T \subset \text{vert } P$ , then  $[T] \in \mathcal{F}(P)$  iff  $\text{aff } T \cap [\text{vert } P \setminus T] = \emptyset$ .

We say that a point  $x \in R^d$  covers a face  $\Phi$  of  $P$  if  $x$  lies beyond all the facets of  $P$  that include  $\Phi$  (see [6, ch. 5]).

By a subpolytope of  $P$  we mean the convex hull of a subset of  $\text{vert } P$ .

We suppose that the reader is familiar with the basic facts about neighborly and cyclic polytopes (see [6, sec. 4.7 and ch. 7], and [8, pp. 82–93]), in particular with Gale’s Evenness Condition.

### 2. Neighborly polytopes

In this section we are concerned with properties of the boundary complex of neighborly polytopes. The main result is that the combinatorial structure of an  $m$ -neighborly  $2m$ -polytope determines the combinatorial structure of every subpolytope.

LEMMA 2.1. Intersection lemma. *Let  $S_1, \dots, S_k$  be subsets of  $R^d$ . If  $\bigcap_{i=1}^k [S_i] \neq \emptyset$  and if  $a = \sum_{j=1}^r \lambda_j a_j \in \bigcap_{i=1}^k [S_i]$  where  $\lambda_{11} > 0, \lambda_{j1} \geq 0, a_j \in S_1$  ( $1 \leq j \leq r$ ),  $\sum_{j=1}^r \lambda_j = 1$ , then there are subsets  $T_i \subset S_i$  ( $1 \leq i \leq k$ ) such that*

$$(1) \quad \bigcap_{i=1}^k [T_i] \neq \emptyset,$$

$$(2) \quad \sum_{i=1}^k |T_i| \leq (k - 1)d + k,$$

$$(3) \quad a_{11} \in T_1.$$

PROOF. Let

$$V = R^{(k-1)(d+1)} = \underbrace{(R \times R^d) \times \dots \times (R \times R^d)}_{k-1 \text{ times}}$$

For  $x \in R^d$  and  $1 \leq i \leq k$  define a point  $x^i \in V$  as follows:

$$x^1 = (-1, -x, 0, \dots, 0); \quad \text{for } 2 \leq i \leq k - 1, \quad x^i = \underbrace{(0, \dots, 0, 1, x, -1, -x, 0, \dots, 0)}_{(i-2)(d+1)};$$

$$\underbrace{\hspace{10em}}_{(k-i-1)(d+1)}$$

finally,  $x^k = (0, \dots, 0, 1, x)$ . Define also  $A = \bigcup_{i=1}^k \{x^i : x \in S_i\}$ .

For  $1 \leq i \leq k$  there are points  $a_{ji} \in S_i$  and numbers  $\lambda_{ji} \geq 0, 1 \leq j \leq r_i$ , such that  $\sum_{j=1}^{r_i} \lambda_{ji} = 1$  and  $a = \sum_{j=1}^{r_i} \lambda_{ji} a_{ji}$ . Then

$$\sum_{i=1}^k \sum_{j=1}^{r_i} \frac{1}{k} \lambda_{ji} = 1, \quad \frac{1}{k} \lambda_{11} > 0 \quad \text{and} \quad \sum_{i=1}^k \sum_{j=1}^{r_i} \frac{1}{k} \lambda_{ji} a_{ji} = 0_V \in V.$$

Hence  $0_V \in \text{conv } A$ . Applying Carathéodory's theorem to the set  $A$  in  $V$  we find subsets  $T_i \subset S_i, 1 \leq i \leq k$ , such that  $a_{11} \in T_1, \sum_{i=1}^k |T_i| \leq (k-1)(d+1)+1$  and  $0_V \in \text{conv}(\bigcup_{i=1}^k \{x^i : x \in T_i\})$ . A close look at the definition of  $x^i$  reveals that  $0_V \in \text{conv}(\bigcup_{i=1}^k \{x^i : x \in T_i\})$  implies  $\bigcap_{i=1}^k \text{conv } T_i \neq \emptyset$ . □

We shall use this lemma later with  $k = 2$ .

**DEFINITION 2.2.** Let  $S$  be a subset of  $\text{vert } P$ .  $[S]$  is a *missing face* (m.f.) of  $P$  if  $[S]$  is not a face of  $P$  but for every proper subset  $T$  of  $S, [T]$  is a face of  $P$ . (Compare [3, def. 2.1].)

We say that  $[S]$  is a *missing  $k$ -face* ( $k$ -m.f.) of  $P$  if  $[S]$  is a m.f. of  $P$  and  $\dim [S] = k$ .

**REMARK.** Definition 2.2 remains meaningful even if we drop the assumption that the polytope  $P$  is simplicial. Also, Theorems 2.3, 2.4, 2.5 and their proofs hold for general polytopes.

**THEOREM 2.3.** If  $S \subset V = \text{vert } P$ , then  $[S]$  is a m.f. of  $P$  iff

- (1)  $P \cap \text{aff } S = [S],$
- (2)  $\emptyset \neq [S] \cap [V \setminus S] \subset \text{relint } [S],$
- (3)  $|S| = \dim [S] + 1.$

**PROOF.** Suppose  $[S]$  is a m.f. of  $P$ . Clearly  $[S] \subset P \cap \text{aff } S$ . If  $x \in P \cap \text{aff } S \setminus [S]$ , then there are points  $y \in \text{relint } [S], z \in \text{relbd } [S]$  such that  $z \in (x, y)$ . There is a proper face  $F$  of  $[S]$  such that  $z \in F$ .  $F$  is a face of  $P$ . Then  $y \in P \setminus F, z \in F$ , hence  $x \notin P$ , a contradiction. Hence  $P \cap \text{aff } S = [S]$ .  $[S]$  is not a face of  $P$ , hence  $\text{aff } S \cap [V \setminus S] \neq \emptyset$ . So  $[S] \cap [V \setminus S] = P \cap \text{aff } S \cap [V \setminus S] = \text{aff } S \cap [V \setminus S] \neq \emptyset$ .

If  $T \subsetneq S$  then  $[T]$  is a face of  $P$ , hence a face of  $[S]$ . So  $[S]$  is a simplex and  $\dim [S] = |S| - 1$ .

If  $x \in \text{relbd } [S]$ , then  $x \in [T]$  for some  $T \subsetneq S$ .  $[T] \in \mathcal{F}(P)$ , hence  $[T] \cap [V \setminus T] = \emptyset$ . So  $x \notin [V \setminus T]$  and *a fortiori*  $x \notin [V \setminus S]$ . Therefore  $[V \setminus S] \cap \text{relbd } [S] = \emptyset$ , hence  $[V \setminus S] \cap [S] \subset \text{relint } [S]$ .

Now we turn to the converse part of the theorem. Suppose  $S \subset V$  satisfies (1), (2), (3). To prove that  $[S]$  is a m.f. of  $P$  it clearly suffices to show that if  $T \subsetneq S$  and  $|T| = |S| - 1$ , then  $[T] \in \mathcal{F}(P)$ . Let  $x \in S, T = S \setminus \{x\}$ .

(A)  $P \cap \text{aff } T = P \cap \text{aff } S \cap \text{aff } T = [S] \cap \text{aff } T = [T]$ .

(B) We claim that  $[T] \cap [V \setminus T] = \emptyset$ . If not, then there is a point  $a \in [T] \cap [V \setminus S, x]$ . There is a point  $b \in [V \setminus S]$  such that  $a \in [b, x]$ .  $b \in \text{aff } \{a, x\}$  since  $a \neq x$ . It follows that  $b \in \text{aff } S \cap [V \setminus S] = [S] \cap [V \setminus S] \subset \text{relint } [S]$ . Since  $a \in [b, x]$ ,  $a \neq x$ , we obtain  $a \in \text{relint } [S]$ , a contradiction. From (A) and (B) we get

$$\text{aff } T \cap [V \setminus T] = P \cap \text{aff } T \cap [V \setminus T] = [T] \cap [V \setminus T] = \emptyset. \quad \square$$

**THEOREM 2.4.** *A  $k$ -neighborly  $d$ -polytope  $P$  has no  $j$ -m.f. with  $j < k$  or  $j > d - k$ .*

**PROOF.** It is obvious that  $P$  has no  $j$ -m.f. with  $j < k$ . Suppose  $S \subset \text{vert } P = V$  and  $[S]$  is a  $j$ -m.f. of  $P$ . Then  $[S] \cap [V \setminus S] \neq \emptyset$  (Theorem 2.3). Applying the intersection Lemma 2.1 we find subsets  $T_1 \subset S$ ,  $T_2 \subset V \setminus S$  such that  $[T_1] \cap [T_2] \neq \emptyset$  and  $|T_1| + |T_2| \leq d + 2$ .  $T_1 = S$ , because otherwise  $[T_1]$  is a face of  $P$ , and then

$$[T_1] \cap [T_2] \subset [T_1] \cap [V \setminus T_1] = \emptyset.$$

$|T_2| \geq k + 1$ , because  $P$  is  $k$ -neighborly. Therefore  $j + 1 = |S| = |T_1| \leq (d + 2) - |T_2| \leq (d + 2) - (k + 1)$ ,  $j \leq d - k$ . □

An immediate consequence of Theorem 2.4 is the well-known fact:

**THEOREM 2.5.** *If  $P$  is a  $k$ -neighborly  $d$ -polytope and  $k > \frac{1}{2}d$ , then  $P$  is a simplex.*

There are many problems concerning the determination of  $\mathcal{F}(P)$  from some partial information about  $\mathcal{F}(P)$ . A natural question is: under what circumstances does  $\text{skel}_j \mathcal{F}(P)$  determine  $\mathcal{F}(P)$ ?

**THEOREM 2.6.** *If  $P$  is a simplicial  $k$ -neighborly  $d$ -polytope, then  $\mathcal{F}(P)$  is determined by its  $(d - k)$ -skeleton; moreover, if  $S \not\subseteq \text{vert } P$  and  $|S| > d - k + 1$ , then  $[S] \in \mathcal{F}(P)$  iff  $[T] \in \mathcal{F}(P)$  for every  $T \subset S$  with  $|T| \leq d - k + 1$ .*

**PROOF.** Assume  $S \not\subseteq \text{vert } P$ ,  $|S| \geq d - k + 2$ . If  $[S] \in \mathcal{F}(P)$ , then  $T \subset S$  implies  $[T] \in \mathcal{F}(P)$ , since  $P$  is simplicial. If  $[S] \notin \mathcal{F}(P)$ , then there is a set  $T \subset S$ , such that  $[T]$  is m.f. By Theorem 2.4,  $|T| \leq d - k + 1$ . □

An immediate consequence is

**THEOREM 2.7.** *If  $Q$  is an  $m$ -neighborly  $2m$ -polytope then  $\text{skel}_m Q$  determines  $\mathcal{F}(Q)$ .*

It is known, though much harder to prove, that the combinatorial structure of every simplicial polytope  $P$  of dimension  $2m$  or  $2m + 1$  is determined by  $\text{skel}_m \mathcal{F}(P)$ .

**THEOREM 2.8.** *Let  $P, P^+$  be  $k$ -neighborly  $d$ -polytopes, not necessarily simplicial. Suppose  $\text{vert } P = V, \text{vert } P^+ = V \cup \{x\}, x \notin P, T \subset V, \dim [T] \geq d - k$ . If for every  $S \subsetneq T, [S, x] \in \mathcal{F}(P^+)$ , then  $[T] \subset \text{bd } P$ .*

**REMARK.** The assumption, that  $[S, x] \in \mathcal{F}(P^+)$  for all  $S \subsetneq T$ , implies that  $T \cup \{x\}$  is affinely independent and therefore  $\dim [T] = |T| - 1$ .

**PROOF.** If  $S \subsetneq T$  then  $[S, x] \in \mathcal{F}(P^+)$ , hence  $[S] \in \mathcal{F}(P)$ . Therefore  $[T]$  is either a face or a m.f. of  $P$ . Assume  $[T]$  is a m.f. of  $P$ . Hence  $\dim [T] \leq d - k$  (Theorem 2.4). But  $\dim [T] \geq d - k$ , hence  $\dim [T] = d - k$  and  $|T| = d - k + 1$ . If  $[V \setminus T] \cap \text{relint}[x, T] \neq \emptyset$  then from the intersection Lemma 2.1 we obtain subsets  $S \subset T$  and  $R \subset V \setminus T$  that  $[R] \cap [x, S] \neq \emptyset$  and  $|R| + |S| + 1 \leq d + 2$ .  $[R] \notin \mathcal{F}(P^+)$ , hence  $|R| \geq k + 1$ .  $|S| \leq d + 1 - |R| \leq d - k$ . Hence  $S \subsetneq T$ .  $[x, S] \in \mathcal{F}(P^+)$  contradicts  $[x, S] \cap [R] \neq \emptyset$ . Therefore  $[V \setminus T] \cap \text{relint}[x, T] = \emptyset$ .

Let  $H$  be a hyperplane that separates  $V \setminus T$  from  $[x, T]$ ,  $V \setminus T \subset H^-, T \cup \{x\} \subset H^+$ .  $[T]$  is a m.f. of  $P$ , hence there is a point  $a \in \text{relint}[T] \cap [V \setminus T]$  (Theorem 2.3). It follows that  $a \in H$ , hence  $T \subset H$ . So  $V \subset H^-, [T] \subset \text{bd } P$ .  $\square$

**COROLLARY 2.9.** *If  $P$  is a simplicial polytope then the hypotheses of Theorem 2.8 imply  $[T] \in \mathcal{B}(P) = \mathcal{F}(P) \setminus \{P\}$ .*

We conclude this section with the following result: The combinatorial structure of an  $m$ -neighborly  $2m$ -polytope determines the combinatorial structure of every subpolytope. A somewhat different proof of this result appears in [10].

**THEOREM 2.10.** *Let  $Q, Q^+$  be  $m$ -neighborly  $2m$ -polytopes,  $\text{vert } Q^+ = \text{vert } Q \cup \{x\}, x \notin Q$ . Then  $\mathcal{F}(Q^+)$  determines  $\mathcal{F}(Q)$ .*

Theorem 2.10 is an immediate consequence of Theorem 2.7 and the following lemma:

**LEMMA 2.11.** *Under the assumption of Theorem 2.10,  $\text{skel}_m \mathcal{F}(Q^+)$  determines  $\text{skel}_m \mathcal{F}(Q)$ ; moreover, if  $T \subset \text{vert } Q, |T| = m + 1$ , then  $[T] \in \mathcal{F}(Q)$  iff either*

- (1)  $[T] \in \mathcal{F}(Q^+)$ , or
- (2)  $[T] \notin \mathcal{F}(Q^+)$ , but  $[x, S] \in \mathcal{F}(Q^+)$  for every  $S \subsetneq T$ .

PROOF. Let  $T \subset \text{vert } Q, |T| = m + 1$ . If (1) holds then obviously  $[T] \in \mathcal{F}(Q)$ . If (2) holds then Corollary 2.9 implies  $[T] \in \mathcal{F}(Q)$ . Conversely, if  $[T] \in \mathcal{F}(Q)$  but  $[T] \notin \mathcal{F}(Q^+)$  then  $x$  lies beyond all the facets of  $Q$  that include  $T$ . Therefore, if  $S \subset T$ , then  $x$  lies beyond some facet of  $P$  that includes  $S$ . But if  $S \subsetneq T$ , then  $[S] \in \mathcal{F}(Q^+)$ . Therefore  $x$  lies beneath at least one facet of  $P$  that includes  $S$ , hence  $[S, x] \in \mathcal{F}(Q^+)$  (see [6, section 5.2]).  $\square$

The following alternative formulation of Theorem 2.10 will be useful in the sequel:

THEOREM 2.12. *Let  $Q_1, Q_2$  be  $m$ -neighborly  $2m$ -polytopes and let the bijection  $\varphi : \text{vert } Q_1 \rightarrow \text{vert } Q_2$  be a combinatorial equivalence between  $Q_1$  and  $Q_2$ .*

*If  $A \subset \text{vert } Q_1$ , then the restriction of  $\varphi$  to  $A$  is a combinatorial equivalence between  $[A]$  and  $[\varphi(A)]$ .*

### 3. Universal faces

DEFINITION 3.1. Suppose  $\Phi \in \mathcal{B}(P) (= \mathcal{F}(P) \setminus \{P\})$ .  $\Phi$  is a  $u$ -universal face ( $u$ -u.f.) of  $P$  if  $[\Phi, S] \in \mathcal{B}(P)$  for every  $S \subset \text{vert } P$  with  $|S| \leq u$ .

Define  $\mathcal{B}(P, u) = \{\Phi \in \mathcal{B}(P) : \Phi \text{ is a } u\text{-u.f. of } P\}$ . Note that  $\mathcal{B}(P, 0) = \mathcal{B}(P)$ , and  $\emptyset \in \mathcal{B}(P, k)$  iff  $P$  is  $k$ -neighborly.

DEFINITION 3.2.  $\Phi$  is a universal face (u.f.) of  $P$  if  $\Phi \in \mathcal{B}(P, u)$  with  $u = \lfloor \frac{1}{2}(d - \dim \Phi - 1) \rfloor = \lfloor \frac{1}{2}(d - |\text{vert } \Phi|) \rfloor$ . A 1-dimensional u.f. is called a universal edge (u.e.).

Definitions 3.1 and 3.2 can be reformulated using the notion of a quotient polytope  $P/\Phi$  introduced in [8, ch. 2, th. 16]. (A quotient polytope  $P/\Phi$  is a polytope  $K$  whose face lattice  $\mathcal{F}(K)$  is isomorphic to the upper segment  $[\Phi, P]$  of  $\mathcal{F}(P)$ .)

DEFINITION 3.1\*.  $\Phi \in \mathcal{B}(P, u)$  iff either (a)  $u = 0$  and  $\Phi \in \mathcal{B}(P)$ , or (b)  $u > 0$  and the quotient polytope  $P/\Phi$  is  $u$ -neighborly, with  $|\text{vert } P| - |\text{vert } \Phi|$  vertices.

REMARK.  $\dim P/\Phi = \dim P - \dim \Phi - 1 = \dim P - |\text{vert } \Phi|$ . Since we assume that  $P$  is not a simplex,  $|\text{vert } P| > \dim P + 1$ . Therefore, if  $\Phi \in \mathcal{B}(P, u)$ ,  $u > 0$ , then  $|\text{vert } P/\Phi| = |\text{vert } P| - |\text{vert } \Phi| > \dim P/\Phi + 1$ , i.e.,  $P/\Phi$  is not a simplex, and therefore  $u \leq \frac{1}{2} \dim P/\Phi \leq \frac{1}{2} \dim P$ .

DEFINITION 3.2\*.  $\Phi$  is a u.f. of  $P$  iff either (a)  $\Phi$  is a facet of  $P$ , or (b)  $\Phi \in \mathcal{B}(P)$  and  $P/\Phi$  is a neighborly (i.e.,  $\lfloor \frac{1}{2} \dim P/\Phi \rfloor$ -neighborly) polytope with  $|\text{vert } P| - |\text{vert } \Phi|$  vertices.

One can use Definitions 3.1 and 3.2 in the case where  $P$  is a simplex. The various remarks that follow these definitions remain valid if we adhere to the convention that a simplex  $\Delta$  is  $u$ -neighborly for  $0 \leq u \leq \dim \Delta$  only.

**THEOREM 3.3.** *Let  $P$  be a  $k$ -neighborly  $d$ -polytope. If  $\Psi_i \in \mathcal{B}(P, u_i)$  for  $i = 1, 2$  and  $t = u_1 + u_2 + k - d \geq 0$ , then  $\Psi = [\Psi_1, \Psi_2] \in \mathcal{B}(P, t)$ .*

**PROOF.** Assume  $A \subset \text{vert } P \setminus \Psi$  and  $[\Psi, A] \notin \mathcal{B}(P)$ . It suffices to show that  $|A| > t$ .

There are three pairwise disjoint sets  $B \subset A$ ,  $T_1 \subset \text{vert } \Psi_1$ ,  $T_2 \subset \text{vert } \Psi_2$  such that  $[T_1, T_2, B]$  is a m.f. of  $P$ . By Theorem 2.4,  $|T_1| + |T_2| + |B| \leq d - k + 1$ .  $[T_1, \Psi_2, B] \notin \mathcal{B}(P)$ , hence  $|T_1| + |B| \geq u_2 + 1$ . By the same reasoning  $|T_2| + |B| \geq u_1 + 1$ . Therefore  $d - k + 1 + |B| \geq |T_1| + |T_2| + 2|B| \geq u_1 + u_2 + 2$ , hence  $|A| \geq |B| \geq u_1 + u_2 + k - d + 1 = t + 1$ .  $\square$

The next theorem is a very useful special case of Theorem 3.3. From this point onward, the letter  $Q$  will always denote an  $m$ -neighborly  $2m$ -polytope.

**THEOREM 3.4.** *Let  $\Psi_1, \dots, \Psi_l$  be pairwise disjoint universal faces of  $Q$ . If  $\Psi_i$  has  $2v_i$  vertices ( $1 \leq i \leq l$ ) and  $v_1 + \dots + v_l \leq m$ , then  $\Psi = [\Psi_1, \dots, \Psi_l]$  is a u.f. of  $Q$ .*

**REMARK.** If  $\Phi$  is a face of  $Q$  with  $2j$  vertices, then  $\Phi$  is a u.f. iff  $\Phi$  is an  $(m - j)$ -u.f.

**PROOF.** Theorem 3.4 in its full generality follows from the case  $l = 2$  by induction on  $l$ . For  $l = 2$ , apply Theorem 3.3 with  $d = 2m$ ,  $k = m$ ,  $u_1 = m - v_1$ ,  $u_2 = m - v_2$ .  $\square$

Let  $C = C(v, 2m)$  be a cyclic  $2m$ -polytope with  $v$  vertices. There is a natural cyclic order on  $\text{vert } C$ . Assume that  $a_1, a_2, \dots, a_v, a_1$  are the vertices of  $C$  in this order. From Gale's Evenness Condition it follows that  $[a_i, a_{i+1}]$  ( $1 \leq i < v$ ) and  $[a_v, a_1]$  are u.e.s of  $C$ , and if  $v \geq 2m + 3$  then  $C(v, 2m)$  has no other u.e.s. See also [3, section 3]. The existence of a hamiltonian circuit of u.e.s characterizes cyclic polytopes:

**THEOREM 3.5.** *Assume  $|\text{vert } Q| \geq 2m + 3$ . If  $Q$  has a simple circuit of length  $v$  consisting of universal edges, then  $Q \cong C(v, 2m)$ .*

**PROOF.** *Case I:  $v < 2m + 3$ .*

Choose  $2m + 3 - v$  vertices of  $Q$  which are not in the given circuit, and add them to the vertices of the circuit. The resulting set spans a subpolytope  $Q'$  of  $Q$ .  $Q'$  is neighborly, hence  $Q' \cong C(2m + 3, 2m)$  (see [6, th. 7.2.3]). The edges of the

given circuit are u.e.s in  $Q'$ , but  $Q'$ , being a cyclic polytope with  $2m + 3$  vertices, has only one simple circuit of u.e.s of length  $2m + 3$ , a contradiction.

*Case II:*  $v \geq 2m + 3$ .

Suppose that the vertices of  $C(v, 2m)$ , in their natural cyclic order, are  $a_1, \dots, a_v, a_1$ . Let  $\varphi$  be a 1 : 1 mapping of vert  $C(v, 2m)$  into vert  $Q$ , which maps the cycle  $a_1, \dots, a_v, a_1$  onto the given circuit of universal edges of  $Q$ . Gale's Evenness Condition and Theorem 3.4 imply that  $\varphi$  induces a 1 : 1 mapping of the set of facets of  $C(v, 2m)$  into the set of facets of  $Q$ . By Lemma 3.6 below,  $\varphi$  is a combinatorial equivalence between  $C(v, 2m)$  and  $Q$ . □

LEMMA 3.6. *Suppose  $\varphi : \text{vert } P \rightarrow \text{vert } P'$  is a bijection. If for every facet  $F$  of  $P$ ,  $F' = [\varphi(\text{vert } F)]$  is a facet of  $P'$ , then  $\varphi$  is a combinatorial equivalence between  $P$  and  $P'$ .*

PROOF. Since the incidence graph of facets and subfacets of the polytope  $P'$  is connected, it is enough to verify the following assertion:

If  $F', G'$  are two adjacent facets of  $P'$  (i.e.,  $F' \cap G'$  is a subfacet of  $P'$ ) and if  $F' = \varphi(F)$  for some facet  $F$  of  $P$ , then  $G' = \varphi(G)$  for some facet  $G$  of  $P$ .

Indeed, consider the set  $\varphi^{-1}(\text{vert } (F' \cap G'))$ . Since  $P$  and  $P'$  are simplicial, this is the set of vertices of a subfacet  $H$  of  $P$ .  $H$  is included in precisely two different facets  $F, G$  of  $P$ . Since  $\varphi(F) = F'$ ,  $\varphi(G)$  must be  $G'$ . □

REMARK. Lemma 3.6 holds even without assuming that  $P$  and  $P'$  are simplicial. The proof in the general case is slightly more involved, and uses induction on  $\dim P$ .

As we mentioned above, every proper face is a 0-u.f. The next two theorems can be considered as generalizations of Theorem 2.6, and Corollary 2.9 to Theorem 2.8.

THEOREM 3.7. *Assume  $P$  is a  $k$ -neighborly  $d$ -polytope,  $S \subsetneq \text{vert } P$ .  $[S]$  is a  $u$ -u.f. of  $P$  iff  $[T]$  is a  $u$ -u.f. of  $P$  for every  $T \subset S$  with  $|T| \leq d - k - u + 1$ .*

PROOF. If  $[S] \notin \mathcal{B}(P, u)$  then there is a set  $A \subset \text{vert } P \setminus S$ ,  $|A| \leq u$  with  $[S, A] \notin \mathcal{B}(P)$ . There are sets  $T \subset S, B \subset A$  such that  $[T, B]$  is a m.f. of  $P$ , hence  $|T| + |B| \leq d - k + 1$  (Theorem 2.4).

*Case I:*  $|T| \leq d - k - u + 1$ .

Then  $[T] \notin \mathcal{B}(P, u)$  a contradiction.

*Case II:*  $|T| > d - k - u + 1$ .



Suppose  $R \subset T$ ,  $|R| = |T| - (d - k - u + 1)$ . Then  $|T \setminus R| = d - k - u + 1$ ,  $T \setminus R \subset S$  and  $[T \setminus R] \notin \mathcal{B}(P, u)$  since  $|B \cup R| = |B| + |R| = |B| + |T| - (d - k - u + 1) \leq (d - k + 1) - (d - k - u + 1) = u$ .  $\square$

**THEOREM 3.8.** *Assume  $P, P^+$  are  $k$ -neighborly  $d$ -polytopes,  $\text{vert } P^+ = \text{vert } P \cup \{x\}$ ,  $x \notin P$ ,  $\emptyset \neq A \subset \text{vert } P$  and  $|A| > d - k - u$ .*

*If  $[A \setminus \{q\}, x] \in \mathcal{B}(P^+, u)$  for every  $q \in A$  then  $[A] \in \mathcal{B}(P, u)$ .*

**PROOF.** We have to show that if  $W \subset \text{vert } P \setminus A$ ,  $|W| \leq u$ , then  $[A, W] \in \mathcal{B}(P)$ .

If  $|W| < u$ , choose any  $q \in A$ ; then  $[A, W, x] = [A \setminus \{q\}, x, W \cup \{q\}] \in \mathcal{B}(P^+)$ , hence  $[A, W] \in \mathcal{B}(P)$ . Now assume that  $|W| = u$ . Then  $|A \cup W| = |A| + u > d - k$ . By Corollary 2.9 it suffices to prove that  $[S, x] \in \mathcal{B}(P^+)$  for all  $S \subsetneq A \cup W$ .

Assume  $S \subsetneq A \cup W$ . If  $S \not\subset A$  choose a point  $q \in A \setminus S$ ; then  $[A \setminus \{q\}, x, W] \in \mathcal{B}(P^+)$ ,  $S \subset (A \setminus \{q\}) \cup W$ , hence  $[S, x] \in \mathcal{B}(P^+)$ . If  $S \subset A$ , then  $W \cap S \subsetneq W$ . Choose any  $q \in A$ ; then  $[S, x] = [A \setminus \{q\}, x, (W \cap S) \cup \{q\}] \in \mathcal{B}(P^+)$ .  $\square$

**COROLLARY 3.9.** *Let  $Q, Q^+$  be neighborly  $2m$ -polytopes,  $\text{vert } Q^+ = \text{vert } Q \cup \{x\}$ ,  $x \notin Q$ . If  $\emptyset \neq A \subset \text{vert } Q$ ,  $|A| \geq 2$  and if  $[A \setminus \{q\}, x]$  is a u.f. of  $Q^+$  for every  $q \in A$ , then  $[A]$  is a u.f. of  $Q$ .*

**PROOF.**  $|A| = 2j$  with  $j \geq 1$  or  $|A| = 2j - 1$  with  $j \geq 2$ . In both cases apply Theorem 3.8 with  $d = 2m$ ,  $k = m$ ,  $u = m - j$ .  $\square$

We shall use this corollary later with  $|A| = 2$ .

**THEOREM 3.10.** *If  $|\text{vert } Q| \geq 2m + 3$ , then the graph of the universal edges of  $Q$  is either a hamiltonian circuit, or a union of disjoint simple paths.*

**PROOF.** In view of Theorem 3.5, it is sufficient to prove that no vertex of  $Q$  is included in three u.e.s of  $Q$ .

Suppose  $x, a, b, c$  are distinct vertices of  $Q$ , and  $[x, a], [x, b], [x, c]$  are u.e.s of  $Q$ . Choose a set  $A$  of  $2m - 1$  vertices of  $Q$  other than  $x, a, b, c$ , and define  $Q' = [x, a, b, c, A]$ .  $Q'$  is a neighborly  $2m$ -polytope with  $2m + 3$  vertices, hence  $Q' \cong C(2m + 3, 2m)$ .  $[x, a], [x, b], [x, c]$  are u.e.s of  $Q'$ , but no vertex of  $C(2m + 3, 2m)$  is included in three u.e.s, a contradiction.  $\square$

#### 4. The sewing construction

In this section we describe a construction, called sewing, and some related notions. This construction, first introduced by the author in [9], will play a central role in the sequel.

The “facet-splitting” operation of Barnette [5] is, in a sense, dual to our sewing construction. We shall discuss the relationship between these two constructions in section 7.4.

**DEFINITION 4.1.** If  $\Psi \in \mathcal{F}(P)$  and  $M \subset \text{vert } P \setminus \Psi$ , then we say that  $M$  is a *missing face of  $P$  relative to  $\Psi$*  if  $[M, \Psi] \notin \mathcal{B}(P)$ , but  $[M', \Psi] \in \mathcal{B}(P)$  for every  $M' \subsetneq M$ .

Define:  $\mathcal{M}(P/\Psi) = \{M : M \text{ is a m.f. of } P \text{ relative to } \Psi\}$ . Finally define:  $\mathcal{M}(P) = \mathcal{M}(P/\emptyset)$ .

**PROPOSITION 4.2.** (1)  $\mathcal{M}(P/P) = \{\emptyset\}$ .

(2) If  $F$  is a facet of  $P$ , then  $\mathcal{M}(P/F) = \{\{q\} : q \in \text{vert } P \setminus F\}$ .

(3)  $M \in \mathcal{M}(P)$  iff  $[M]$  is a m.f. of  $P$ .

(4) If  $M \in \mathcal{M}(P/\Psi)$ , then there is a set  $S$  in  $\mathcal{M}(P)$  such that  $M \cup \Psi \supset S \supset M$ .

(5) If  $\Psi \in \mathcal{F}(P)$ , then for every vertex  $b$  in  $\text{vert } P \setminus \Psi$  there are sets  $M$  and  $N$  in  $\mathcal{M}(P/\Psi)$  such that  $b \in M$  and  $b \notin N$ .

If  $Q$  is a neighborly  $2m$ -polytope then the following properties hold too:

(6) If  $M \in \mathcal{M}(Q)$  then  $|M| = m + 1$ .

(7) If  $\Psi$  is a u.f. of  $Q$ ,  $|\text{vert } \Psi| = 2j$  and  $M \in \mathcal{M}(Q/\Psi)$ , then  $|M| = m - j + 1$ .

(8)  $\Psi$  is a u-u.f. of  $Q$  iff  $|M \cap \Psi| \leq m - u$  for every  $M \in \mathcal{M}(Q)$ , or equivalently, if  $|M \setminus \Psi| \geq u + 1$  for every  $M \in \mathcal{M}(Q)$ .

*In particular*

(9)  $E$  is a u.e. of  $Q$  iff no element of  $\mathcal{M}(Q)$  includes  $\text{vert } E$ .

**PROOF.** (1)–(4) follow immediately from Definitions 4.1 and 2.2. In order to establish (5), take  $M = \{b\} \cup C$ , where  $C$  is a minimal subset (with respect to inclusion) of  $\text{vert } P \setminus (\Psi \cup \{b\})$  such that  $[b, C, \Psi] \notin \mathcal{B}(P)$  but  $[C, \Psi] \in \mathcal{B}(P)$ , and take  $N$  to be a minimal subset of  $\text{vert } P \setminus (\Psi \cup \{b\})$  such that  $[N, \Psi] \notin \mathcal{B}(P)$ .

(6) and (7) follow from Theorem 2.4, and (6) implies (8) and (9). □

**DEFINITION 4.3.** A *tower in  $P$*  is a strictly increasing sequence  $\mathcal{T} = \{\Phi_j\}_{j=1}^k$  of non-empty proper faces of  $P$ . Sometimes we shall adjoin the empty face as a first element  $\Phi_0$  of  $\mathcal{T}$ . If  $\Phi \in \mathcal{F}(P)$ , denote by  $\mathcal{F}_\Phi$  the set of all facets of  $P$  which include  $\Phi$ . We denote  $\mathcal{F}_\Phi$  by  $\mathcal{F}_j$  (in order to avoid double subscripts). Note that  $\mathcal{F}_1 \supset \mathcal{F}_2 \supset \dots \supset \mathcal{F}_k$ . Define  $\mathcal{C} = \mathcal{C}(P, \mathcal{T}) = \mathcal{F}_1 \setminus (\mathcal{F}_2 \setminus (\dots \setminus \mathcal{F}_k \dots))$ . It is easy to see that  $\mathcal{C} = (\mathcal{F}_1 \setminus \mathcal{F}_2) \cup (\mathcal{F}_3 \setminus \mathcal{F}_4) \cup \dots$  where the last term in the union is  $\mathcal{F}_{k-1} \setminus \mathcal{F}_k$  if  $k$  is even or  $\mathcal{F}_k$  if  $k$  is odd. With the convention that  $\mathcal{F}_j = \emptyset$  and  $\Phi_j = P$  for  $j > k$  we can simply write  $\mathcal{C} = \bigcup_{i=1}^\infty (\mathcal{F}_{2i-1} \setminus \mathcal{F}_{2i})$  and similarly  $\mathcal{F}_0 \setminus \mathcal{C} = \bigcup_{i=0}^\infty (\mathcal{F}_{2i} \setminus \mathcal{F}_{2i+1})$ , where  $\mathcal{F}_0$  is the set of all facets of  $P$ .

We say that  $\mathcal{T} = \{\Phi_j\}_{j=1}^m$  is a *universal tower (u.t.) in  $Q$*  if

- (1)  $Q$  is a neighborly  $2m$ -polytope,
- (2)  $\Phi_j$  is a u.f. of  $Q$  for  $1 \leq j \leq m$ ,
- (3)  $|\text{vert } \Phi_j| = 2j$  for  $1 \leq j \leq m$ .

Let  $\mathcal{D}$  be a set of facets of  $P$ . We say that a point  $x \in R^d$  lies exactly beyond  $\mathcal{D}$  with respect to  $P$  if  $x$  lies beyond every facet of  $P$  that is in  $\mathcal{D}$  and beneath every other facet of  $P$ . If it is clear from the context what is the polytope  $P$ , we omit the phrase “with respect to  $P$ ”.

LEMMA 4.4. *Let  $\mathcal{T}$  be a tower in  $P$ ,  $\mathcal{C} = \mathcal{C}(P, \mathcal{T})$ . Then there is a point  $x \in R^d$  which lies exactly beyond  $\mathcal{C}$ .*

PROOF. By induction on the height  $k$  of  $\mathcal{T}$ . If  $k = 0$ , define  $\mathcal{C} = \emptyset$ . In that case every point  $x \in \text{int } P$  lies exactly beyond  $\mathcal{C}$ . If  $k \geq 1$ , let  $\mathcal{T}' = \mathcal{T} \setminus \{\Phi_1\}$  and  $\mathcal{C}' = \mathcal{C}(P, \mathcal{T}')$ . By the induction hypothesis, there is a point  $x' \in R^d$ , which lies exactly beyond  $\mathcal{C}'$ . Note that  $\mathcal{C}' \subset \mathcal{F}_1$  and  $\mathcal{C} = \mathcal{F}_1 \setminus \mathcal{C}'$ . Choose a point  $p \in \text{relint } \Phi_1$  and let  $x = (1 + \varepsilon)p - \varepsilon x'$ . If  $\varepsilon$  is positive and sufficiently small, then  $x$  lies exactly beyond  $\mathcal{C}$ . □

The construction which we have just described will enable us to construct a large variety of neighborly polytopes by adding new vertices to existing neighborly polytopes.

From here until the end of section 4 we adhere to the following convention:

CONVENTION 4.5.  $Q$  is a neighborly  $2m$ -polytope,  $Q$  is not simplex,  $\mathcal{T} = \{\Phi_j\}_{j=1}^m$  is a u.t. in  $Q$ ,  $\mathcal{C} = \mathcal{C}(Q, \mathcal{T})$ ,  $x$  lies exactly beyond  $\mathcal{C}$  with respect to  $Q$ , and  $Q^+ = [Q, x]$ .

Define  $S_j = \text{vert } \Phi_j \setminus \Phi_{j-1}$  for  $j = 1, 2, \dots, m + 1$  (recall that  $\Phi_0 = \emptyset$  and  $\Phi_j = Q$  for  $j > m$ ).

THEOREM 4.6. (1)  $Q^+$  is a simplicial  $2m$ -polytope and  $\text{vert } Q^+ = \text{vert } Q \cup \{x\}$ .

- (2)  $Q^+$  is neighborly.
- (3) If  $0 < j \leq m$  is even, then  $\Phi_j$  is a u.f. of  $Q^+$ .
- (4) If  $0 < j \leq m$  is odd, then  $\Phi_j$  is not a u.f. of  $Q^+$ , but if  $j < m$  then  $\Phi_j$  is still a face of  $Q^+$ .
- (5) If  $a \in S_j$  for some  $1 \leq j \leq m$ , then  $[\Phi_{j-1}, a, x]$  is a u.f. of  $Q^+$ .

LEMMA 4.7. (1)  $\text{vert } Q^+ = \text{vert } Q \cup \{x\}$ .

(2) If  $M \subset \text{vert } Q \cup \{x\}$ , then  $M \in \mathcal{M}(Q^+)$  iff either (a)  $M = \bigcup_{\nu=1}^j S_{2\nu-1} \cup A$  for some integer  $0 \leq j \leq (m + 1)/2$  and some  $A \in \mathcal{M}(Q/\Phi_{2j})$ , or (b)  $M =$

$\bigcup_{\nu=1}^j S_{2\nu} \cup A \cup \{x\}$  for some integer  $0 \leq j \leq m/2$  and some set  $A \in \mathcal{M}(Q/\Phi_{2j+1})$ .  
 (Note that  $\Phi_{m+1} = Q$  and  $\mathcal{M}(Q, Q) = \{\emptyset\}$ .)

PROOF. *Step 1:* We show that if  $M$  is of type (a), then  $[M] \notin \mathcal{B}(Q^+)$ .

Assume  $M = \bigcup_{\nu=1}^j S_{2\nu-1} \cup A$ ,  $A \in \mathcal{M}(Q/\Phi_{2j})$ ,  $0 \leq j \leq (m+1)/2$ . If  $j = 0$ , then  $[M] = [A] \notin \mathcal{B}(Q)$ . Assume  $j > 0$ , hence  $S_1 \subset M$ .

Let  $F$  be a facet in  $\mathcal{F}_{[M]}$ , and let  $\mu$  be the maximal integer  $\nu$  such that  $1 \leq \nu \leq j$  and  $F \in \mathcal{F}_{2\nu-1}$ . Then  $F \notin \mathcal{F}_{2\mu}$ , since otherwise, if  $\mu < j$  then  $F \in \mathcal{F}_{2\mu+1}$ , because  $S_{2\mu+1} \subset M \subset F$ , and if  $\mu = j$  then  $F \supset A \cup \Phi_{2j}$ , in contradiction to  $A \in \mathcal{M}(Q/\Phi_{2j})$ . Therefore  $\mathcal{F}_{[M]} \subset \bigcup_{\nu=1}^j (\mathcal{F}_{2\nu-1} \setminus \mathcal{F}_{2\nu}) \subset \mathcal{C}$ . It follows that if  $M$  is of type (a), then  $[M] \notin \mathcal{B}(Q^+)$ .

*Step 2:* We show that if  $M$  is of type (b), then  $[M] \notin \mathcal{B}(Q^+)$ . Define  $M^- = M \setminus \{x\}$ . The rest of step 2 is similar to step 1: we prove that  $\mathcal{F}_{[M^-]} \subset \bigcup_{\nu=0}^j (\mathcal{F}_{2\nu} \setminus \mathcal{F}_{2\nu+1}) \subset \mathcal{F}_0 \setminus \mathcal{C}$  and conclude that  $[M] \notin \mathcal{B}(Q^+)$ .

*Step 3:* Now we show that if  $S \subset \text{vert } Q$  and  $[S] \notin \mathcal{B}(Q^+)$ , then  $S$  includes a set  $M$  of type (a). Since  $[S] \notin \mathcal{B}(Q^+)$ , it follows that if  $F \in \mathcal{F}_0$  and  $F \supset S$ , then  $F \in \mathcal{C}$ .

Let  $j$  be the first nonnegative integer such that  $[S, \Phi_{2j}] \notin \mathcal{B}(Q)$ . Clearly  $2j \leq m+2$ . Since  $\Phi_m \in \mathcal{B}(Q^+)$  for even  $m$ , it follows that  $2j \leq m+1$ .

We proceed to show that  $S_{2\nu-1} \subset S$  for  $1 \leq \nu \leq j$ . Since  $[S, \Phi_{2\nu-2}] \in \mathcal{B}(Q)$ , if  $S_{2\nu-1} \not\subset S$  then  $Q$  has a facet  $F$  such that  $F \supset S \cup \Phi_{2\nu-2}$ ,  $F \not\subset \Phi_{2\nu-1}$ , hence  $F \notin \mathcal{C}$  — a contradiction. Since  $[S, \Phi_{2j}] \notin \mathcal{B}(Q)$ ,  $S$  includes a set  $A \in \mathcal{M}(Q/\Phi_{2j})$ .  $S \supset M = \bigcup_{\nu=1}^j S_{2\nu-1} \cup A$ ,  $M$  is of type (a). From this it follows that  $\text{vert } Q \subset \text{vert } Q^+$ .  $x \in \text{vert } Q^+$ , by the definition of  $x$ , hence assertion (1) follows.

*Step 4:* Now prove that if  $x \in S \subset \text{vert } Q^+$  and  $[S] \notin \mathcal{B}(Q^+)$ , then  $S$  includes a set  $M$  of type (a) or (b). Denote  $S^- = S \setminus \{x\}$ . If  $[S^-] \notin \mathcal{B}(Q^+)$ , then  $S^-$  includes a set of type (a), by step 3. If  $[S^-] \in \mathcal{B}(Q^+)$ , then one can show, as in step 3, that  $S$  includes a set of type (b).

*Step 5:* Note that all the sets of type (a) or (b) have  $m+1$  elements (see 4.2(7)).

*Step 6:* Let  $S$  be an element of  $\mathcal{M}(Q^+)$ .  $S$  includes a set  $M$  of type (a) or (b) (by steps 3, 4). But  $[M] \notin \mathcal{B}(Q^+)$  (by steps 1, 2). Hence  $M = S$ , and  $S$  is of type (a) or (b).

*Step 7:* Conversely, assume  $M$  is of type (a) or (b).  $[M] \notin \mathcal{B}(Q^+)$ , hence  $M$  includes a set  $S \in \mathcal{M}(Q^+)$ . By step 6,  $S$  is of type (a) or (b), hence  $|S| = |M| = m+1$ . It follows that  $M = S \in \mathcal{M}(Q^+)$ .

PROOF OF THEOREM 4.6. (1) and (2) follow immediately from Lemma 4.7.

(3) Assume  $0 < 2p \leq m$ . By 4.2(8), in order to prove that  $\Phi_{2p}$  is a u.f. of  $Q^+$ , it suffices to show that  $|M \setminus \Phi_{2p}| \geq m - 2p + 1$  or, equivalently,  $|M \cap \Phi_{2p}| \leq 2p$  for every  $M \in \mathcal{M}(Q^+)$ .

Case I: Assume  $M = S_1 \cup \dots \cup S_{2j-1} \cup A$ ,  $A \in \mathcal{M}(Q/\Phi_{2j})$ . If  $j \leq p$ , then  $|M \setminus \Phi_{2p}| = |A \setminus \Phi_{2p}| \geq m - 2p + 1$ , since  $[A, \Phi_{2p}] \notin \mathcal{B}(Q)$ . If  $j \geq p$ , then  $|M \cap \Phi_{2p}| = |S_1 \cup \dots \cup S_{2p-1}| = 2p$ .

Case II: Assume  $M = S_2 \cup \dots \cup S_{2j} \cup A \cup \{x\}$ ,  $A \in \mathcal{M}(Q/\Phi_{2j+1})$ . If  $j < p$ , then  $|M \setminus \Phi_{2p}| = 1 + |A \setminus \Phi_{2p}| \geq m - 2p + 2$ . If  $j \geq p$ , then  $|M \cap \Phi_{2p}| = 2p$ .

(4) Assume  $1 \leq 2p + 1 \leq m$ . Take  $M = S_1 \cup \dots \cup S_{2p+1} \cup A$  for some  $A \in \mathcal{M}(Q/\Phi_{2p+2})$ . Then  $M \in \mathcal{M}(Q^+)$ ,  $|\Phi_{2p+1} \cap M| = 2p + 2$ , hence  $\Phi_{2p+1}$  is not a u.f. of  $Q^+$ . If  $j$  is odd and  $< m$ , then  $\Phi_{j+1} \in \mathcal{B}(Q^+)$ , hence  $\Phi_j \in \mathcal{B}(Q^+)$ . In fact,  $\Phi_j$  is an  $(m - j - 1)$ -u.f. of  $Q^+$ .

(5) Case I. Assume  $2 \leq 2p \leq m$ ,  $a \in S_{2p}$  and consider  $\Psi = [\Phi_{2p-1}, a, x]$ .

Subcase Ia.  $M = S_1 \cup \dots \cup S_{2j-1} \cup A$ ,  $A \in \mathcal{M}(Q/\Phi_{2j})$ . If  $j \leq p$ , then  $|M \setminus \Psi| \geq |A \setminus \Phi_{2p}| \geq m - 2p + 1$ . If  $j \geq p$ , then  $|M \cap \Psi| = |S_1 \cup \dots \cup S_{2p-1}| = 2p$ .

Subcase Ib.  $M = S_2 \cup \dots \cup S_{2j} \cup A \cup \{x\}$ ,  $A \in \mathcal{M}(Q/\Phi_{2j+1})$ . If  $j < p$ , then  $|M \setminus \Psi| \geq |A \setminus \Phi_{2p}| \geq m - 2p + 1$ . If  $j \geq p$ , then  $|M \cap \Psi| = |S_2 \cup \dots \cup S_{2p-2} \cup \{a, x\}| = 2p$ .

Therefore  $\Psi$  is  $(m - 2p)$ -universal in  $P^+$ .

Case II. Assume  $1 \leq 2p + 1 \leq m$ ,  $a \in S_{2p+1}$ , and consider  $\Psi = [\Phi_{2p}, a, x]$ .

Subcase IIa.  $M = S_1 \cup \dots \cup S_{2j-1} \cup A$ ,  $A \in \mathcal{M}(Q/\Phi_{2j})$ . If  $j \leq p$ , then  $|M \setminus \Psi| \geq |A \setminus \Phi_{2p+1}| \geq m - (2p + 1) + 1$ . If  $j > p$ , then  $|M \cap \Psi| = |S_1 \cup \dots \cup S_{2p-1} \cup \{a\}| = 2p + 1$ .

Subcase IIb.  $M = S_2 \cup \dots \cup S_{2j} \cup A \cup \{x\}$ ,  $A \in \mathcal{M}(Q/\Phi_{2j+1})$ . If  $j \leq p$ , then  $|M \setminus \Psi| \geq |A \setminus \Phi_{2p+1}| \geq m - (2p + 1) + 1$ . If  $j \geq p$ , then  $|M \cap \Psi| = 1 + |S_2 \cup \dots \cup S_{2p}| = 2p + 1$ .

Therefore  $\Psi$  is  $(m - (2p + 1))$ -universal in  $Q^+$ . □

THEOREM 4.8. If  $\Psi \in \mathcal{B}(Q, u)$ ,  $\Psi \cap \Phi_{m-1} = \emptyset$  and  $|\Psi \cap S_m| \leq 1$ , then  $\Psi \in \mathcal{B}(Q^+, u)$ .

PROOF. Assume  $M = \bigcup_{\nu=1}^j S_{2\nu-1} \cup A$ ,  $A \in \mathcal{M}(Q/\Phi_{2j})$  or  $M = \bigcup_{\nu=1}^j S_{2\nu} \cup A \cup \{x\}$ ,  $A \in \mathcal{M}(Q/\Phi_{2j+1})$ . Define  $r = 2j - 1$  in the first case,  $r = 2j$  in the second case. There is a set  $\bar{M} \in \mathcal{M}(Q)$  such that  $A \cup \Phi_{r+1} \supset \bar{M} \supset A$ . If  $r < m$ , then  $|M \cap \Psi| = |A \cap \Psi| \leq |\bar{M} \cap \Psi| \leq m - u$ . If  $r = m$ , then  $|M \cap \Psi| \leq 1$ . If  $u = m$ , then  $\Psi = \emptyset$ , hence  $0 = |M \cap \Psi| \leq m - u = 0$ . If  $u < m$ , then  $|M \cap \Psi| \leq 1 \leq m - u$ . □

**THEOREM 4.9.** *If  $\Psi \in \mathcal{B}(Q, u)$ ,  $\Psi \cap \Phi_m = \emptyset$ ,  $a \in S_p$ ,  $b \in S_{p+1}$ ,  $1 \leq p < m$  and  $[\Psi, a, b] \in \mathcal{B}(Q, u - 1)$ , then  $[\Psi, a, b] \in \mathcal{B}(Q^+, u - 1)$ .*

**PROOF.** We use the same notations as in the proof of Theorem 4.8. If  $r \geq p$ , then  $|M \cap (\Psi \cup \{a, b\})| = |A \cap \Psi| + 1 \leq |\bar{M} \cap \Psi| + 1 \leq m - u + 1$ . If  $r < p$  then  $|M \cap (\Psi \cup \{a, b\})| = |A \cap (\Psi \cup \{a, b\})| \leq |\bar{M} \cap (\Psi \cup \{a, b\})| \leq m - (u - 1)$ .  $\square$

**THEOREM 4.10.** *An edge  $E$  of  $Q^+$  is a u.e. of  $Q^+$  iff either*

- (1)  $E = [a, x]$  and  $a \in S_1$ , or
- (2)  $E$  is a u.e. of  $Q$  and either (a)  $E \cap \Phi_m = \emptyset$  or (b)  $E = [a, b]$  with  $a \in S_p$  and  $b \in S_{p+1}$  for some  $1 \leq p \leq m$ .

**PROOF.** In view of Theorems 4.6(5), 4.8, 4.9 and 3.10 it suffices to prove that if  $E = [a, b]$  is an edge of  $Q$  and either

- ( $\alpha$ )  $a \in S_p$ ,  $b \notin \Phi_{p+1}$  for some  $1 \leq p < m$ , or
- ( $\beta$ )  $E = [S_p]$  for some  $1 \leq p \leq m$ ,

then  $E$  is not universal in  $Q^+$ .

**PROOF OF ( $\alpha$ ).** If  $a \in S_p$ ,  $b \notin \Phi_{p+1}$ ,  $1 \leq p < m$ , then by 4.2(5)  $b$  belongs to a set  $A \in \mathcal{M}(Q/\Phi_{p+1})$ .  $\{a\} \cup A$  is included in a m.f. of  $Q^+$  (of type (a) if  $p$  is odd and of type (b) if  $p$  is even). But  $[a, b] \subset [a, A]$ , hence  $[a, b]$  is not universal in  $Q^+$ .

( $\beta$ ) follows immediately from Lemma 4.7.  $\square$

In the rest of this section we deal with the problem of embedding the complex  $\mathcal{B}(Q, u)$  of  $u$ -universal faces of  $Q$  (see Definition 3.1) in the boundary complex of a neighborly  $2(m - u)$ -polytope.

This material will not be needed in subsequent sections.

If  $|\text{vert } Q| = 2m + 2$ , then the structure of  $Q$  is well-known (see [6, pp. 98, 108]).  $Q$  has two vertex-disjoint missing  $m$ -faces  $\Delta_1, \Delta_2$ , and  $\mathcal{B}(Q, u)$  is the free join of  $\text{skel}_{m-u-1}\Delta_1$  and  $\text{skel}_{m-u-1}\Delta_2$ , i.e. if  $S \subset \text{vert } Q$  then  $[S] \in \mathcal{B}(Q, u)$  iff  $|S \cap \Delta_1| \leq m - u$  and  $|S \cap \Delta_2| \leq m - u$ .

It follows that every  $u$ -universal  $(2m - 2u - 2)$ -face of  $Q$  is included in precisely  $u + 2$   $u$ -universal  $(2m - 2u - 1)$ -faces of  $Q$ , consequently,  $\mathcal{B}(Q, u)$  is not embeddable in the boundary complex of any  $2(m - u)$ -polytope, unless  $u = 0$ .

From now on, assume that  $|\text{vert } Q| = v \geq 2m + 3$ .

For cyclic polytopes the situation is simple: If  $Q = C(v, 2m)$ ,  $v \geq 2m + 3$ , then  $\mathcal{B}(Q, u) \cong \mathcal{B}(C(v, 2(m - u)))$  for  $u = 0, 1, \dots, m - 1$ , and the isomorphism is given by any mapping of  $\text{vert } Q$  onto  $\text{vert } C(v, 2(m - u))$  which preserves the natural cyclic order of the vertices. Moreover, if the vertices of  $Q$  lie on the

moment curve  $x(t) = (t, t^2, \dots, t^{2m})$ , or on the trigonometric moment curve  $x(\theta) = (\cos \theta, \sin \theta, \dots, \cos m\theta, \sin m\theta)$ , then  $C(v, 2(m - u))$  can be taken as the image of  $Q$  under the orthogonal projection of  $R^{2m}$  onto the subspace  $R^{2(m-u)} = \{(x_1, \dots, x_{2(m-u)}, 0, \dots, 0) : x_i \in R\}$ .

In the case  $u = m - 1$ , Theorem 3.10 solves the problem stated above:  $\mathcal{B}(Q, m - 1)$  is isomorphic to a subcomplex of the boundary complex of a convex  $v$ -gon.

Now we shall see that the embeddability of  $\mathcal{B}(Q, u)$  is preserved by the sewing construction.

**THEOREM 4.11.** *Suppose  $Q_u$  is a  $2(m - u)$ -polytope ( $0 < u < m$ ),  $|\text{vert } Q_u| = |\text{vert } Q|$ , and  $\alpha$  is an isomorphism of  $\mathcal{B}(Q, u)$  into  $\mathcal{B}(Q_u)$ . (This implies that  $Q_u$  is neighborly.) Assume also that  $Q^+ = [Q, x]$  is obtained from  $Q$  by sewing at  $x$  through a u.t.  $\mathcal{T}$ .*

*Then there is a polytope  $Q_u^+ = [Q_u, x_u]$ , obtained from  $Q_u$  by sewing at  $x_u$  through an appropriate u.t.  $\mathcal{T}_u$ , and an isomorphism  $\alpha^+$  of  $\mathcal{B}(Q^+, u)$  into  $\mathcal{B}(Q_u^+)$ . Moreover,  $\alpha^+(q) = \alpha(q)$  for all  $q \in \text{vert } Q$ .*

**PROOF.** Define  $\mathcal{T}_u = \{\alpha\Phi_j\}_{j=1}^{m-u}$ . It is easily checked that  $\mathcal{T}_u$  is u.t. in  $Q_u$ . Choose  $x_u$  to be a point that lies exactly beyond  $\mathcal{C}(Q_u, \mathcal{T}_u)$ , and let  $Q_u^+ = [Q_u, x_u]$ .

Define  $\alpha^+ : \text{vert } Q^+ \rightarrow \text{vert } Q_u^+$  by  $\alpha^+(x) = x_u, \alpha^+(q)$  for  $q \in \text{vert } Q$ . In order to show that  $\alpha^+$  extends to an isomorphism of  $\mathcal{B}(Q^+, u)$  into  $\mathcal{B}(Q_u^+)$ , we must prove that if  $S \subset \text{vert } Q^+$  and  $[\alpha^+S] \notin \mathcal{B}(Q_u^+)$  then  $[S] \notin \mathcal{B}(Q^+, u)$ . It suffices to prove this for  $S \subset \text{vert } Q^+$  such that  $[\alpha^+S]$  is a m.f. of  $Q_u^+$ .

Assume  $[\alpha^+S]$  is a m.f. of  $Q_u^+$ .

*Case I:*  $\alpha^+S = \alpha(S_1 \cup S_3 \cup \dots \cup S_{2j-1}) \cup \alpha A$  where  $0 \leq j < \frac{1}{2}(m - u + 1)$  and  $\alpha A \in \mathcal{M}(Q_u / \alpha\Phi_{2j})$ , or  $j = \frac{1}{2}(m - u + 1)$  and  $A = \emptyset$  (see Lemma 4.7). It follows that  $S = S_1 \cup S_3 \cup \dots \cup S_{2j-1} \cup A$  and  $[\alpha A, \alpha\Phi_{2j}] \notin \mathcal{B}(Q_u)$ . Therefore  $[A, \Phi_{2j}] \notin \mathcal{B}(Q, u)$ , hence there is a set  $B \subset \text{vert } Q, |B| \leq u$  and  $[A, B, \Phi_{2j}] \notin \mathcal{B}(Q)$ .

If  $F$  is a facet of  $Q$  and  $F \supset S \cup B$ , then  $F \in (\mathcal{F}_1 \setminus \mathcal{F}_2) \cup \dots \cup (\mathcal{F}_{2j-3} \setminus \mathcal{F}_{2j-2}) \cup \mathcal{F}_{2j-1}$  and  $F \notin \mathcal{F}_{2j}$ . Hence  $[S, B] \notin \mathcal{B}(Q^+)$ , and consequently  $[S] \notin \mathcal{B}(Q^+, u)$ .

*Case II:*  $\alpha^+S = \alpha(S_2 \cup S_4 \cup \dots \cup S_{2j}) \cup \alpha A \cup \{x_u\}$  where  $0 \leq j < \frac{1}{2}(m - u)$  and  $\alpha A \in \mathcal{M}(Q_u / \alpha\Phi_{2j+1})$ , or  $j = \frac{1}{2}(m - u)$  and  $A = \emptyset$ .

As above, we find that  $S = S_2 \cup S_4 \cup \dots \cup A \cup \{x\}$  and  $[A, \Phi_{2j+1}] \notin \mathcal{B}(Q, u)$ . There is a set  $B \subset \text{vert } Q$  such that  $|B| \leq u$  and  $[A, B, \Phi_{2j+1}] \notin \mathcal{B}(Q)$ .

If  $F$  is a facet of  $Q$  and  $F \supset B \cup (S \setminus \{x\})$ , then  $F \in (\mathcal{F}_0 \setminus \mathcal{F}_1) \cup \dots \cup (\mathcal{F}_{2j} \setminus \mathcal{F}_{2j+1})$ , hence  $[S, B] \notin \mathcal{B}(Q^+)$ , and therefore  $[S] \notin \mathcal{B}(Q^+, u)$ .  $\square$

Theorem 4.11 states that, in an appropriate sense,  $(Q^+)_u = (Q_u)^+$ .

It is probably not true that for every neighborly  $2m$ -polytope  $Q$  (with  $|\text{vert } Q| \geq 2m + 3$ ) and for all  $1 \leq u \leq m - 2$ , the complex  $\mathcal{B}(Q, u)$  can be embedded in the boundary complex of a neighborly  $2(m - u)$ -polytope  $Q_u$ , though we have no counterexample.

### 5. Reconstruction theorems

If  $Q, \mathcal{T}, \mathcal{C}, x$  and  $Q^+$  are as in Convention 4.5, then we say that  $Q^+$  is obtained from  $Q$  at  $x$  by sewing through the tower  $\mathcal{T}$ . We claim that the tower  $\mathcal{T}$  is determined by  $Q^+$  and  $x$  in the following sense:

If  $Q^+$  is obtained from  $Q$  at  $x$  by sewing through any tower  $\mathcal{T}'$ , then  $\mathcal{T}' = \mathcal{T}$ .

In order to prove this it suffices to show that if  $\mathcal{C}(Q, \mathcal{T}) = \mathcal{C}(Q, \mathcal{T}')$  then  $\mathcal{T} = \mathcal{T}'$ , because  $Q^+$  and its vertex  $x$  determine  $\mathcal{C}$  (see Theorem 2.10).

LEMMA 5.1. *Let  $\Phi$  be a u.f. of  $Q$  with  $2j$  vertices,  $\Psi$  a u.f. of  $Q$  with  $2j + 2$  vertices and  $\Psi \supset \Phi$ . Then  $\Phi = \cap (\mathcal{F}_\Phi \setminus \mathcal{F}_\Psi)$  (see Definition 4.3).*

PROOF. Obviously  $\Phi \subset \cap (\mathcal{F}_\Phi \setminus \mathcal{F}_\Psi) \subset Q$ . Suppose  $y \in \text{vert } Q \setminus \Phi$ . We have to show that  $Q$  has a facet  $F$  such that  $F \supset \Phi, F \not\supset \Psi$  and  $y \notin F$ . If  $y \in \Psi \setminus \Phi$  then every facet of  $Q$  which includes  $\Phi$  and does not contain  $y$  is in  $\mathcal{F}_\Phi \setminus \mathcal{F}_\Psi$ .

Suppose  $y \notin \Psi$ . By 4.2(5) there is a set  $A \subset \text{vert } Q \setminus (\Psi \cup \{y\})$  such that  $[A, \Psi] \notin \mathcal{B}(Q)$  and  $|A| = m - j$  (Proposition 4.2(7)). Since  $y \notin [\Phi, A]$ , it follows that  $Q$  has a facet  $F$  such that  $\Phi \subset F, A \subset F, y \notin F$  and necessarily  $\Psi \not\subset F$ .  $\square$

THEOREM 5.2. *For  $i = 1, 2$ , let  $\mathcal{T}_i$  be a universal tower in  $Q$ . If  $\mathcal{C}(Q, \mathcal{T}_1) = \mathcal{C}(Q, \mathcal{T}_2)$  then  $\mathcal{T}_1 = \mathcal{T}_2$ .*

PROOF. Let  $\mathcal{T} = \{\Phi_i\}_{i=1}^m$  be a u.t. in  $Q$ . We shall prove that  $\mathcal{C} = \mathcal{C}(Q, \mathcal{T})$  and  $\mathcal{F}(Q)$  determine  $\Phi_i$  ( $i = 1, \dots, m$ ).

Denote by  $\mathcal{F}_0$  the set of all facets of  $Q$ . We proceed to define by induction a sequence  $\mathcal{C}_1, \mathcal{C}_2, \dots$  of subsets of  $\mathcal{F}_0$ , as follows:

$$\mathcal{C}_1 = \mathcal{C}, \text{ and } \mathcal{C}_j = \{F \in \mathcal{F}_0 \setminus \mathcal{C}_{j-1} : \cap \mathcal{C}_{j-1} \subset F\} \text{ for } j > 1.$$

We claim that  $\Phi_j = \cap \mathcal{C}_j$  for  $j = 1, 2, \dots, m$ . Assume  $1 \leq j \leq m$  and  $\Phi_i = \cap \mathcal{C}_i$  for all  $i, 1 \leq i < j$ . It is easy to prove, by induction on  $i$ , that  $\mathcal{C}_i = \bigcup_{\nu \geq 0} (\mathcal{F}_{i+2\nu} \setminus \mathcal{F}_{i+2\nu+1})$  for  $1 \leq i \leq j$ . In particular  $\mathcal{F}_j \setminus \mathcal{F}_{j+1} \subset \mathcal{C}_j \subset \mathcal{F}_j$ . For  $j < m$ , Lemma 5.1 yields  $\Phi_j = \cap (\mathcal{F}_j \setminus \mathcal{F}_{j+1}) \supset \cap \mathcal{C}_j \supset \cap \mathcal{F}_j = \Phi_j$ , hence  $\Phi_j = \cap \mathcal{C}_j$ . If



$j = m$ , then  $\mathcal{C}_m = \mathcal{F}_m$ , hence  $\Phi_m = \cap \mathcal{C}_m$ . For  $j > m$ , it can be easily checked that  $\mathcal{C}_j = \emptyset$ . □

**THEOREM 5.3.** *Let  $Q, Q^+$  be neighborly  $2m$ -polytopes,  $\text{vert } Q^+ = \text{vert } Q \cup \{x\}$ ,  $x \notin Q$ . Suppose  $a, b \in \text{vert } Q$ ,  $a \neq b$ . If  $[a, x]$  and  $[b, x]$  are universal edges of  $Q^+$ , then  $[a, b]$  is a universal edge of  $Q$ .*

Theorem 5.3 is a special case of Corollary 3.9. It can also be deduced directly from the following theorem:

**THEOREM 5.4.** *Let  $Q, Q^+$  and  $x$  be as in Theorem 5.3. If  $a \in \text{vert } Q$ , then  $[a, x]$  is a u.e. of  $Q^+$  iff all the facets of  $Q$  that  $x$  covers contain  $a$ .*

**PROOF.** Suppose  $[a, x]$  is a u.e. of  $Q^+$ . For  $0 < \lambda < 1$  let  $z = z(\lambda) = (1 - \lambda)a + \lambda x$ . It is easy to see that  $[Q, z]$  is an  $m$ -neighborly polytope. If  $Q$  had a facet  $F$  such that  $a \notin F$  and  $x$  lies beyond  $F$ , then  $\lambda$  could be chosen such that  $z \in \text{aff } F$ , contradicting the simpliciality of  $[Q, z]$ .

Now suppose that all the facets of  $Q$  that  $x$  covers contain  $a$ . Assume  $A \subset \text{vert } Q$ ,  $|A| = m - 1$ . We have to show that  $[a, x, A] \in \mathcal{F}(Q^+)$ . Since  $[a, A] \in \mathcal{F}(Q^+)$ , it suffices to prove that  $Q$  has a facet  $F$  such that  $x$  lies beyond  $F$  and  $\{a\} \cup A \subset F$ . But  $[A, x] \in \mathcal{F}(Q^+)$ , hence  $Q$  has a facet  $F$  that includes  $A$  and is covered by  $x$ . Necessarily,  $a \in F$ . □

**PROOF OF THEOREM 5.3.** Suppose  $[a, x], [b, x]$  are u.e.s of  $Q^+$ . If  $A \subset \text{vert } Q$ ,  $|A| = m - 1$ , then  $[A, x] \in \mathcal{F}(Q^+)$ , hence  $Q$  has a facet  $F$  such that  $F \supset A$  and  $x$  lies beyond  $F$ . By Theorem 5.4,  $F \supset \{a, b\}$ , and therefore  $[a, b, A] \in \mathcal{F}(Q)$ . □

The same technique leads to a proof of the next theorem.

**THEOREM 5.5.** *Let  $\mathcal{C}$  and  $\mathcal{D}$  be sets of facets of  $Q$ . Let  $x, y$  be points in  $R^{2m} \setminus Q$ , such that  $x$  lies exactly beyond  $\mathcal{C}$  and  $y$  lies exactly beyond  $\mathcal{D}$ . Assume both  $[Q, x]$  and  $[Q, y]$  are neighborly  $2m$ -polytopes with  $|\text{vert } Q| + 1$  vertices. If  $\mathcal{C} \subset \mathcal{D}$  then  $\mathcal{C} = \mathcal{D}$ .*

**PROOF.** For  $0 < \lambda < 1$ , define  $z = z(\lambda) = (1 - \lambda)x + y$ . It is easy to see that  $[Q, z]$  is a neighborly polytope ( $z$  covers enough, but not too many facets of  $Q$ ) and the simpliciality of  $[Q, z]$  for every choice of  $\lambda$ ,  $0 < \lambda \leq 1$ , implies that  $\mathcal{D} \setminus \mathcal{C} = \emptyset$ . □

As a matter of fact,  $|\mathcal{C}|$  is a function of  $v = |\text{vert } Q|$  and  $m$  only:

$$|\mathcal{C}| = \binom{v - m - 1}{m - 1}.$$

The next theorem is a converse to Theorem 4.6.

**THEOREM 5.6.** *Let  $Q, Q^+$  be neighborly  $2m$ -polytopes. Assume  $\text{vert } Q^+ = \text{vert } Q \cup \{x\}$ ,  $x \notin Q$  and let  $\mathcal{T} = \{\Phi_j\}_{j=1}^m$  be a tower in  $Q$  with  $|\text{vert } \Phi_j| = 2j$  for  $j = 1, 2, \dots, m$ . If*

- (1)  $\Phi_j$  is a u.f. of  $Q^+$  for every even  $j$ ,  $1 \leq j \leq m$ , and
- (2)  $[\Phi_j, p, x]$  is a u.f. of  $Q^+$  for every even  $j$ ,  $0 \leq j < m$  and  $p \in \text{vert } \Phi_{j+1} \setminus \Phi_j$  (here  $\Phi_0 = \emptyset$ ),

*then  $\mathcal{T}$  is a universal tower in  $Q$ , and  $Q^+$  is obtained from  $Q$  at  $x$  by sewing through  $\mathcal{T}$ .*

**REMARK.** If  $P$  is a subpolytope of  $P^+$ , and  $\Phi$  is a face of both  $P$  and  $P^+$ , then there is a flat  $H$  such that the correspondence  $\Psi \rightarrow \Psi \cap H$  is an isomorphism of the upper segment  $[\Phi, P^+]$  of  $\mathcal{F}(P^+)$  onto  $\mathcal{F}(P^+ \cap H)$ , and the same correspondence is an isomorphism of the upper segment  $[\Phi, P]$  of  $\mathcal{F}(P)$  onto  $\mathcal{F}(P \cap H)$  (see [8, pp. 72–73]).

**PROOF.** Denote by  $\mathcal{D}$  the set of facets of  $Q$  covered by  $x$ . We have to show that  $\mathcal{T}$  is a u.t. in  $Q$  and that  $\mathcal{D} = \mathcal{C}(Q, \mathcal{T})$  (see Definition 4.3).

Assume  $j$  is even,  $0 \leq j < m$ . Consider the polytopes  $Q/\Phi_j$  and  $Q^+/\Phi_j$ . They are neighborly  $2(m - j)$ -polytopes, and  $\text{vert } Q^+/\Phi_j = \text{vert } Q/\Phi_j \cup \{[\Phi_j, x]/\Phi_j\}$  by the remark above. By Corollary 5.3,  $\Phi_{j+1}/\Phi_j$  is a u.f. of  $Q/\Phi_j$ . Therefore  $[\Phi_{j+1}, A]/\Phi_j$  is a face of  $Q/\Phi_j$  for every  $A \subset \text{vert } Q \setminus \Phi_j$ ,  $|A| \leq m - j - 1$ . Hence  $\Phi_{j+1}$  is a u.f. of  $Q$ . It follows that  $\mathcal{T}$  is a u.t. in  $Q$ .

Denote by  $x'$  the vertex  $[\Phi_j, x]/\Phi_j$  of  $Q^+/\Phi_j$ . From condition (2) and Theorem 5.4 it follows that if  $x'$  lies beyond the facet  $F' = F/\Phi_j$  of  $Q/\Phi_j$  then  $F \supset \Phi_{j+1}$ . (Note that  $x'$  lies beyond  $F'$  iff  $x$  lies beyond  $F$ .) It follows that  $\mathcal{D} \cap \mathcal{F}_j \subset \mathcal{F}_{j+1}$  for even  $j$ ,  $0 \leq j < m$ . The same conclusion follows from condition (1) for  $j = m$ , if  $m$  is even ( $\mathcal{F}_{m+1} = \emptyset$ ). Therefore  $\mathcal{D} \cap (\mathcal{F}_j \setminus \mathcal{F}_{j+1}) = \emptyset$  for even  $j$ ,  $0 \leq j \leq m$ , hence  $\mathcal{D} \cap \cup\{(\mathcal{F}_j \setminus \mathcal{F}_{j+1}) : j \text{ even}\} = \emptyset$ , i.e.,  $\mathcal{D} \subset \mathcal{C}(Q, \mathcal{T})$ . By Theorem 4.6 and Theorem 5.5 we conclude that  $\mathcal{D} = \mathcal{C}(Q, \mathcal{T})$ . □

Theorem 5.6 says that we can tell from partial information about the structure of  $\mathcal{F}(Q)$  whether  $Q$  is obtained by sewing at a given vertex  $x$  through a given tower  $\mathcal{T}$ . The next theorem shows how this implies “commutativity” of our sewing procedure.

**DEFINITION 5.7.** If  $\mathcal{T}$  is a u.t. in  $Q$ , let  $Q(\mathcal{T})$  denote the class of all polytopes that are obtained from  $Q$  by sewing through  $\mathcal{T}$ .

**THEOREM 5.8.** *For  $1 \leq i \leq p$ , let  $\mathcal{T}_i$  be a u.t. in  $Q$ . Assume the sets  $\cup \mathcal{T}_i$ ,  $1 \leq i \leq p$ , are pairwise disjoint (i.e.,  $A \in \mathcal{T}_i$ ,  $B \in \mathcal{T}_j$  and  $i \neq j$  imply*

$A \cap B = \emptyset$ ). Then there are points  $x_i, 1 \leq i \leq p$ , such that  $[Q, \{x_i : i \in \Gamma\}, x_j] \in [Q, \{x_i : i \in \Gamma\}](\mathcal{T}_j)$  for every subset  $\Gamma$  of  $\{1, \dots, p\}$  and for every  $j$  in  $\{1, \dots, p\} \setminus \Gamma$ .

PROOF. By Lemma 4.4 and Theorem 4.8 it follows easily, by induction on  $k, 1 \leq k \leq p$ , that there are points  $x_1, \dots, x_p$  such that the sequence  $(x_1, \mathcal{T}_1), \dots, (x_p, \mathcal{T}_p)$  satisfies

$$(*) \quad [Q, x_1, \dots, x_k] \in [Q, x_1, \dots, x_{k-1}](\mathcal{T}_k) \quad \text{for } 1 \leq k \leq p.$$

Theorem 5.8 follows from the next lemma. □

LEMMA 5.9. Under the assumptions of Theorem 5.8, if the sequence  $\{(x_j, \mathcal{T}_j)\}_{j=1}^p$  satisfies  $(*)$ , then the sequence  $\{(x_{\pi(j)}, \mathcal{T}_{\pi(j)})\}_{j=1}^p$  satisfies  $(*)$  for any permutation  $\pi$  of  $\{1, \dots, p\}$ .

PROOF. For  $p = 1$  there is nothing to prove.

Assume  $p = 2$ . Consider the faces of  $[Q, x_1, x_2]$  whose universality is asserted in Theorem 4.6, parts (3) and (5), with  $x = x_2$  and  $\mathcal{T} = \mathcal{T}_2$ . Obviously, each such face is a u.f. of  $[Q, x_2]$ , and by Theorem 5.6,  $[Q, x_2] \in Q(\mathcal{T}_2)$ . Now consider the faces of  $[Q, x_1]$  whose universality is asserted in Theorem 4.6, parts (3) and (5), with  $x = x_1$  and  $\mathcal{T} = \mathcal{T}_1$ . By Theorem 4.8, these faces are u.f.s also in  $[Q, x_1, x_2]$ , and by Theorem 5.6,  $[Q, x_2, x_1] \in [Q, x_2](\mathcal{T}_1)$ . Therefore the sequence  $(x_2, \mathcal{T}_2), (x_1, \mathcal{T}_1)$  satisfies  $(*)$ .

Now assume  $p > 2$ . We say that  $\pi$  is an admissible permutation of  $\{1, \dots, p\}$  if  $\{(x_{\pi(j)}, \mathcal{T}_{\pi(j)})\}_{j=1}^p$  satisfies  $(*)$  whenever  $\{(x_j, \mathcal{T}_j)\}_{j=1}^p$  satisfies  $(*)$ . Since the set of admissible permutations is closed under multiplication, it suffices to show that the transpositions  $(i, i + 1), 1 \leq i < p$ , are admissible.

Let  $x_1, \dots, x_p$  be points such that  $(*)$  holds for  $(x_1, \mathcal{T}_1), \dots, (x_p, \mathcal{T}_p)$ . Assume  $1 \leq i < p$ . We shall show that the sequence  $(x_1, \mathcal{T}_1), \dots, (x_{i+1}, \mathcal{T}_{i+1}), (x_i, \mathcal{T}_i), \dots, (x_p, \mathcal{T}_p)$  also satisfies  $(*)$ . The only parts of  $(*)$  that are not self-evident for this sequence are  $[Q', x_{i+1}] \in Q'(\mathcal{T}_{i+1})$  and  $[Q', x_{i+1}, x_i] \in [Q', x_{i+1}](\mathcal{T}_i)$ , where  $Q' = [Q, x_1, \dots, x_{i-1}]$ . But these assertions follow from the case  $p = 2$ , applied to  $Q'$  and to the sequence  $(x_i, \mathcal{T}_i), (x_{i+1}, \mathcal{T}_{i+1})$ . □

Note that the proof of Theorem 5.8 given here holds even if the assumption that the sets  $\bigcup \mathcal{T}_i$  are pairwise disjoint is replaced by the following slightly weaker condition: Assume  $\mathcal{T}_i = \{\Phi_{i,1}, \dots, \Phi_{i,m}\}$  for  $1 \leq i \leq p$ , then  $\Phi_{i,m-1} \cap \Phi_{j,m} = \emptyset$  and  $|\Phi_{i,m} \cap \Phi_{j,m}| \leq 1$  for  $i \neq j, 1 \leq i, j \leq p$ .

It would be interesting to know what is the freedom in choosing the points  $x_1, \dots, x_p$ , in case we do assume that the towers  $\mathcal{T}_1, \dots, \mathcal{T}_p$  have pairwise disjoint unions.

The following might be true:

If  $[Q, x_i] \in Q(\mathcal{T}_i)$  for  $1 \leq i \leq p$ , then  $[Q, \{x_i : i \in \Gamma\}, x_j] \in [Q, \{x_i : i \in \Gamma\}](\mathcal{T}_j)$  for every  $\Gamma \subset \{1, 2, \dots, p\}$  and  $j \in \{1, 2, \dots, p\} \setminus \Gamma$ .

**6. Lower bounds**

Repeated application of the sewing construction described in section 4 yields lower bounds for the number  $g(v, 2m)$  of combinatorial types of neighborly  $2m$ -polytopes with  $v$  vertices.

We start with a cyclic  $2m$ -polytope  $C(v, 2m)$  with  $v$  vertices,  $v \geq 2m + 3$ , and “sew” it repeatedly.

The main results are as follows:

**THEOREM 6.1.**  $g(2m + 4, 2m) > \frac{(2m + 2)!}{3 \cdot 2^{m+3}(m + 2)!}$ .

(The right-hand side is asymptotic to  $(\sqrt{2}/6)(2m/e)^m$  as  $m \rightarrow \infty$ .)

**THEOREM 6.2.**  $g((2m + 1)p + 2, 2m) \geq \frac{1}{2}(pm - p)!$  for  $p = 2, 3, \dots$ .

**PROOF OF THEOREM 6.1.** Let  $K$  be  $C(2m + 3, 2m)$ . Assume that  $\text{vert } K = \{a_1, a_2, \dots, a_{2m+3}\}$ , and that  $a_1, a_2, \dots, a_{2m+3}, a_1$  is the circuit of universal edges of  $K$ .  $\text{Aut } K$ , the group of combinatorial automorphisms of  $K$ , is precisely the dihedral group of order  $2(2m + 3)$  consisting of rotations and reflections of the circuit  $a_1, a_2, \dots, a_{2m+3}, a_1$ .

Consider pairs  $(Q, x)$  where  $Q$  is a neighborly  $2m$ -polytope with  $2m + 4$  vertices and  $x$  is a distinguished vertex of  $Q$ . Two such pairs  $(Q, x), (Q', x')$  are considered isomorphic if there is a combinatorial equivalence  $\varphi : \text{vert } Q \rightarrow \text{vert } Q'$  with  $\varphi(x) = x'$ . The number of isomorphism types of such pairs is clearly at most  $(2m + 4)g(2m + 4, 2m)$ .

Now let us count the number of isomorphism types of pairs  $(K^+, x)$ , where  $K^+$  is obtained from  $K$  at  $x$  by sewing through a u.t.  $\mathcal{T}$ . Consider two such pairs  $(K_1^+, x_1)$ , with  $K_i^+ \in K(\mathcal{T}_i), i = 1, 2$ . If these pairs are isomorphic by a mapping  $\varphi : \text{vert } K_1^+ \rightarrow \text{vert } K_2^+$  such that  $\varphi(x_1) = x_2$ , then  $\psi = \varphi \upharpoonright K$  is an automorphism of  $K$ , and  $\psi$  maps  $\mathcal{C}(K, \mathcal{T}_1)$  onto  $\mathcal{C}(K, \mathcal{T}_2)$ . It follows from Theorem 5.2 that  $\psi(\mathcal{T}_1) = \mathcal{T}_2$ .

The converse is obvious: If  $\mathcal{T}_1$  is mapped onto  $\mathcal{T}_2$  by an automorphism of  $K$ , then the pairs  $(K_1^+, x_1), (K_2^+, x_2)$  are isomorphic.

Thus, the number of isomorphism types of pairs  $(K^+, x)$  considered above is precisely the number of equivalence classes of u.t.s of  $K$  under  $\text{Aut } K$ .

Every u.t. in  $K$  can be transformed by a suitable rotation to a tower with  $\Phi_1 = [a_1, a_2]$ . With  $\Phi_1$  fixed,  $\Phi_2$  can be chosen in  $2m + 1$  ways, then  $\Phi_3$  in  $2m - 1$  ways and so on. (Note that  $K/\Phi_i$  is a cyclic polytope of type  $C(2(m - i) + 3, 2(m - i))$ , for  $0 \leq i < m$ .) Therefore the number of u.t.s in  $K$  with  $\Phi_1 = [a_1, a_2]$  is  $(2m + 1)(2m - 1) \cdots 7 \cdot 5$ . There is only one automorphism  $\psi$  of  $K$ , except the identity, that maps the edge  $\Phi_1 = [a_1, a_2]$  onto itself ( $\psi(a_i) = a_{2m+6-i}$  for  $3 \leq i \leq 2m + 3$ ).

Only one u.t. is fixed by  $\psi$ . It follows that the number of equivalence classes of u.t.s in  $K$  under  $\text{Aut } K$  is precisely  $\frac{1}{2}(1 + (2m + 1)(2m - 1) \cdots 5)$ .

We conclude that  $(2m + 4)g(2m + 4, 2m) > \frac{1}{6} \prod_{i=1}^m (2i + 1)$ . □

If we repeat the reasoning in the proof of Theorem 6.1 with  $K = C(2m + 3, 2m)$  replaced by  $K = C(v - 1, 2m)$ , we conclude that  $g(v, 2m) \rightarrow \infty$  as  $v \rightarrow \infty$ . But the following construction yields a better lower bound.

**PROOF OF THEOREM 6.2.** Let  $K$  be a cyclic polytope  $C(v, 2m)$  with  $v = 2mp + 2$  vertices,  $p \geq 2$ . Assume that  $\text{vert } K = \{a_1, a_2, \dots, a_v\}$  and  $a_1, a_2, \dots, a_v, a_1$  is the circuit of u.e.s of  $K$ . We say that  $a_i$  and  $a_j$  are successive vertices if  $j = i + 1$ . A family  $\sigma = \{\sigma(i, j), 1 \leq i \leq p, 1 \leq j \leq m\}$  is called a partition of  $\text{vert } K$  if:

- (1)  $\sigma(i, j)$  is a set of two successive vertices of  $K$ ,  $1 \leq i \leq p, 1 \leq j \leq m$ .
- (2)  $\sigma(i, 1) = \{a_{2i}, a_{2i+1}\}, 1 \leq i \leq p$ .
- (3)  $\sigma(i, j) \subset \{a_{2p+3}, a_{2p+4}, \dots, a_v\}, 1 \leq i \leq p, 1 < j \leq m$ .
- (4) Every vertex of  $K$ , except  $a_1$  and  $a_{2p+2}$ , is contained in some  $\sigma(i, j)$ .

From (1)–(4) it follows that the sets  $\sigma(i, j)$  are pairwise disjoint.

For every partition  $\sigma$  define:  $\mathcal{T}_i = \mathcal{T}_i(\sigma) = \{[\sigma(i, 1), \dots, \sigma(i, l)]\}_{l=1}^m, 1 \leq i \leq p$ . For  $1 \leq i \leq p, \mathcal{T}_i$  is a u.t. in  $K$ , and the sets  $\bigcup \mathcal{T}_i$ , for  $1 \leq i \leq p$ , are pairwise disjoint. By Theorem 5.8 there are points  $x_i = x_i(\sigma), 1 \leq i \leq p$ , such that  $[K, \{x_j : j \in \Gamma\}, x_i] \in [K, \{x_j : j \in \Gamma\}](\mathcal{T}_i)$  for  $\Gamma \subset \{1, \dots, p\} \setminus \{i\}$ .

Define  $K^\sigma = [K, x_1, \dots, x_p]$ .  $K^\sigma$  is a neighborly  $2m$ -polytope with  $v + p = (2m + 1)p + 2$  vertices. We shall show that the number of distinct combinatorial types of polytopes  $K^\sigma$  is at least one half the number  $(pm - p)!$  of partitions.

Define an equivalence relation on the set of partitions as follows:  $\sigma \sim \tau$  if  $K^\sigma \cong K^\tau$ .

In order to complete the proof, it suffices to show that an equivalence class of a partition contains at most two elements.

Assume  $\sigma \sim \tau$ , and let  $f : K^\sigma \rightarrow K^\tau$  be a combinatorial equivalence.

If  $E$  is a u.e. of  $K^\sigma$  then either  $E$  is a u.e. of  $K$  or  $x_i = x_i(\sigma) \in \text{vert } E$  for some  $1 \leq i \leq p$ .  $K^\sigma$  is sewed at  $x_i$  through  $\mathcal{T}_i$  (Theorem 5.8), hence  $[x_i, a_{2i}]$  and