labile factors important to the nutritional value of the solubles.

As solubles are used as supplements in animal feeds, consideration should be given to the relative nutritional value of solubles prepared by acid and enzyme treatment of stickwater. When the two types of solubles were used to supplement both natural and synthetic rations for the growth of chicks, no significant differences in the nutritive value of the products could be detected (11).

## Acknowledgment

The authors wish to thank the Rohm and Haas Co., Philadelphia, Pa., for samples of Rhozyme B-6; Merck & Co., Ltd., Montreal, for a supply of ficin; Takamine Laboratory, Inc., Clifton, N. J., for samples of bromelin and protease; Nutritional Biochemicals Corp., Cleveland, Ohio, for supplying papain; and Wallerstein Laboratories, New York, N. Y., for samples of colla-

The authors also acknowledge the assistance of P. J. G. Carrothers and P. B. Crean, who designed and constructed the pilot plant; the late N. A. W. LePage, who assisted in the operation of the pilot plant; and W. A. B. Thomson, who carried out some of the earlier phases of this investigation. C. J. Tai and L. Lukey provided technical assist-

ance. The cooperation of the management and staff of British Columbia Packers, Ltd., in making samples available has been greatly appreciated.

## References

- (1) Block, R. J., Durrum, E. L., Zweig, G., "A Manual of Paper Chromatography and Paper Electrophoresis, p. 67, Academic Press, New York,
- (2) Burley, R. W., Solomons, C. C.,
- S. African Ind. Chemist 11, 154 (1957).

  (3) Carrothers, P. J. G., "Design of a Pilot Plant for Processing Stickwater," unpublished rept., Fisheries Research Board of Canada, Technological Station, Vancouver, 1957
- (4) Eastoe, J. E., Biochem. J. 65, 363 (1957).
- (5) Gunther, J. K., Sair, L., U. S. Patent **2,525,294** (Oct. 10, 1950).
- (6) Hart, J. L., Tester, A. L., Trans. Am. Fisheries Soc. 64, 307 (1934).
  (7) Idler, D. R., MacLeod, R. A.,
- Thomson, W. A. B., Appl. Microbiol. 3, 205 (1955).
- (8) Jackson, D. S., Biochem. J. 65, 277 (1957).
- (9) Kuntzel, A., Koepff, H., J. Intern. Soc. Leather Trades' Chemists 24, 92
- (10) Lassen, H. S., U. S. Patent 2,372,-677 (April 3, 1945).
- (11) March, B. E., Biely, J., McBride, J., MacLeod, R. A., Idler, D. R., Fisheries Research Board Can. Progr.

- Repts. Pacific Coast States No. 111, 23 (1958).
- (12) Martin, C. J., Axelrod, A. E., Proc. Soc. Exptl. Biol. Med. 83, 461 (1953).
- (13) Meldinger S.S.F. 1, 5 (Jan. 20, 1950).
  (14) Neuman, R. E., Logan, M. A., J. Biol. Chem. 186, 549 (1950).
- (15) Rigg, G., Carpenter, J. L., J. Ind. Eng. Chem. 4, 901 (1912).
- (16) Sahyun, M., "Outline of the Amino Acids and Proteins," 2nd ed., p. 926, Reinhold, New York, 1948.
- (17) Thomson, W. A. B., Jonas, R. E. E., MacLeod, R. A., Idler, D. R., Fisheries Research Board Can., Circ. 21 (April 1958).

Received for review February 21, 1959. Accepted June 15, 1959.

## Fungicide Distribution and Disinfection Efficiency in Seed Treatment—Correction

On page 327 [J. AGR. FOOD CHEM. 7, 326 (1959)], the caption for Figure 2

Results from seed treatment tests with  $\log \Phi$  as a function of  $\log \operatorname{dose}(O, \Phi)$ 

**1**, **(**) compared to probit analysis  $(\bullet)$ 

In section III of the subcaption of Figure 2, the last symbol (•) should be O. Olle Lindström