

B₂, G₁, G₂, and B₁A under the conditions of the experiment. Differences in the behavior of strains of *C. pyrenoidosa* and *C. vulgaris* to the action of various herbicides can be and have been attributed to strain differences (Vance and Smith, 1969; Thomas *et al.*, 1971) and to differences in growth conditions. By changing from autotrophic conditions to heterotrophic conditions by the addition of a reduced carbon source such as glucose, it has been shown that the inhibitory effects of some herbicides which affect the photosynthetic process on strains of *C. pyrenoidosa* and *C. vulgaris* can be reversed (Geoghagen, 1957; Ashton *et al.*, 1966; Sikka and Pramer, 1968). In order to reduce the time of the assay and in order to avoid special provisions for a carbon dioxide source, glucose has been used as a carbon source under our conditions.

The results presented here emphasize the importance of strain differences when *Chlorella* is used as a test organism.

LITERATURE CITED

Ashton, F. N., Bisalputra, T., Risley, E. B., *Amer. J. Bot.* **53**, 217 (1966).

Geoghagen, M. J., *New Phytol.* **56**, 71 (1957).
Ikawa, M., Ma, D. S., Meeker, G. B., Davis, R. P., *J. Agr. Food Chem.* **17**, 425 (1969).
Sikka, H. C., Pramer, D., *Weed Sci.* **16**, 296 (1968).
Starr, R. C., *Amer. J. Bot.* **51**, 1013 (1964).
Thomas, V. M., Jr., Sullivan, J. D., Jr., Ikawa, M., Univ. of N.H., Durham, N.H., unpublished results (1971).
Vance, B. D., Smith, D. L., *Texas J. Sci.* **20**, 329 (1969).

James D. Sullivan, Jr.
Miyoshi Ikawa*

Department of Biochemistry
University of New Hampshire
Durham, New Hampshire 03824

Received for review January 12, 1972. Accepted March 3, 1972. Published as Scientific Contribution No. 599 of the New Hampshire Agricultural Experiment Station. This work was supported in part by Grant 616-15-4 from the Cooperative State Research Service, U.S. Department of Agriculture, and Grant EC-00294 from the Environmental Control Administration, U.S. Public Health Service.

Correction

DETERMINATION OF CARBETAMIDE RESIDUES
AND ITS ANILINE METABOLITE

In this article by Alvaro Guardigli, William Chow, and Morton S. Lefar [*J. Agr. Food Chem.* **20**(2), 348 (1972)], on page 348 the first sentence, last paragraph of the second column, should read: "To the acidified hydrolyzed aqueous extract was added 25 ml of benzene followed by 10 ml of a 0.01% solution of 4-bromobenzoyl chloride in benzene."

Correction

AMINO ACID COMPOSITION OF BUCKWHEAT

In this article by Yeshajahu Pomeranz and George S. Robbins [*J. Agr. Food Chem.* **20**(2), 270 (1972)], on page 272 the footnote to Table III should read: "All *absolute* figures above 0.632 are in italics and all figures above 0.765 are in bold face to indicate statistical significance."