

## CORRECTIONS

INHIBITION OF *CARPOPHILUS FREEMANI* DOBSON (COLEOPTERA: NITIDULIDAE) AGGREGATION PHEROMONE RESPONSE BY A Z-DOUBLE-BOND PHEROMONE ANALOG, by R. J. Petroski\* and D. Weisleder. *J. Agric. Food Chem.* **1997**, 45, 943–945.

Table 1 (below) was missing.

**Table 1. Inhibition of Beetle Response to Aggregation Pheromone (*E*) by the 6*Z* Isomer (*Z*)**

condition <sup>a</sup>			treatment effect <sup>b</sup>			
ng		SYN	mean counts		% inh	<i>F</i>
<i>E</i>	<i>Z</i>		<i>E</i>	<i>E</i> + <i>Z</i>		
0.13	0.13	Y	6	5	17	0.8
0.13	1.3	Y	32	17	47	10.1**
1.3	1.3	N	119	98	18	5.3*
1.3	1.3	Y	75	61	19	1.3
1.3	13.0	N	23	17	26	15.4**
1.3	13.0	Y	22	15	32	15.4**

<sup>a</sup> The synergist (SYN) was propyl acetate (20  $\mu$ L, 1% solution in mineral oil); Y when present, N when absent. <sup>b</sup> Bioassay counts are the numbers of beetles flying upwind to the filter paper baits and alighting during the 3-min tests. Test samples containing both the natural pheromone (*E*) and the 6*Z*-isomer (*Z*) are designated *E* + *Z*. Percent inhibition (% inh) is defined as  $100 \times (\text{mean counts } E - \text{mean counts } E + Z) / \text{mean counts } E$ .  $N = 16$ . The symbol *F* is the *F* statistic. df: treatment = 1,14; position = 1,14. \*\*,  $P < 0.01$ ; \*,  $P < 0.05$ . Average MSE (treatment  $\times$  test) was 0.028 (range 0.010–0.055).

JF980001Z

S0021-8561(98)00001-6

Published on Web 02/13/1998

EFFECT OF BLANCHING ON STRUCTURAL QUALITY OF DRIED POTATO SLICES, by J. I. Maté,\* C. G. M. I. Quataert, G. Meerdink, and K. van't Riet. *J. Agric. Food Chem.* **1998**, 46, 676.

C. G. M. I. Quataert's name was incorrectly spelled.

JF980073Q

S0021-8561(98)00073-9

Published on Web 02/19/1998