

Simple Preparation of 8–5-Coupled Diferulate

John Ralph,^{*,†,‡} Maria Teresa Garcia Conesa,[§] and Gary Williamson[§]

U.S. Dairy Forage Research Center, Agricultural Research Service, U.S. Department of Agriculture, and Department of Forestry, University of Wisconsin—Madison, Madison, Wisconsin 53706, and Department of Biochemistry, Institute of Food Research, Colney, Norwich NR4 7UA, United Kingdom

Diferulates are important products of ferulate-mediated cross-linking of plant cell walls. Coupling of ferulates by radical mechanisms produces a range of diferulates. One of the major isomers, 8–5-coupled diferulate, can be synthesized by single-electron oxidants, but the yields are relatively low and purification is nontrivial. Significant quantities of this dimer are required for various analytical and reactivity studies. A simpler preparation utilizing the biomimetic peroxidase–H₂O₂ system allows the production of gram quantities of pure diethyl 8–5-diferulate.


Keywords: Ferulate; diferulate; ferulic acid; diferulic acid; peroxidase; radical coupling; cross-linking

INTRODUCTION

Ferulates play important roles in plant growth and development. Polysaccharide–ferulate esters laid down in the plant cell wall undergo radical coupling reactions with other ferulates to produce ferulate dimers and effect polysaccharide–polysaccharide cross-linking (Ralph et al., 1994; Geissmann and Neukom, 1971; Fry, 1979) or with lignin units to effect lignin–polysaccharide cross-linking (Ralph et al., 1992, 1995). Ferulate dimers can also radically cross-couple with lignin units to effect even more substantial lignin–polysaccharide cross-linking (Quideau and Ralph, 1997; Ralph et al., 1997). Arising from radical processes, diferulates are produced in several regioisomeric forms, although this has only recently been recognized (Ralph et al., 1994). The 8–8-, 8–5-, and 8–O–4-coupled isomers, are now invariably found in higher concentrations than the 5–5-coupled isomer, which was previously the only diferulate noted (with the exception of photochemical [2 + 2] products).

Syntheses of all of the diferulates were reported with the initial discovery of the range of diferulates (Ralph et al., 1994). However, significant quantities are now required by cell wall researchers, and larger scale and simpler preparations are required. The 5–5-coupled dimer is relatively easily prepared, but the 8–O–4- and 8–8-coupled isomers remain in the domain of synthetic organic chemists. The 8–5-coupled product was prepared by a relatively simple method, utilizing silver oxide as the single-electron oxidant, but product separation was not trivial. It has been well established (Ralph et al., 1992; Teutonico et al., 1991; Chioccare et al., 1993; Wallace and Fry, 1995) that the primary ferulate dimer generated utilizing the biomimetic peroxidase–H₂O₂ system, or a range of single-electron oxidants, is the 8–5-coupled dimer, but a practical gram-scale prepara-

Scheme 1. Biomimetic Synthesis of 8–5-Coupled Diethyl Diferulate I^a



^a (i) EtOH/HCl, yield ~100%. (ii) peroxidase, H₂O₂, pH 4, 10 min, yield ~50+%.

tion has not emerged. Here we report a simple preparation that produces pure 8–5-coupled diethyl diferulate I cheaply and in reasonable yields (Scheme 1).

EXPERIMENTAL PROCEDURES

Ferulic acid and peroxidase (EC 1.11.1.7, type II, 150–200 units/mg, from horseradish) were obtained from Sigma; 30% H₂O₂ was from Mallinckrodt and acetyl chloride from Aldrich. Solvents were of AR grade. Petroleum ether was the 35–60 °C boiling fraction. All were used without further purification. Analytical TLC was performed with Alugram Sil-G/UV₂₅₄ plates (Macherey-Nagel) with visualization by UV light. Silica gel flash chromatography was on a Biotage Flash 40M flash chromatography unit using a 70 g prepacked column (32–63 μm particles, 60 Å average pore size, 570 m²/g surface area), utilizing an ISCO UA-6 UV–vis detector and an ISCO Foxy 200 fraction collector.

Ethyl Ferulate. Ethyl ferulate was prepared as described previously (Ralph et al., 1994) from ethanolic HCl, prepared conveniently according to Fieser's method (Fieser and Fieser, 1967). Thus, ferulic acid (10 g) was dissolved in absolute ethanol (100 mL) and acetyl chloride (5 mL) added. The solution was stirred gently overnight, and the volatiles were removed by rotary evaporation at 40 °C. Addition of further ethanol and evaporation several times removed the HCl. The product was then used directly for the diferulate step or allowed to crystallize from EtOAc/petroleum ether.

Dimerization. The ethyl ferulate (2 g) was dissolved, with heating (to ~60 °C), into pH 4.0 acetate buffer (2 L). After

* Address correspondence to this author at the U.S. Dairy Forage Research Center [fax (608) 264-5147; e-mail jralph@facstaff.wisc.edu].

[†] U.S. Dairy Forage Research Center.

[‡] Department of Forestry.

[§] Department of Biochemistry.

the solution cooled to $\sim 40^\circ\text{C}$, H_2O_2 (0.76 mL) was added and then peroxidase (10 mg in 2 mL of phosphate buffer). Within seconds, crude product began to precipitate out. After 10 min, the product was filtered through sintered glass yielding crude product (~ 2 g). Alternatively, the solution can be extracted directly with EtOAc.

Purification. TLC indicated that the required product was well separated from trace faster moving and more major slower moving uncharacterized contaminants. Flash chromatography using EtOAc–petroleum ether (30:70, ~ 30 mL/min) as eluant produced the product in fractions 37–50 (each 15 mL). Product **I** spontaneously crystallized on evaporation of the solvent (purified yield, 1.05 g, $\sim 50\%$). Although not really required, recrystallization from acetone–petroleum ether produced white needle crystals. NMR data and melting point, 152.8–153.1 $^\circ\text{C}$, were as detailed previously (Ralph et al., 1994).

DISCUSSION

Preparing phenylcoumarans of this type by purely synthetic methods without utilizing radical coupling reactions is tedious (Ede et al., 1987; Ralph et al., 1987; Brunow and Lundquist, 1984; Nakatsubo and Higuchi, 1979; Hise et al., 1985; Hassi et al., 1987). The silver(I) oxide method used previously (Ralph et al., 1994), and presumably other methods, produced the required product in $\sim 30\%$ yield, but it was contaminated by components from which separation was not trivial. The biomimetic peroxidase– H_2O_2 system, described under Experimental Procedures, produced the required 8–5-coupled product **I** quickly in reasonable yield; from the NMR spectra, the yield appeared to be very high, but polymeric components must have accounted for a substantial fraction. Importantly, those contaminants that were present had significantly different mobilities on silica, allowing simple separation. The yield of purified product was $\sim 50\%$. Although we used a commercial flash chromatography system, the separation could be easily carried out by simpler flash systems (Taber, 1982; Still et al., 1978), the sintered glass funnel method (Leopold, 1982), or any other nonpressurized chromatography system. We loaded the full 2 g of product onto Biotage's silica 40M packed column and, within ~ 20 min, had the pure 8–5-coupled diferulate separated. Scaling up the reaction did not produce greater quantities of the required product; optimization studies at different scales were not pursued, however.

This simple method should allow laboratories to synthesize their own stocks of this valuable compound. Unfortunately, the only other dimer so readily prepared at this point is the 5–5-coupled dimer. Methods to produce high yields of 8–O–4- or 8–8-coupling, utilizing the single-electron systems described by Landucci (1995) for favoring these products in monolignol coupling reactions, are being explored but have so far been disappointing for diferulates.

LITERATURE CITED

- Brunow, G.; Lundquist, K. A new synthesis of model compounds for the β -5 structural unit in lignins. *Acta Chem. Scand.* **1984**, *B38*, 335–336.
- Chioccare, F.; Poli, S.; Rindone, B.; Pilati, T.; Brunow, G.; Pietikainen, P.; Setälä, H. Regio- and diastereoselective synthesis of dimeric lignans using oxidative coupling. *Acta Chem. Scand.* **1993**, *47*, 610–616.
- Ede, R. M.; Ralph, J.; Wilkins, A. L. The stereochemistry of β -5 lignin model compounds. *Holzforchung* **1987**, *41*, 239–45.
- Fieser, L. F.; Fieser, M. Reagents for Organic Synthesis. In *Reagents for Organic Synthesis*; Wiley: New York, 1967; p 192.
- Fry, S. C. Phenolic components of the primary cell wall and their possible role in the hormonal regulation of growth. *Planta* **1979**, *146*, 343–351.
- Geissmann, T.; Neukom, H. Vernetzung von Phenolcarbonsäureestern von Polysacchariden durch oxydative phenolische Kupplung. *Helv. Chim. Acta* **1971**, *54*, 1108–1112.
- Hassi, H. Y.; Aoyama, M.; Tai, D.; Chen, C. L.; Gratzl, J. S. Substituent effects on carbon-13 chemical shifts of aromatic carbons in β -O-4 and β -5 type lignin model compounds. *J. Wood Chem. Technol.* **1987**, *7*, 555–581.
- Hise, R. G.; Chen, C. L.; Gratzl, J. S. Synthesis of β -aryl lignin model compounds. *J. Wood Chem. Technol.* **1985**, *5*, 379–390.
- Landucci, L. L. Reactions of *p*-hydroxycinnamyl alcohols with transition metal salts. 1. Oligolignols and poly(lignols) (DHPs) from coniferyl alcohol. *J. Wood Chem. Technol.* **1995**, *15*, 349–368.
- Leopold, E. J. Vacuum dry column chromatography. *J. Org. Chem.* **1982**, *47*, 4592–4594.
- Nakatsubo, F.; Higuchi, T. Syntheses of phenylcoumarans. *Mokuzai Gakkaishi* **1979**, *25*, 735–742.
- Quideau, S.; Ralph, J. Lignin-ferulate cross-links in grasses. Part 4. Incorporation of 5–5-coupled diferulate into lignin. *J. Chem. Soc., Perkin Trans. 1* **1997**, 2351–2358.
- Ralph, J.; Ede, R. M.; Robinson, N. P.; Main, L. Reactions of β -aryl lignin model quinone methides with anthrahydroquinone and anthranol. *J. Wood Chem. Technol.* **1987**, *7*, 133–60.
- Ralph, J.; Helm, R. F.; Quideau, S.; Hatfield, R. D. Lignin-feruloyl ester cross-links in grasses. Part 1. Incorporation of feruloyl esters into coniferyl alcohol dehydrogenation polymers. *J. Chem. Soc., Perkin Trans. 1* **1992**, 2961–2969.
- Ralph, J.; Quideau, S.; Grabber, J. H.; Hatfield, R. D. Identification and synthesis of new ferulic acid dehydrodimers present in grass cell walls. *J. Chem. Soc., Perkin Trans. 1* **1994**, 3485–3498.
- Ralph, J.; Grabber, J. H.; Hatfield, R. D. Lignin-ferulate cross-links in grasses: active incorporation of ferulate polysaccharide esters into ryegrass lignins. *Carbohydr. Res.* **1995**, *275*, 167–178.
- Ralph, J.; Hatfield, R. D.; Grabber, J. H.; Jung, H. G.; Quideau, S.; Helm, R. F. Cell Wall Cross-linking in Grasses by Ferulates and Diferulates. In *Lignin and Lignan Biosynthesis*; Lewis, N. G., Sarkanen, S., Eds.; American Chemical Society: Washington, DC, 1998; pp 209–236.
- Still, W. C.; Kahn, M.; Mitra, A. Rapid chromatographic technique for preparative separations with moderate resolution. *J. Org. Chem.* **1978**, *43*, 2923–2925.
- Taber, D. F. TLC mesh column chromatography. *J. Org. Chem.* **1982**, *47*, 1351–1352.
- Teutonico, R. A.; Dudley, M. W.; Orr, J. D.; Lynn, D. G.; Binns, A. N. Activity and accumulation of cell division-promoting phenolics in tobacco tissue cultures. *Plant Physiol.* **1991**, *97*, 288–297.
- Wallace, G.; Fry, S. C. In vitro peroxidase-catalyzed oxidation of ferulic acid esters. *Phytochemistry* **1995**, *39*, 1293–1299.

Received for review February 10, 1998. Revised manuscript received April 30, 1998. Accepted May 19, 1998. We are grateful to IFR for funding M.T.G.C.'s visit to the Dairy Forage Research Center to synthesize ferulate dimers and to Fachuang Lu for valuable help. G.W. and M.T.G.C. were funded by the Biotechnology and Biological Sciences Research Council, U.K., and by the European Union (FAIR-CT96-1099). The work was also supported in part by a USDA-NRI Competitive Grant 96-35304 (Plant Growth and Development section).

JF980123R