

## Fungal Flora and Ochratoxin A Production in Grapes and Musts from France

LUCILE SAGE<sup>†</sup>, SERGE KRIVOBOK<sup>†</sup>, ÉMILIE DELBOS<sup>†</sup>,  
FRANÇOISE SEIGLE-MURANDI,<sup>\*,†</sup> AND EDMOND E. CREPPY<sup>‡</sup>

Groupe pour l'Étude du Devenir des Xénobiotiques dans l'Environnement (GEDEXE), UFR de Pharmacie, Université Joseph Fourier, BP 138, 38243 Meylan Cedex, France, and Laboratoire de Toxicologie et d'Hygiène Appliquée, UFR de Pharmacie, Université Bordeaux II, 3 ter Place de la Victoire, 33076 Bordeaux Cedex, France

Eleven samples of grapes and musts used in red table wines were investigated for the occurrence of potential ochratoxin A (OTA)-producing molds. From these samples, 59 filamentous fungi and 2 yeasts were isolated. Among the 30 genera isolated, Deuteromycetes were the most frequent (70%) followed by Ascomycetes (10%). Six of the eleven grapes samples were contaminated by potentially ochratoxinogenic strains (*Penicillium chrysogenum* and *Aspergillus carbonarius*). When cultivated in vitro on solid complex media, the 14 strains of *A. carbonarius* produced OTA. No other species produced OTA under the same conditions. Among must samples, eight of eleven were found to be contaminated by OTA (concentrations from <10 to 461 ng/L). There is a strong correlation between the presence of ochratoxin-producing strains on grapes and OTA in musts. These findings should be connected with the OTA contamination of human blood in these areas and in France.

**KEYWORDS:** Ochratoxin A; *Penicillium*; *Aspergillus*; grapes; musts

### INTRODUCTION

First isolated by van der Merwe et al. (1) in cultures of *Aspergillus ochraceus* Wilhelm, ochratoxin A (OTA) was later found in the culture media of various species of *Aspergillus* and *Penicillium* (2–5). OTA is a widespread contaminant in human food and animal feed, as reviewed by IARC (4), Pittet (6), SCOOP (7), and Speijers and van Egmond (8). It has been shown that besides various toxic effects, OTA was nephrotoxic for several animal species, especially swine (9, 10). A kidney disease (Balkan endemic nephropathy) was found to be associated with the presence of OTA in food and the blood of humans suffering from tubulo interstitial nephropathy associated with urinary tract tumors (11, 12).

OTA is also immunosuppressive (13, 14), teratogenic (15, 16), genotoxic (17–20), and carcinogenic (21, 22) in rodents. Therefore, the International Agency for Research on Cancer has classified it in group 2B as a possible human carcinogen (4). OTA had been found in various countries in a large variety of foods: cereals, vegetables, dried fruits, nuts, and meats (pork, poultry, fish, etc.) (23–30).

Several studies have also found OTA in beverages (wine, grape juice, beer, milk, etc.) (31–34). First found in various Swiss table wines (35), OTA was also found in wines from different regions (36, 37) and in dried vine fruit (38). The goal of this study was to obtain further quantitative data on the

presence of OTA in grapes and musts used in the Aude department (Southern France) for red table wines and of ochratoxinogenic strains on grapes to estimate the potential contribution of wine in the daily intake of OTA in humans, especially in Southern France.

### MATERIALS AND METHODS

**Isolation of Fungi from Grapes.** We collected 11 samples (1 kg each) of grapes from Southern France used for the making of Aude wines. For each of them, 50 g was collected to ensure a representative sample according to Horwitz et al. (39) and suspended in 100 mL of sterile water containing SDS (0.05%, w/v). After 1 h of magnetic shaking, 1 mL of each suspension was deposited in a Petri dish (90 mm diameter) on malt extract (1.5%)/agar (1.5%) medium (MEA) complemented with chloramphenicol (0.05%, w/v), and 1 mL of each of the same suspensions was deposited in a Petri dish and the culture medium poured over it following the soil plates method of Warcup (40). The plates were incubated at 24, 30, 37, and 45 °C. *Aspergilli* were identified on Czapek–Dox medium (41), *Penicillia* on media proposed by Pitt (42), and *Fusaria* on PDA medium (43). The purity of each strain and its identity were checked following the morphological criteria of Zycha and Siepman (44), Booth (45), Ellis (46), Raper and Fennel (47), Pitt (48), Samson (49), Al-Musallam (50), and Von Arx (51).

**OTA Production by Isolated Strains.** Strains were grown on MEA medium for 1 week before inoculation at three points in 9 cm Petri dishes containing Yeast Extract Sucrose (YES) agar with 0.5 g/L magnesium sulfate and Czapek Yeast Autolysate (CYA) agar. OTA production was analyzed after 7 and 14 days of growth at 24 °C as

<sup>†</sup> Université Joseph Fourier.

<sup>‡</sup> Université Bordeaux II.

Table 1. Fungal Strains Isolated from Grape and Their Production of OTA

class/genus (nomenclature)	A	B	C	D	E	F	G	H	J	K	M
Ascomycetes											
<i>Coniochaeta velutina</i> (Fuckel) Cooke		+									
<i>Emericella nidulans</i> (Eidam) Vuillemin		+									
<i>Neosartorya fischeri</i> var. <i>fischeri</i> (Wehmer) Malloch & Cain								+			
<i>Sordaria conoidea</i> (Ehrenberg) Hughes								+			
<i>Sordaria fimicola</i> (Rob.) Cesati & De Notaris								+			+
<i>Sordaria lappae</i> Potebnia									+		
Basidiomycete											
Deuteromycetes											
<i>Acremonium alternatum</i> Link	+										
<i>Alternaria alternata</i> (Fries) Keissler	+	+	+		+	+	+	+	+	+	
<i>Aspergillus aculeatus</i> Iizuka	+	+	+		+						
<i>Aspergillus carbonarius</i> (Bainier) Thom <sup>a</sup>	+	+		+	+						
<i>Aspergillus fumigatus</i> Fresenius	+	+									
<i>Aspergillus</i> section <i>Nigri</i>	+	+	+	+	+		+			+	+
<i>Aspergillus terreus</i> Thom				+							
<i>Aspergillus ustus</i> (Bainier) Thom & Church											+
<i>Aureobasidium pullulans</i> (de Bary) Arnaud		+						+			
<i>Botrytis cinerea</i> Persoon ex Fries	+			+	+	+	+	+	+		
<i>Cladosporium cladosporioides</i> (Fresenius) de Vries	+	+		+	+		+				
<i>Cladosporium elatum</i> (Harz) Nannfeldt								+			
<i>Curvularia clavata</i> Jain		+									
<i>Curvularia inaequalis</i> (Shaer) Boedijn	+	+									
<i>Curvularia lunata</i> (Wakker) Boedijn								+			
<i>Drechslera spicifera</i> (Bainier) von Arx		+	+								
<i>Epicoccum nigrum</i> Link		+	+				+	+			+
<i>Fusarium moniliforme</i> Sheldon	+										
<i>Fusarium solani</i> (Martius) Saccardo											+
<i>Geotrichum candidum</i> Link ex Persoon								+			
<i>Gliocladium roseum</i> Bainier	+							+			
<i>Harzia verrucosa</i> (Tognini) Holubova Jechova								+			
<i>Microdochium bolleyi</i> (Sprague) de Hoog & Hermanides-Nijhof								+			
<i>Penicillium atramentosum</i> Thom		+				+					
<i>Penicillium brevicompactum</i> Dierckx		+	+			+	+	+	+	+	+
<i>Penicillium canescens</i> Sopp								+			
<i>Penicillium chrysogenum</i> Thom <sup>b</sup>		+						+	+		
<i>Penicillium citreonigrum</i> Dierckx		+									
<i>Penicillium citrinum</i> Thom								+			
<i>Penicillium echinulatum</i> Fassatiava								+			
<i>Penicillium expansum</i> Link					+		+	+	+	+	+
<i>Penicillium glabrum</i> (Wehmer) Westling	+	+			+		+				
<i>Penicillium griseofulvum</i> Dierckx								+			+
<i>Penicillium miczynskii</i> Zaleski		+								+	+
<i>Penicillium minioluteum</i> Dierckx		+	+								
<i>Penicillium oxalicum</i> Currie & Thom		+									
<i>Penicillium paxilli</i> Bainier	+										
<i>Penicillium purpurogenum</i> Stoll	+	+		+							+
<i>Penicillium rugulosum</i> Thom				+							
<i>Penicillium spinulosum</i> Thom		+									
<i>Penicillium thomii</i> Maire	+			+	+		+		+		+
<i>Phoma eupyrena</i> Saccardo	+		+			+		+			
<i>Phoma putaminum</i> Spegazzini	+										
<i>Rhodotorula aurantiaca</i> (Saito) Lodder	+		+					+			
<i>Scopulariopsis brevicaulis</i> (Saccardo) Bainier								+	+		
<i>Stemphylium botryosum</i> Wallroth	+										
<i>Trichoderma harzianum</i> Rifai	+	+						+			+
<i>Ulocladium chartarum</i> (Preuss) Simmons	+	+	+					+			
<i>Verticillium fungicola</i> (Preuss) Hassebrauk		+	+					+			
White Yeast spp.	+					+			+		
Zygomycetes											
<i>Actinomyces elegans</i> (Eidam) Benjamin & Hesseltine			+								
<i>Rhizopus stolonifer</i> (Ehrenberg ex Fries) Lind											+

<sup>a</sup> Strains actually producing OTA in solid YES and/or CYA media. <sup>b</sup> Potentially toxinogenic strains.

previously described (52). The YES medium contained (g/L) sucrose (Prolabo) (40.0) and yeast extract (Difco, Detroit, MI) (20.0). The CYA medium contained (g/L) sucrose (Prolabo) (40.0) and yeast extract (Difco) (20.0). Sterilization was carried out by autoclaving for 20 min at 121 °C. After sterilization, the pH of the media was 6.5.

**OTA Extraction from Culture.** At each sampling, three agar plugs were removed from the central area of the colony. Each plug was weighed and introduced into a small vial, methanol (0.5 mL) was added,

and the vial was shaken for 1 h (orbital shaker, 200 rpm). The extracts were filtered (Millipore Corp., Bedford, MA) and injected into an HPLC instrument.

**OTA Extraction from Musts.** The method described by Visconti et al. (53) was used. Each sample (100 mL) was diluted with a solution containing 1% poly(ethylene glycol) and 5% sodium hydrogenocarbonate, filtered, and applied to an OchraTest immunoaffinity column (Vicom Inc., Watertown, MA). The column was washed with a solution

**Table 2.** Levels of OTA Found in Must Samples and Its Production by Toxinogenic Strains Isolated from the Same Samples

must sample	A	B	C	D	E	F	G	H	J	M	K
[OTA] (ng/L)	a	58		318	461	<10	150	51	20		
				[OTA] ( $\mu\text{g/g}$ of medium)							
				YES <sup>a</sup>				CYA			
isolated strain	must sample			7 days	14 days			7 days	14 days		
<i>A. carbonarius</i>	A			62 ± 2.1	81.1 ± 0.6			66.4 ± 1.0	87.5 ± 1.1		
<i>A. carbonarius</i>	B			traces	0.1 ± 0.0						
<i>A. carbonarius</i>	B			traces	0.2 ± 0.0			2.0 ± 1.1	1.2 ± 0.1		
<i>A. carbonarius</i>	D				1.5 ± 0.3			19.9 ± 0.6	18.5 ± 0.6		
<i>A. carbonarius</i>	D			0.6 ± 0.1	0.6 ± 0.2			12.0 ± 1.4	11.7 ± 0.9		
<i>A. carbonarius</i>	D			0.1 ± 0.0	0.4 ± 0.1			2.5 ± 0.1	3.4 ± 0.2		
<i>A. carbonarius</i>	D				0.1 ± 0.0			5.5 ± 0.0	6.9 ± 1.0		
<i>A. carbonarius</i>	E			0.2 ± 0.0	1.5 ± 0.1				0.3 ± 0.1		
<i>A. carbonarius</i>	E			0.3 ± 0.0	0.9 ± 0.1			7.8 ± 0.0	9.0 ± 0.4		
<i>A. carbonarius</i>	E			0.1 ± 0.0	0.2 ± 0.0			0.5 ± 0.2	0.3 ± 0.1		
<i>A. carbonarius</i>	E			1.9 ± 0.5	0.7 ± 0.0			13.1 ± 2.0	8.8 ± 0.1		
<i>A. carbonarius</i>	E			24.6 ± 2.5	54.6 ± 2.0			65.4 ± 2.2	74.7 ± 1.8		
<i>A. carbonarius</i>	E			1.2 ± 0.4	0.9 ± 0.4			9.5 ± 0.3	7.3 ± 0.4		
<i>A. carbonarius</i>	E			0.6 ± 0.1	0.9 ± 0.1			15.1 ± 2.4	9.2 ± 0.6		
<i>A. section Nigri</i>	A–E, G, K, M										
<i>P. chrysogenum</i>	B, H, J										
controls											
<i>A. alliaceus</i>								239.5 ± 6.5			
<i>A. ochraceus</i>								15.4 ± 0.3			

<sup>a</sup> OTA not quantified.

containing sodium chloride (2.5%) and sodium hydrogenocarbonate (0.5%) followed by water. OTA was eluted with methanol and quantified by reversed-phase HPLC. OTA-free samples were spiked with OTA and treated in the same way. About 90% of the OTA was extracted.

**Quantification of OTA by HPLC.** HPLC was performed with a liquid chromatograph (Shimadzu) equipped with an LC 6A, an SIL-9A automatic injector, a diode array UV detector (SPD-M10A VP), and an RF-530 fluorescence detector with a column ( $\varnothing = 4.6$  mm,  $L = 250$  mm) filled with  $C_{18}$  ODS-Hypersil, 5  $\mu\text{m}$ . The mobile phase was acetonitrile/water/acetic acid (99:99:2; pH 3.5). Flow rate: 1 mL/min. Detection: (excitation) 333 nm, (emission) 460 nm. Quantification was achieved with a computing integrator (RC-6A). Following Soleas et al. (34), a five-point calibration prepared by adding different volumes of OTA stock standard solution to an OTA-free sample was used with concentrations ranging from 0.5 to 10  $\mu\text{g/L}$ . For the confirmation of OTA production, diode array detection was used. OTA was identified by comparison with pure toxin (Sigma Chemical, St. Louis, MO) used as both external and internal standards.

## RESULTS AND DISCUSSION

**Fungal Strain Isolation and Production of OTA Detection by Strains.** Out of the 11 samples of grapes, 59 filamentous fungi and 2 yeasts were isolated (Table 1). The 30 identified genera belong to Deuteromycetes (70%) and Ascomycetes (10%). *Penicillium* (31%) and *Aspergillus* (10%) were the most often encountered genera. *Alternaria alternata* (9/11 samples), *Aspergillus* gr. *niger* (8/11), *Penicillium brevicompactum* (8/11), and *Botrytis cinerea* (7/11) were the natural microfungi from the grape samples most often identified. Though grape sample B was weakly contaminated by fungi, 25 different strains were identified in the cultures. On the contrary, heavily moldy sample E gave only 9 different fungal strains, especially *A. gr. niger* and *Penicillium thomii*.

The toxinogenic potential of the isolated fungal strains was compared to *Aspergillus alliaceus* and *A. ochraceus*. Among the strains isolated from grapes, potentially OTA producers were *Aspergillus carbonarius* (14 strains out of 15) and *Penicillium chrysogenum* (3 strains) (Tables 1 and 2). Six of the eleven

grape samples (A, B, D, E, H, and J) were contaminated by these ochratoxinogenic fungi (Table 2).

*A. carbonarius* was the only species that produced OTA media after 7 or 14 days when cultivated on YES or CYA. According to Bragulat et al. (52), the CYA medium is better adapted for OTA production than the YES medium except for two weakly ochratoxinogenic strains. However, concentrations of OTA in CYA medium were much lower than those given by the *A. alliaceus* reference strain (239.5  $\mu\text{g/g}$ ) adapted to the culture on a solid medium: the production ranged from 0.5 to 87.5  $\mu\text{g/g}$  for *A. carbonarius* strains and 15.4  $\mu\text{g/g}$  for *A. ochraceus*. As the production of OTA by *A. niger* var. *niger* has recently been reported (54), 73 *Aspergillus* section *Nigri* strains isolated from A–E, G, K, and M grape samples were also evaluated for OTA production. After 7 and 14 days of cultivation on YES or CYA medium, none produced any OTA.

The three strains of *P. chrysogenum* isolated from grapes were not toxinogenic, which is consistent with the results of Frisvad (55). He performed a survey for the production of mycotoxins by *Penicillium* strains supposed to be toxinogenic selected from the main mycotheques and found that, for several reasons, many *Penicillia* had been misidentified. Finally, they could not detect OTA production by any of the 230 *P. chrysogenum* strains examined. Several other surveys on the fungal contamination of foods have also revealed the low percentage of OTA-producing *Penicillia* (3, 56–58).

The high frequency of the genus *Aspergillus* in the strains isolated from grapes is consistent with the results of Creppy et al. (59), who outline that, in France, ochratoxicosis seems to be connected with the presence of *Aspergilli* while, in Germany and in Scandinavia, it would be connected with the presence of *Penicillia*, owing to their different temperature needs (60): ochratoxinogenic *Penicillia* grow well over a wide range of temperature (4–31 °C), whereas *Aspergilli* that produce OTA require higher temperatures (12–39 °C).

**OTA Detection in Musts.** In this study, 8 out of 11 must samples were found to be contaminated by OTA (Table 2).

The concentrations were estimated from 461 to less than 10 ng/L, the higher concentration being found in sample E. There is a strong correlation between the presence of ochratoxin-producing strains on grapes and the finding of OTA in musts. OTA has been found in various agricultural commodities (almost all cereals, animal feeds, other food commodities), as reviewed by Pittet (6). Recently, there have been several reports of the presence of OTA in wine and grape juice (35, 37, 61–63). Surprisingly high concentrations of this mycotoxin were also detected in raisins from Greece and Turkey (36). Otteneder and Majerus (37) have realized a wide review to describe extensively the situation of OTA contamination of wine. They analyzed 400 samples themselves and collected more than 450 results from the literature. According to these data, OTA is much more commonly detected in red wines (54%) than in rosé (40%) and white (25%) wines. The OTA concentration is remarkably higher in red wines (22–1153 ng/L) than in rosé (119 ng/L) and white (12–108 ng/L) wines, depending on the regions. This situation can be explained by the different processing methods: whereas, for white wines, the grapes are immediately pressed, red grapes are left mashed for several days, which obviously permits fungal growth and production of the toxin. Provisional estimates of the Codex Alimentarius Commission, based on limited European data, suggest that red wine is the second major source of human exposure to OTA following cereals and preceding coffee and beer (64).

**Risk Assessment.** Health concerns regarding mycotoxins depend on the toxicity and the amount of mycotoxin consumed, body weight, the general health conditions of the individual, the presence of other mycotoxins, and other dietary factors. In the present work, except for OTA production, no potentially mycotoxinogenic strains were found in grape samples. It is important to note that others mycotoxins have already been found in wines: trichothecenes from the fungus *Trichothecium roseum* (65) and patulin (66).

OTA is genotoxic (17–19) and carcinogenic (21, 22) in rodents. It can be considered as one of the most potent renal carcinogens. Owing to the high level of potential occurrence of OTA in foods and beverages and its relatively long half-life in humans, 35 days in serum, following Studer-Rohr (67), the high incidence of this mycotoxin in human sera and milk is not surprising (7, 68–70). All these data show that OTA is among the most frequent mycotoxins found in human blood in Europe, the U.S., Canada, and elsewhere where it has been investigated (71).

At the beginning of the 1990s, the occurrence of OTA was not systematically searched for, and therefore it is almost impossible to establish any conclusion on the possible health hazard for the population resulting from OTA intake. In 1994, the European Commission set up a Scientific Co-Operation on Questions Relating to Food (7) task force to provide the Scientific Committee for Food with information on European dietary exposure to OTA. As a result, the mean dietary intake of humans, based on occurrence and consumption data, was established as 1.8 ng/kg bw per day, and based on human blood data, as 0.9 ng/kg bw per day. Because of the increase of scientific reports on OTA contamination in beverages and many kinds of food, the WHO/FAO Joint Expert Committee on Food Additives (JECFA) proposed 100 ng/kg bw per day as a provisional tolerable weekly intake (PTWI) for OTA (72). Ueno et al. (33) stated that this exposure estimate is not deemed to represent a health hazard. In 1998, The Scientific Committee on Food of the European Union considered it would be advisable to reduce the tolerable daily OTA intake from food as low as

possible, i.e., between 1.2 and 14 ng/kg bw per day or lower than 5 ng/kg bw per day (73).

## CONCLUSION

The aim of this research was to record the fungal population in grapes, particularly ochratoxinogenic strains, and to see the possible connection between the presence of OTA in wine. In the present work, 73% of the must samples contained OTA, produced by strains of *A. carbonarius*. The rate of contamination ranged from 461 to less than 10 ng/L.

These OTA contamination levels were in accordance with previous studies (35, 37, 61–63). They could explain the human blood content contamination in France of 22–25% of the general population with levels from 0.1 to 130 ng/mL (68). Although the number of samples was limited to 11, this study shows a real hazard for the population in the southwestern part of France. Wine was pointed out as one of the possible sources of OTA in human plasma (33, 37).

The most efficient way to protect consumers against OTA health hazards is therefore to implement good agricultural practice to lower the presence of fungal strains on the grape and therefore reduce the possibility of OTA production during the wine-making process. An appropriate antifungal treatment of vines and a tight control of the wine-making process would be the main factors.

A new study on grapes (end of veraison and harvest) and musts is in progress in various regions in France. It should allow inventory in each region of ochratoxinogenic strains on grapes at various stages of ripeness, to establish the connection with antifungal treatments applied on the vine and the presence of OTA in musts.

## LITERATURE CITED

- (1) Merwe, van der, K. J.; Steyn, P. S.; Fourie, L.; de Scott, B.; Theron, J. J. Ochratoxin A, a toxic metabolite produced by *Aspergillus ochraceus* Wilh. *Nature (London)* **1965**, *205*, 1112–1113.
- (2) Krogh, P. Ochratoxins. In *Mycotoxins in human and animal health*; Rodricks, J. V., Hesseltine, C. W., Mehlman, M. A., Eds.; Pathotox Publishers, Inc.: Peoria, Ill, 1977; pp 489–498.
- (3) Pitt, J. I. *Penicillium viridicatum*, *Penicillium verrucosum* and production of ochratoxin A. *Appl. Environ. Microbiol.* **1987**, *53*, 266–269.
- (4) IARC (International Agency for Research on Cancer). *Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins*; World Health Organization: Lyon, France, 1993; Vol. 56.
- (5) Varga, J.; Kevei, E.; Rinyu, E.; Téren, J.; Kozakiewicz, Z. Ochratoxin production by *Aspergillus* species. *Appl. Environ. Microbiol.* **1996**, *62*, 4461–4464.
- (6) Pittet, A. Natural occurrence of mycotoxins in foods and feeds. An updated review. *Rev. Méd. Vét.* **1998**, *149*, 479–492.
- (7) *Assessment of dietary intake of ochratoxin A by the population of EU member states*; SCOOP Report; European Commission, DG III Publisher: 1996.
- (8) Speijers, G. J. A.; van Egmond, H. P. Worldwide ochratoxin A levels in food and feeds. In *Human ochratoxicosis and its pathologies*; Creppy, E. E., Castegnaro, M., Dirheimer, G., Eds.; J. Libbey Eurotext No. 231; Libbey: Montrouge, France, 1993; pp 199–207.
- (9) Krogh, P.; Axelsen, N. H.; Elling, F.; Gyrd Hansen, N.; Hald, B.; Hyldgaard-Jensen, J.; Larsen, A. E.; Madsen, A.; Mortensen, H. P.; Moller, T.; Petersen, O. K.; Ravnskov, U.; Rostgaard, M.; Aalund, O. Experimental porcine nephropathy: Changes of renal function and structure induced by ochratoxin A-contaminated feed. *Acta Pathol. Microbiol. Scand., Sect. A* **1974**, Suppl. 246, 9 pp.

- (10) Mortensen, H. P.; Hald, B.; Madsen, A. Feeding experiments with ochratoxin A contaminated barley for bacon pigs. 5. Ochratoxin A in pig blood. *Acta Agric. Scand.* **1983**, *33*, 235–239.
- (11) Krogh, P.; Hald, B.; Plestina, R.; Ceovic, S. Balkans (endemic) nephropathy and foodborne ochratoxin A: preliminary results of a survey of foodstuffs. *Acta Pathol. Microbiol. Scand., Sect. B* **1977**, *85*, 238–240.
- (12) Castegnaro, M.; Bartsch, H.; Chernozemsky, I. N. Endemic nephropathy and urinary-tract tumors in the Balkans. *Cancer Res.* **1987**, *47*, 3608–3609.
- (13) Haubeck, H. D.; Lorkowski, G.; Kölsh, E.; Rösenthaller, R. Immunosuppression by ochratoxin A and its prevention by phenylalanine. *Appl. Environ. Microbiol.* **1981**, *41*, 1040–1042.
- (14) Creppy, E. E.; Stormer, F. C.; Rösenthaller, R.; Dirheimer, G. Effects of two metabolites of ochratoxin A, (4R)-4-hydroxyochratoxin A and ochratoxin alpha on immune response in mice. *Infect. Immunol.* **1983**, *39*, 1015–1018.
- (15) Arora, R. G.; Frörelén, H. Interference of mycotoxins with prenatal development of the mouse. II. Ochratoxin A induced teratogenic effects in relation to the dose and stage of gestation. *Acta Vet. Scand.* **1981**, *22*, 535–552.
- (16) Mayura, K.; Edwards, J. F.; Maull, E. A.; Phillips, T. D. The effects of ochratoxin A on postimplantation rat embryos in culture. *Arch. Environ. Contam. Toxicol.* **1989**, *18*, 411–415.
- (17) Creppy, E. E.; Kane, A.; Dirheimer, G.; Lafarge-Frayssinet, C.; Mousset, S.; Frayssinet, C. Genotoxicity of ochratoxin A in mice: DNA single strand break evaluation in spleen, liver and kidney. *Toxicol. Lett.* **1985**, *28*, 29–35.
- (18) Pfohl-Leskowicz, A.; Chakor, K.; Creppy, E. E.; Dirheimer, G. DNA-adducts formation in mice treated with ochratoxin A. In *Mycotoxins, Endemic nephropathy and urinary tract tumours*; Castegnaro, M., Plestina, R., Dirheimer, G., Chernozemsky, I. N., Bartsch, H., Eds.; IARC Scientific publication No. 115; IARC: Lyon, France, 1991; pp 25545–253.
- (19) Obrecht-Pflumio, S.; Chassat, T.; Dirheimer, G.; Marzin, D. Genotoxicity of ochratoxin A by *Salmonella* mutagenicity test after bioactivation by mouse kidney microsomes. *Mutat. Res.* **1999**, *446*, 95–102.
- (20) Dirheimer, G. Recent advances in the genotoxicity of mycotoxins. *Rev. Méd. Vét.* **1998**, *149*, 605–616.
- (21) Kanisawa, M.; Suzuki, S. Induction of renal and hepatic tumours in mice by ochratoxin A, a mycotoxin. *Gann* **1978**, *69*, 599–600.
- (22) Boorman, G., Ed. NTP Technical report on the toxicology and carcinogenesis studies of ochratoxin A (CAS No. 303-47-9) in F344/N Rats (Gavage Studies) (NIH Publication No. 89-2813); National Toxicology Program, U.S. Department of Health and Human Services: Research Triangle Park, NC, 1989.
- (23) Krogh, P. Ochratoxin A residues in tissues of slaughter pigs with nephropathy. *Nord. Vet. Med.* **1977**, *29*, 402–405.
- (24) Chelkowski, J.; Golinski, P. Mycotoxins and toxicogenic fungi in mixed feeds and their cereal components. *Proceedings of the Fifth IUPAC Symposium on Mycotoxins and Phycotoxins. I*; Technical University: Vienna, 1982; pp 68–71.
- (25) Krogh, P.; Nesheim, S. Ochratoxin A. In *Environmental Carcinogens. Selected methods of analysis, Vol. 5, Mycotoxins*; Egan, H., Stoloff, L., Castegnaro, M., Scott, P., O'Neill, I. K., Bartsch, H., Eds.; IARC Scientific Publication No. 44; International Agency for Research on Cancer: Lyon, France, 1983; pp 247–253.
- (26) Scudamore, K. A.; Clarke, J. H.; Atkin, P. M. Natural occurrence of the fungal naphthoquinones in cereals and animal feedstuffs. In *Spoilage and mycotoxins of cereals and other stored products*; Flannigan, B., Ed.; International Biodeterioration 22 (supplement); 1986; pp 81–84.
- (27) Krogh, P. Ochratoxin in food. In *Mycotoxins in food*. Krogh, P., Ed.; Academic Press: London, 1987; pp 97–120.
- (28) Kuiper-Goodman, T.; Scott, P. M. Risk assessment of the mycotoxin ochratoxin A. *Biomed. Environ. Sci.* **1989**, *2*, 179–248.
- (29) Pohland, A. E.; Nesheim, S.; Friedman, L. Ochratoxin A: A review. *Pure Appl. Chem.* **1992**, *64*, 1029–1046.
- (30) Krivobok, S.; Seigle-Murandi, F.; Steiman, R.; Creppy, E. Fungal flora and ochratoxin A in various food and feed in France. *Syst. Appl. Microbiol.* **1995**, *18*, 455–459.
- (31) Breitholtz-Emanuelsson, A.; Olsen, M.; Oskarsson, A.; Palminge, I.; Hult, K. Ochratoxin A in cows milk and human milk with corresponding human blood samples. *J. Assoc. Off. Anal. Chem. Int.* **1993**, *76*, 842–846.
- (32) Scott, P. M.; Kanhere, S. R. Determination of ochratoxin A in beer. *Food Addit. Contam.* **1995**, *12*, 592–598.
- (33) Ueno, Y.; Maki, S.; Lin, J.; Furuya, M.; Sugiura, Y.; Kawamura, O. A 4-year study of plasma ochratoxin A in a selected population in Tokyo by immunoassay and immunoaffinity column-linked HPLC. *Food Chem. Toxicol.* **1998**, *36*, 445–449.
- (34) Soleas, G. J.; Yan, J.; Goldberg, D. M. Assay of ochratoxin A in wine and beer by high-pressure liquid chromatography photodiode array and gas chromatography mass selective detection. *J. Agric. Food Chem.* **2001**, *49*, 2733–2740.
- (35) Zimmerli, B.; Dick, R. Ochratoxin A in table wine and grape-juice: occurrence and risk assessment. *Food Addit. Contam.* **1996**, *13*, 655–668.
- (36) MAFF (Ministry of Agriculture, Fisheries and Foods). *Survey of aflatoxins and ochratoxin A in cereals and retail products*; Food Surveillance Information Sheet No. 130; Nov 1997.
- (37) Otteneder, H.; Majerus, P. Occurrence of ochratoxin A (OTA) in wines: influence of the type of wine and its geographical origin. *Food Addit. Contam.* **2000**, *17*, 793–798.
- (38) MacDonald, S.; Wilson, P.; Barnes, K.; Damant, A.; Massey, R.; Mortby, E.; Shepherd, M. J. Ochratoxin A in dried vine fruit: method development and survey. *Food Addit. Contam.* **1999**, *16*, 253–260.
- (39) Horwitz, W.; Senzel, A.; Reynolds, H.; Park, D. L., Eds. Natural poisons. *Official Methods of Analysis of the Association of Official Analytical Chemists*; AOAC: Washington, DC, 1975; Chapter 26, p 24.
- (40) Warcup, J. H. Methods for isolation and estimation of activity of fungi in soil. In *The ecology of Soil Fungi*; Parkinson, D., Waid, J. S., Eds.; Liverpool University Press: Liverpool, 1960; pp 3–21.
- (41) Dox, A. W. Intracellular enzymes of lower fungi, especially those of *Penicillium camemberti*. *J. Biol. Chem.* **1909**, *6*, 461–467.
- (42) Pitt, J. I. An appraisal of identification methods for *Penicillium species*—Novel taxonomic criteria based on temperature and water relations. *Mycologia* **1973**, *65*, 1135–1157.
- (43) Gordon, W. L. The occurrence of *Fusarium* species in Canada. 2. Prevalence and taxonomy of *Fusarium* species in cereal seed. *Can. J. Bot.* **1952**, *30*, 209–251.
- (44) Zycha, H.; Siepmann, R. *Mucorales*; Cramer: 1969; 355 pp.
- (45) Booth, C. *The genus Fusarium*; Commonwealth Mycological Institute: Kew, England, 1966; 237 pp.
- (46) Ellis, M. B. *Dematiaceous Hyphomycetes*; Commonwealth Mycological Institute: Kew, England, 1971; 608 pp.
- (47) Raper, K. B.; Fennell, D. I. In *The genus Aspergillus*; Krigegeger, Ed.; New York, 1973; 626 pp.
- (48) Pitt, J. I. *The genus Penicillium and its teleomorphic states Eupenicillium and Talaromyces*; Academic Press: London, 1979; 634 pp.
- (49) Samson, R. A. A compilation of the *Aspergilli* described since 1965. *Stud. Mycol.* **1979**, *18*, 1–40.
- (50) Al-Musallam, A. Revision of the black *Aspergillus* species. Dissertation, University of Utrecht, The Netherlands, 1980; 92 pp.
- (51) Von Arx, J. A. *The genus of fungi sporulating in pure culture*; Cramer: 1981; 424 pp.
- (52) Bragulat, M. R.; Abarca, M. L.; Accensi, F.; Cabanes, F. J. New screening method for ochratoxigenic molds in pure cultures. *Acta Vet. Scand.* **1998**, *22*, 515.

- (53) Visconti, A.; Pascale, M.; Centonze, G. Determination of ochratoxin A in wine by means of immunoaffinity column cleanup and high-performance liquid chromatography. *J. Chromatogr.* **1999**, *864*, 89–101.
- (54) Abarca, M. L.; Bragulat, M. R.; Castella, G.; Cabanes, F. J. Ochratoxin A production by strains of *Aspergillus niger* var. *niger*. *Appl. Environ. Microbiol.* **1994**, *60*, 2650–2652.
- (55) Frisvad, J. C. The connection between the *Penicillia* and *Aspergilli* and mycotoxins with special emphasis on misidentified isolates. *Arch. Environ. Contam. Toxicol.* **1989**, *18*, 452–467.
- (56) Lillehoj, E. B.; Goransson, B. Occurrence of ochratoxin- and citrinin-producing fungi on developing Danish barley grain. *Acta Pathol. Microbiol. Scand., Sect. B* **1980**, *88*, 133–7.
- (57) Mills, J. T.; Abramson, D. Ochratoxigenic potential of *Penicillium* spp. from stored rapessed and cereals in western Canada. *Can. J. Plant Pathol.* **1982**, *4*, 37–41.
- (58) Mills, J. T.; Abramson, D.; Frohlich, A. A.; Marquardt, R. R. Citrinin and ochratoxin A production by *Penicillium* spp. from stored durum wheat. *Can. J. Plant Pathol.* **1989**, *11*, 357–360.
- (59) Creppy, E. E.; Betbeder, A. M.; Gharbi, A.; Counord, J.; Castegnaro, M.; Bartsch, H.; Montcharmont, P.; Fouillet, B.; Chambon, P.; Dirheimer, G. *Human ochratoxicosis in France*; IARC Scientific Publication No. 115; International Agency for Research of Cancer: Lyon, France, 1991; pp 145–151.
- (60) IPCS (International Program on Chemical Safety). Selected mycotoxins: ochratoxins, trichothecenes, ergot. *Environ. Health Criter.* **1990**, *105*, 27–69.
- (61) Höhler, D. Ochratoxin A in food and feed: occurrence, legislation and mode of action. *Z. Ernährungswiss.* **1998**, *37*, 2–12.
- (62) Ospital, M.; Cazabeil, J.-M.; Betbeder, A.-M.; Tricard, C.; Creppy, E. E.; Medina, B. L'ochratoxine A dans les vins. *Rev. Fr. Oenol.* **1998**, *169*, 16–18.
- (63) Burdaspal, P. A.; Legarda, T. M. Ochratoxina a en vinos, mostos y zumos de uva elaborados en España y en otros países europeos. *Alimentaria Enero-Febrero*, **1999**, 107–113.
- (64) Walker, R. Presented at the 3rd Joint FAO/UNEP International Conference on Mycotoxins, Tunis, Tunisia, March 3–6, 1999.
- (65) Schwenk, S.; Altmayer, B.; Eichhorn, K. W. Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary gas chromatography. *J. Chromatogr.* **1988**, *448*, 424–7.
- (66) Scott, P. M.; Fuleki, T.; Harwig, J. Patulin content of juice and wine produced from moldy grape. *J. Agric. Food Chem.* **1977**, *25*, 434–437.
- (67) Studer-Rohr, I. Ochratoxin A in humans: Exposure, kinetics and risk assessment. Dissertation No. 11071 of the Swiss Federal Institute of Technology (ETH), Zürich, 1995.
- (68) Creppy, E. E.; Castegnaro, M.; Dirheimer, G.; Grosse, Y.; Mériaux, J.; Manier, C.; Montcharmont, P.; Waller, C.; et al. Study of human ochratoxicosis in three French regions: Alsace, Aquitaine and Rhône-Alpes. In *Human ochratoxicosis and its pathologies*; Creppy, E. E., Castegnaro, M., Dirheimer, G. Eds.; Colloque INSERM; INSERM: Paris, France, 1993; Vol. 231, pp 147–158.
- (69) Zimmerli, B.; Dick, R. Determination of ochratoxin A at the ppt level in human blood, serum, milk and some foodstuffs by high-performance liquid chromatography with enhanced fluorescence detection and immunoaffinity column cleanup: methodology and Swiss data. *J. Chromatogr.* **1995**, *666*, 85–99.
- (70) Miraglia, M.; de Dominicis, A.; Brera, C.; Corneli, S.; Cavas, E.; Menghetti, E.; Miraglia, E. Ochratoxin A levels in human milk and related food samples: an exposure assessment. *Nat. Toxins* **1995**, *3*, 436–444.
- (71) Petzinger, E.; Ziegler, K. Ochratoxin A from a toxicological perspective. *J. Vet. Pharmacol. Ther.* **2000**, *23*, 91–8.
- (72) WHO (World Health Organization). *Evaluation of certain food additives and contaminants*; Thirty-seventh report of the joint FAO/WHO Expert Committee on Food Additives (WHO technical report series 806); World Health Organization: Geneva, 1991; pp 29–31.
- (73) Anonymous. Opinion on ochratoxin A. European Commission DG XXIV, CS/CNTM/MYC14, 1998.

---

Received for review July 31, 2001. Revised manuscript received November 21, 2001. Accepted November 26, 2001.

JF011015Z