

Puerins A and B, Two New 8-C Substituted Flavan-3-ols from  
Pu-er Tea

ZHI-HONG ZHOU, YING-JUN ZHANG,\* MIN XU, AND CHONG-REN YANG

State Key Laboratory of Phytochemistry and Plant Resources of West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, People's Republic of China

Pu-er tea is a special treated fermented tea produced from crude green tea, which is prepared from the leaves of *Camellia sinensis* var. *assamica*. It is a traditional beverage having been used in China, particularly the southern areas, for a long time. Chemical investigation led to the identification of two new 8-C substituted flavan-3-ols, puerins A (**1**) and B (**2**), and two known cinchonain-type phenols, epicatechin-[7,8-*bc*]-4 $\alpha$ -(4-hydroxyphenyl)-dihydro-2(3*H*)-pyranone (**3**) and cinchonain Ib (**4**), together with other seven known phenolic compounds, 2,2',6,6'-tetrahydroxydiphenyl (**5**), (-)-epicatechin-8-C- $\beta$ -D-glucopyranoside (**6**), (-)-epicatechin (EC) (**7**), (-)-epigallocatechin (EGC) (**8**), ( $\pm$ )-gallocatechin (GC) (**9**), gallic acid (**10**), and myricetin (**11**), in addition to caffeine (**12**). Their structures were determined by spectroscopic methods. The compounds **1–5**, which might be formed in the postfermentative process of Pu-er tea, were isolated from tea and theaceous plants for the first time.

**KEYWORDS:** Pu-er tea; *Camellia sinensis* var. *assamica*; phenolic compounds

## INTRODUCTION

Tea is one of the most popular beverages consumed in the world, and its biological activities and health functions have been widely explored. It is produced from the leaves of tea plants, *Camellia sinensis* L. and *C. sinensis* var. *assamica* (Masters) Kitamura (Theaceae), which are evergreen shrubs or trees and usually pruned to 2–5 feet for cultivation. On the basis of the processing procedures, tea can generally be divided into green tea (nonfermented), oolong tea (semifermented), and black tea (fully fermented by oxidizing enzyme). Up to now, many chemical investigations have been carried out on tea, and a series of flavan-3-ols and hydrolyzable tannins in green tea (**1**, **2**), 8-C-ascorbyl(-)-epigallocatechin 3-*O*-gallate and novel flavan-3-ols derivatives in oolong tea (**3–5**), and benzotropolone-type pigments in black tea (**6**, **7**), were reported.

Pu-er tea, a well-known tea from ancient times, was originally produced in the Yunnan province of China through a special postfermentative process, using crude green tea as raw materials, which were prepared from the leaves of *C. sinensis* var. *assamica*. It is increasingly liked by consumers not only because of the special flavor and taste but also the health-giving biological function. Pu-er tea was initially formed from long-distance transport of crude green tea on horseback. However, now, it is produced by a standardized postfermentative process with high temperatures and high humidity. The fresh leaves of *C. sinensis* var. *assamica*, the original plant material of Pu-er tea, were found to contain chalcane-flavan dimers and flavan-3-ol derivatives (**8**), and some flavonols and catechins were

reported from the raw material (the crude green tea) of Pu-er tea by our group (**9**). However, the chemical constituents of Pu-er tea are thus far not known. In this investigation, we isolated a number of flavanoids and phenolic compounds from Pu-er tea and determined their structures by spectroscopic methods.

## MATERIALS AND METHODS

**General.** Optical rotations were measured on a P-1020 Polarimeter (JASCO, Tokyo, Japan). IR spectra were measured on an IR-450 spectrometer (Shimadzu, Kyoto, Japan) with KBr pellets. UV spectra were obtained on a 210A double-beam spectrophotometer (Shimadzu, Kyoto, Japan).  $^1\text{H}$  and  $^{13}\text{C}$  NMR,  $^1\text{H}$ - $^1\text{H}$  COSY, HMQC, and HMBC spectra were recorded in acetone- $d_6$  with Bruker AM-400 and DRX-500 spectrometers operating at 500 and 400 MHz for  $^1\text{H}$  and 125 and 100 MHz for  $^{13}\text{C}$ , respectively. Coupling constants are expressed in Hertz, and chemical shifts are given on a  $\delta$  (ppm) scale with tetramethylsilane as an internal standard. FABMS and HRFABMS were recorded on an AutoSpe 3000 spectrometer (VG, Manchester, U.K.) with glycerol as the matrix. Column chromatography was done on Diaion 101 resin (Shandong Lukang Pharmaceutical Co., Ltd.), Sephadex LH-20, 25–100  $\mu\text{m}$  (Pharmacia Fine Chemical Co., Ltd.), MCI-gel CHP20P, 75–100  $\mu\text{m}$  (Mitsubishi Chemical Co., Ltd.), and Toyopearl HW-40F (Tosoh Co., Ltd.). Thin-layer chromatography (TLC) was performed on precoated kieselgel 60 F254 plates, 0.2 mm thick (Merck), with benzene/ethyl formate/formic acid (3:6:1 or 2:7:1, v/v), and spots were detected by spraying with a 2% ethanolic  $\text{FeCl}_3$  or anisaldehyde- $\text{H}_2\text{SO}_4$  reagent, followed by heating.

\* To whom correspondence should be addressed. Telephone: +86-871-5223235. Fax: +86-871-5150124. E-mail: zhangyj@mail.kib.ac.cn.

**Table 1.**  $^{13}\text{C}$  NMR Spectroscopic Data of Compounds **1** and **2** ( $\delta$  Value, in Acetone- $d_6$ )

C	1	2	C	1	2
2	79.4	80.2	13	31.9	32.6
3	66.5	67.2	14	23.8	24.4
4	29.2	29.5	15	29.2	29.5
5	155.3	157.6	16	35.7	36.4
6	95.8	96.1	17	12.5	12.6
7	156.7	157.6	1'	131.5	131.4
8	100.9	100.4	2'	114.7	106.9
9	156.3	156.0	3'	145.2	146.7
10	104.9	105.0	4'	145.2	135.0
11	53.1	54.0	5'	118.7	146.7
12	176.0	177.4	6'	115.7	106.9

**Plant Materials.** Pu-er tea, produced from the crude green tea prepared with leaves of *Camellia sinensis* var. *assamica*, was purchased at Linchang Tea Factory, Linchang County, Yunnan Province of China, and used for extraction and isolation of the chemical constituents.


**Extraction and Isolation.** Pu-er tea (1 kg) was extracted with 80% aqueous acetone at room temperature for 3 times (6, 3, and 3 L). After removal of the acetone under reduced pressure, the aqueous solution afforded precipitates, which were removed by filtration. The filtrate was extracted with  $\text{CHCl}_3$  to give the crude compound **12** (8 g). The aqueous mother liquor was concentrated under reduced pressure to afford a residue (49 g), which was applied to a Diaion 101 resin column, eluting with  $\text{H}_2\text{O}$  and MeOH (each 1 L), successively. The MeOH eluate was concentrated to dryness and subjected to column chromatography over Sephadex LH-20, eluting with MeOH– $\text{H}_2\text{O}$  (1:0–0:1) (each gradient at 300 mL) to afford five fractions. Further repeated column chromatography over MCI-gel CHP20P, Toyopearl HW-40F, and Sephadex LH-20 gave **3** (35 mg), **4** (40 mg), and **10** (60 mg) from fraction 1, **1** (40 mg), **2** (25 mg), and **7** (60 mg) from fraction 2, **6** (45 mg) and **11** (20 mg) from fraction 3, **8** (22 mg) and **9** (19 mg) from fraction 4, and **5** (80 mg) from fraction 5, respectively.

**Puerin A (1).** Pale brown amorphous powder,  $[\alpha]_D^{25} +10.0$  ( $c$  0.005, MeOH). UV  $\lambda_{\text{max}}$  (MeOH): 207.5, 225.5, 282 nm. IR (KBr)  $\nu_{\text{max}}$  ( $\text{cm}^{-1}$ ): 3279, 1611, 1456, 1355, 1283, 1205, 1161, 1101, 1066, 975, 932.  $^1\text{H}$  NMR (400 MHz, acetone- $d_6$ )  $\delta$ : 6.80 (1H, br s, H-2'), 6.76 (1H, d,  $J = 8.0$  Hz, H-5'), 6.67 (1H, br d,  $J = 8.0$  Hz, H-6'), 6.13 (1H, s, H-6), 5.36 (1H, br s, H-11), 4.80 (1H, br s, H-2), 4.20 (1H, m, H-3), 3.57, 2.52 (each 1H, m, H-13), 2.87 (1H, br d,  $J = 16.5$  Hz, H-4b), 2.79 (2H, m, H-15), 2.72 (1H, br d,  $J = 16.5$  Hz, H-4a), 2.38, 2.12 (each 1H, m, H-14), 2.21 (2H, m, H-16), 1.00 (3H, br s, H-17).  $^{13}\text{C}$  NMR (100 MHz, acetone- $d_6$ ): see **Table 1**. FABMS (negative-mode):  $m/z$  400  $[\text{M}]^-$ . HRFABMS  $m/z$  400.1514  $[\text{M}]^-$ , calcd for  $\text{C}_{22}\text{H}_{24}\text{O}_7$ , 400.1522.

**Puerin B (2).** Pale brown amorphous powder,  $[\alpha]_D^{25} +10.0$  ( $c$  0.005, MeOH).  $^1\text{H}$  NMR (400 MHz, acetone- $d_6$ )  $\delta$ : 7.01 (2H, s, H-2', 6'), 6.13 (1H, s, H-6), 5.38 (1H, br s, H-11), 4.76 (1H, br s, H-2), 4.22 (1H, m, H-3), 3.60, 2.52 (each 1H, m, H-13), 2.85 (1H, br d,  $J = 16.5$  Hz, H-4b), 2.78 (2H, m, H-15), 2.72 (1H, br d,  $J = 16.5$  Hz, H-4a), 2.37, 2.12 (each 1H, m, H-14), 2.22 (2H, m, H-16), 1.09 (3H, br s, H-17).  $^{13}\text{C}$  NMR (100 MHz, acetone- $d_6$ ): see **Table 1**. FABMS (negative mode):  $m/z$  415  $[\text{M}-\text{H}]^-$ . HRFABMS:  $m/z$  415.1553  $[\text{M}-\text{H}]^-$ , calcd for  $\text{C}_{22}\text{H}_{23}\text{O}_8$ , 415.1393.

## RESULTS AND DISCUSSION

The 80% aqueous acetone extract of commercial Pu-er tea was separated successively by partitioning with  $\text{CHCl}_3$  and  $\text{H}_2\text{O}$ ,

**Figure 1.** Structures of compounds isolated from Pu-er tea.

and repeated Diaion 101 resin, Sephadex LH-20, MCI-gel CHP20P, and Toyopearl HW-40F column chromatography afforded two new phenolic compounds **1** and **2** (**Figure 1**). In addition, 10 known compounds were also obtained and identified as epicatechin-[7,8-*bc*]-4 $\alpha$ -(4-hydroxyphenyl)-dihydro-2-(3*H*)-pyranone (**3**) (*10*), cinchonain Ib (**4**) (*11*, *12*), 2,6,2',6'-tetrahydroxybiphenyl (**5**) (*13*), (–)-epicatechin-8-*C*- $\beta$ -D-glucopyranoside (**6**) (*14*), (–)-epicatechin (EC) (**7**) (*15*), (–)-epigallocatechin (EGC) (**8**) (*16*), ( $\pm$ )-gallocatechin (GC) (**9**) (*17*), gallic acid (**10**), myricetin (**11**) (*15*), and caffeine (**12**), respectively, on the basis of their physical and spectroscopic data and by comparison with reference values and authentic samples.

Compound **1** was obtained as a pale brown amorphous powder and showed UV absorptions at 207.5, 225.5, 282 nm. The negative HRFABMS exhibited a molecular ion peak at  $m/z$  400.1514  $[\text{M}]^-$ , corresponding to the molecular formula  $\text{C}_{22}\text{H}_{24}\text{O}_7$ . The occurrence of a flavan-3-ol skeleton in the molecule could be easily deduced from the  $^1\text{H}$  NMR spectrum. The signals at  $\delta$  4.80 (s), 4.20 (m), and 2.70–2.85 (2H, m) ascribable to  $\text{C}_2$ ,  $\text{C}_3$ , and  $\text{C}_4$  protons on a flavan C ring, respectively, and ABX-type aromatic signals at  $\delta$  6.80 (1H, br s, H-2'), 6.76 (1H, br d,  $J = 8.0$  Hz, H-5'), and 6.67 (1H, d,  $J = 8.0$  Hz, H-6') arising from the 3',4'-dihydroxylated B-ring protons were closely related to those in (–)-epicatechin (**7**) (*15*). A singlet signal at high field in the aromatic region ( $\delta$  6.13) suggested that the A ring is pentasubstituted. The  $^{13}\text{C}$  NMR and DEPT spectra (**Table 1**) revealed, in addition to 15 signals similar to those of epicatechin, compounds **3** (*10*) and **4** (*11*), the presence of a methine [ $\delta_{\text{C}}$  53.1 (d),  $\delta_{\text{H}}$  5.36] and *n*-pentyl [ $\delta$  23.8 (t), 29.2 (t), 31.9 (t), 35.7 (t), 12.5 (q)] groups, which were confirmed to be linked to each other by the  $^1\text{H}$ - $^1\text{H}$  COSY spectra. Furthermore, the




Figure 2. Important  $^1\text{H}$ - $^1\text{H}$  COSY and HMBC correlations of **1**.

appearance of a signal at  $\delta$  176.0 suggested the presence of a carboxyl function. This was supported by the IR spectrum showing a strong band at  $1611\text{ cm}^{-1}$ , probably because of a  $\beta,\gamma$ -unsaturated five-membered lactone. In the HMBC spectrum of **1** (Figure 2), the methylene protons at  $\delta$  2.87 and 2.72 (H-4) were correlated with the carbon signal at  $\delta$  155.3, which was shown to correlate with the aromatic proton at  $\delta$  6.13, assigning the carbon signal at  $\delta$  155.3 as C-5 and the aromatic proton ( $\delta$  6.13) as C<sub>6</sub>-H. In addition, HMBC correlation of the methine proton at  $\delta$  5.36 (H-11) with  $\delta$  156.3 (C-9) indicated the occurrence of a carbon-carbon linkage at the C<sub>8</sub> position in the epicatechin moiety. Other HMBC correlations including the methine at  $\delta$  5.36 (H-11) with the carboxyl group at  $\delta$  176.0 were also observed (Figure 2). On the basis of the above evidence, the five-membered lactone ring with an *n*-pentyl group attached was determined to be located at C-7 and C-8 positions of the epicatechin moiety and the structure of puerin A was established as shown in molecular **1**.

The molecular formula of compound **2** was determined to be C<sub>22</sub>H<sub>24</sub>O<sub>8</sub> based on the negative HRFABMS, which had one more oxygen atom than that of **1**. The  $^1\text{H}$  and  $^{13}\text{C}$  (Table 1) NMR spectroscopic features were consistent with those of **1**, suggesting a similar structure of **2** to that of **1**. The only difference was observed for their B ring in flavan-3-ol moiety. The typical  $^1\text{H}$  and  $^{13}\text{C}$  chemical shifts [ $\delta_{\text{H}}$  7.01 (2H, s),  $\delta_{\text{C}}$  131.4 (C-1'), 106.9 (C-2', 6'), 146.7 (C-3', 5') and 135.0 (C-4')] indicated the presence of a pyrogallol moiety as the B ring in **2**. Therefore, the structure of **2** was determined to be that of puerin B.

The configurations of C-11 methine in **1** and **2** were deduced by comparing their  $[\alpha]_{\text{D}}$  with those of the known structural-related complex flavan-3-ols, which have the same substituent location in the flavan A ring and similar structures to **1** and **2**. Both epicatechin- (cinchonain Ia) and catechin-[7,8-*bc*]-4 $\beta$ -(3,4-dihydroxyphenyl)-dihydro-2(3*H*)-pyranone with a different C-3 configuration of the flavan-3-ol moiety but the same 4 $\beta$  configuration of the substituent in the flavan A ring showed the same negative  $[\alpha]_{\text{D}}$ , while epicatechin- (cinchonain Ib, **4**) and catechin-[7,8-*bc*]-4 $\alpha$ -(3,4-dihydroxyphenyl)-dihydro-2(3*H*)-pyranone with the same 4 $\alpha$  configuration of the substituent also gave the same positive  $[\alpha]_{\text{D}}$  (11, 12). This observation revealed that the positive or negative symbol of the  $[\alpha]_{\text{D}}$  of these complex flavan 3-ols was determined by the substituent configuration (4 $\alpha$  or 4 $\beta$ ) in the flavan A ring. Accordingly, the configurations of the C-11 methine in **1** and **2** were deduced to be  $\alpha$  because of their positive  $[\alpha]_{\text{D}}$  values.

It is noticeable that the chemical constituents of Pu-er tea are distinctly different from those of the other kinds of tea, e.g., green tea, oolong tea, and black tea. Most of the characteristic flavan-3-ol derivatives in green tea, such as epigallocatechin 3-*O*-gallate (EGCG), EGC, and epicatechin 3-*O*-gallate (ECG), have disappeared, whereas the content of polyphenol polymers is remarkably enhanced in Pu-er tea. At the same time, several polyoxygenated compounds such as **1**–**5** are formed from the

related catechin derivatives during the processing procedure of Pu-er tea. Compounds **1** and **2** are the first acylated flavan-3-ols with an aliphatic acid moiety linked to the flavan-3-ol A ring as a five-membered lactone through a C–C linkage. Both of the compounds **3** and **4** belonging to cinchonain-type flavan-3-ols are isolated from tea and tea plants for the first time. The former (**3**) is a phenylpropanoid-catechin isolated from the bark of *Ocotea porosa* (Lauraceae) (10), while the latter (**4**) was reported to be isolated from *Cinchona succirubra* (Rubiaceae) (11). Compound **5**, known as a synthetic chemical, is obtained from a natural source for the first time (13).

#### ACKNOWLEDGMENT

The authors are grateful to the members of the Analytical Group of our Laboratory for measurements of all spectra.

#### LITERATURE CITED

- Nonaka, G.; Kawahara, O.; Nishioka, I. Tannins and related compounds. XV. A new class of dimeric flavan-3-ol gallates, theasinensins A and B, and proanthocyanidin gallates from green tea leaf. *Chem. Pharm. Bull.* **1983**, *31*, 3906–3914.
- Nonaka, G.; Sakai, R.; Nishioka, I. Hydrolyzable tannins and proanthocyanidins from green tea. *Phytochemistry* **1984**, *23*, 1753–1755.
- Hashimoto, F.; Nonaka, G.; Nishioka, I. Tannins and related compounds. LVI. Isolation of four new acylated flavan-3-ols from oolong tea (**1**). *Chem. Pharm. Bull.* **1987**, *35*, 611–616.
- Hashimoto, F.; Nonaka, G.; Nishioka, I. Tannins and related compounds. LXIX. Isolation and structure elucidation of B, B'-linked bisflavanoids, theasinensins D-G and oolongtheanin from oolong tea (**2**). *Chem. Pharm. Bull.* **1988**, *36*, 1676–1684.
- Hashimoto, F.; Nonaka, G.; Nishioka, I. Tannins and related compounds. XC. 8-*C*-ascorbyl (-)-epigallocatechin 3-*O*-gallate and novel dimeric flavan-3-ols, oolonghomobisflavans A and B, from oolong tea (**3**). *Chem. Pharm. Bull.* **1989**, *37*, 3255–3263.
- Nonaka, G.; Hashimoto, F.; Nishioka, I. Tannins and related compounds. XXXVI. Isolation and structures of theaflagallins, new red pigments from black tea. *Chem. Pharm. Bull.* **1986**, *34*, 61–65.
- Hashimoto, F.; Nonaka, G.; Nishioka, I. Tannins and related compounds. CXIV. Structure of novel fermentation products, theogallinin, theaflavinonin, and desgalloyl theaflavinonin from black tea, and changes of tea leaf polyphenols during fermentation. *Chem. Pharm. Bull.* **1992**, *40*, 1383–1389.
- Nonaka, G.; Hashimoto, F.; Nishioka, I. Tannins and related compounds. LXVII. Novel chalcone-flavan dimmers, assamicains A, B, and C, and a new flavan-3-ol and proanthocyanidins from the fresh leaves of *Camellia sinensis* L. var. *assamica* Kitamura. *Chem. Pharm. Bull.* **1989**, *37*, 77–85.
- Zhou, Z.-H.; Yang, C.-R. Chemical constituents of crude green tea, the material of Pu-er tea in Yunnan. *Acta Bot. Yunnanica* **2000**, *22*, 343–350.
- David, J. M.; Yoshida, M.; Gottlieb, O. R. Phenylpropanoid-catechins from bark of *Ocotea porosa*. *Phytochemistry* **1994**, *35*, 545–546.
- Nonaka, G. I.; Nishioka, I. Tannins and related compounds. VII. Phenylpropanoid-substituted epicatechins, cinchonains from *Cinchona succirubra* (**1**). *Chem. Pharm. Bull.* **1982**, *30*, 4268–4276.
- Chen, H. F.; Tanaka, T.; Nonaka, G. I.; Fujioka, T.; Mihashi, K. Phenylpropanoid-substituted catechins from *Castanopsis hystrix* and structure revision of cinchonains. *Phytochemistry* **1993**, *33*, 183–187.
- Harada, T.; Tuyet, T. M. T.; Oku, A. Asymmetric desymmetrization of 2,2',6,6'-tetrahydroxybiphenyl through annulation with enantiomerically pure bis(mesyate). *Org. Lett.* **2000**, *2*, 1319–1322.

- (14) Kashiwada, Y.; Nishioka, I. Tannins and related compounds. XLV. Rhubarb (5). Isolation and characterization of flavan-3-ol and procyanidin glucosides. *Chem. Pharm. Bull.* **1986**, *34*, 3208–3222.
- (15) Zhang, W.-J.; Liu, Y.-Q.; Li, X.-C.; Yang, C.-R. Chemical constituents of “Ecological Tea” from Yunnan. *Acta Bot. Yunnanica* **1995**, *17*, 204–208.
- (16) Nonaka, G.; Nishioka, I.; Nagasawa, T.; et al. Tannins and related compounds. I. Rhubarb (1). *Chem. Pharm. Bull.* **1981**, *29*, 2862–2870.
- (17) Nonaka, G.; Muta, M.; Nishioka, I. Myricatin, a galloyl flavanone sulfate and prodelphinidin gallates from *Myrica rubra*. *Phytochemistry* **1983**, *22*, 237–241.

---

Received for review June 12, 2005. Revised manuscript received August 22, 2005. Accepted August 22, 2005.

JF051390H