

Antioxidant Activity and Bioactive Compounds of Tea Seed (*Camellia oleifera* Abel.) Oil

CHIA-PU LEE AND GOW-CHIN YEN*

Department of Food Science and Biotechnology, National Chung Hsing University,
 250 Kuokuang Road, Taichung 40227, Taiwan, Republic of China

The oil of tea seed (*Camellia oleifera* Abel.) is used extensively in China as cooking oil. The objectives of this study were to investigate the antioxidant activity of tea seed oil and its active compounds. Of the five solvent extracts, methanol extract of tea seed oil exhibited the highest yield and the strongest antioxidant activity as determined by DPPH scavenging activity and Trolox equivalent antioxidant capacity (TEAC). Two peaks separated from the methanol extract by HPLC contributed the most significant antioxidant activity. These two peaks were further identified as sesamin and a novel compound: 2,5-bis-benzo[1,3]dioxol-5-yl-tetrahydro-furo [3,4-*d*][1,3]dioxine (named compound B) by UV absorption and characterized by MS, IR, ¹H NMR, and ¹³C NMR techniques. Sesamin and compound B decreased H₂O₂-mediated formation of reactive oxygen species in red blood cells (RBCs), inhibited RBCs hemolysis induced by AAPH, and increased the lag time of conjugated dienes formation in human low-density lipoprotein. The results indicate that both compounds isolated from tea seed oil exhibit remarkable antioxidant activity. Apart from the traditional pharmacological effects of *Camellia oleifera*, the oil of tea seed may also act as a prophylactic agent to prevent free radical related diseases.

KEYWORDS: Tea seed oil; antioxidant activity; sesamin; LDL; oxidative damage; comet assay

INTRODUCTION

The oil of tea seed (*Camellia oleifera* Abel.) is used extensively in China as cooking oil (1). It was also traditionally applied as a medicine for stomachache and burning injury in China. The seeds of plants usually contain various antioxidant factors; the naturally occurring active components in tea seed are different from other kinds of oil seeds. Chen et al. (2) indicated that the saponin in tea seed could lower the cholesterol, triglycerides, and low density-lipoprotein in the blood of rat. Zhang and Zhou (3) found that it has the antioxidant ability to reduce liver reactive oxygen species (ROS) in rats. Some oils, such as olive oil, are rich in health beneficial compounds, except they do not supply energy for humans. The Mediterranean diet could affect several degenerative pathologies, including coronary heart diseases and cancer. In epidemiological and clinical researches, the phenolic compounds of olive oil play an antioxidative role in health (4). Phenolics and flavonoids are major materials of non-nutritive compounds and possess antioxidant, antimutagenic, antiinflammatory, and anticarcinogenic activities, and reduce atherosclerosis (5). There are many kinds of phenolic compounds in olive oil (6). Hydroxytyrosol and tyrosol, which are major bioactive phenolic compounds of olive oil, appear to account for the important part of this protective effect in vivo (7). Several reports have demonstrated that hydroxytyrosol and tyrosol have remarkably protective effects against oxidative stress-related damages. Ramirez-

Tortosa et al. (8) showed that olive oil might efficaciously counteract chemical oxidation of low-density lipoproteins. The major fatty acids of tea seed oil, oleic and linoleic acid, are very similar to those in olive oil (9). We suggest that some non-nutritive constituents in tea seed oil may provide the same effect as olive oil. Therefore, the purposes of this study were to investigate the antioxidant activity and to identify the major antioxidant compounds of tea seed oil, and to further characterize the antioxidant properties of these compounds and their effect on health.

MATERIALS AND METHODS

Materials and Chemicals. The sample of tea seed (*Camellia oleifera* Abel.) was supplied from the Hsin-I country farmer's association (Nantou, Taiwan). The tea seed was sealed in a plastic bag and stored at -20 °C until use. Methanol, ethanol, acetone, ethyl acetic, and acetonitrile (HPLC grade) were obtained from Tedia Co. (Fairfield, OH). 2,2'-Azobis-(2-amidinopropane)-dihydropropane (AAPH), copper sulfate pentahydrate (CuSO₄·5H₂O), sodium dihydrogen phosphate, disodium hydrogen phosphate, potassium bromide, and ethylenediaminetetraacetic acid disodium salt dihydrate (EDTA) were obtained from Wako Pure Chemical Co. (Osaka, Japan). Sodium bromide and sodium chloride were obtained from Merck Co. (Darmstadt, Germany). α,α-Diphenyl-β-picrylhydrazyl (DPPH), 2',7'-dichlorofluorescein diacetate (DCFH-DA), 2,2'-azinobis-(3-ethylbenzthiazoline-6-sulfonate) (ABTS), 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), dimethyl sulfoxide (DMSO), Triton X-100, hydrogen peroxide, Trypan blue, ethidium bromide, and phenol-red and horseradish peroxidase (HRPase, type II) were obtained from Sigma Co (St. Louis, MO). Nycoprep separation media (density 1.077 Kg/L) was purchased

* Author to whom correspondence should be addressed [telephone +886-4-22879755; fax +886-4-22854378; e-mail geyen@nchu.edu.tw].

from Pharmacia Biotech Co. (Sweden). Hydrogen peroxide was purchased from Hayasahi Co. (Tokyo, Japan).

Preparation of Tea Seed Oil and Extracts. Tea seeds were roasted at 120 °C for 20 min and then pressed by twin screw extruder to obtain the tea seed oil. Tea seed oil (100 g) was extracted with 200 mL each of methanol, ethanol, acetone, ethyl acetic, and acetonitrile, respectively, for 1 h in a shaking incubator at room temperature. The extracts were filtered, and the residue was re-extracted three times under the same conditions. The combined filtrates from each solvent were evaporated under vacuum using a rotary evaporator at 40 °C, and then weighed to determine the extraction yield and stored at -20 °C until use.

DPPH Scavenging Effect of Extracts. The scavenging effect of different solvent extracts of tea seed oil on DPPH radical was estimated according to the method of Duh et al. (10). The extracts were added to a methanolic solution (0.5 mL) of DPPH radical (final concentration of DPPH was 0.2 mmol/L). The mixture was shaken vigorously and kept at room temperature for 30 min; the absorbance of the resulting solution was then measured spectrophotometrically at 517 nm.

Total Antioxidant Activity Assay (Trolox Equivalent Antioxidant Capacity, TEAC Assay). The total antioxidant activity of tea seed oil extracts was measured using the TEAC assay as described by Yeh and Yen (11). ABTS⁺ radical cation was generated by the interaction of ABTS (100 μM), H₂O₂ (50 μM), and peroxidase (4.4 unit/mL). To measure antioxidant capacity, 0.25 mL of different solvent extracts (0.2 mg/mL) of tea seed oil was mixed with an equal volume of 2,2'-azino-bis[3-ethyl-n-benz-thiazoline-6-sulfonic acid] (ABTS), H₂O₂, peroxidase, and deionized water. Absorbance was monitored at 734 nm for 10 min. The decrease in absorption at 734 nm after the addition of reactant was used to calculate the TEAC value. TEAC value is expressed as millimolar concentration of Trolox solution having the antioxidant equivalent to a 1000 ppm solution of the sample under investigation. The higher TEAC value of the sample means stronger antioxidant ability.

Isolation of Active Compounds from Extracts of Tea Seed Oil. Isolation of active compounds from methanol extract of tea seed oil was performed with a Hitachi liquid chromatograph (Hitachi, Ltd., Tokyo, Japan), consisting of a model L-6200 pump, a Rheodyne model 7125 syringe-loading sample injector, a model D-2000 integrator, and a diode array detector (Hitachi, L-7455 model) set at 280 nm. A 250 × 4 mm i.d., 5 μm, LiChrospher RP-18 (Merck Co., Ltd., Darmstadt, Germany) was used for analysis and at a flow rate of 0.8 mL/min. The elution solvents were (A) H₂O and (B) MeOH. The solvent gradient elution program used was: 75% A for 20 min; then decreasing to 50% A in 40 min; then decreasing to 10% A in 60 min; then decreasing to 0% A in 65 min, and for 20 min, followed by 5 min isocratic wash at 75% A. A monitoring UV detector (L-4200) was set at 280 nm. Two major peaks were obtained by this separation condition. The same elution solvents were employed for collection of both two peaks in the methanol extract using a preparative 250 × 20 mm, i.d., 5 μm, Mightysil RP-18 column (Kanto Chemical Co., Tokyo, Japan) and a flow rate of 4 mL/min. Peaks containing the major compounds were collected and concentrated in a rotary evaporator and crystallized. Both compounds were identified by means of spectrometry.

Spectrometry. The UV-vis absorption spectra of the active compounds in methanol were recorded on a Hitachi U-3000 spectrophotometer. IR data were recorded in potassium bromide (KBr) disks with Bruker Equinox 55 Fourier Transform infrared spectrophotometer (Bruker, Karlsruhe, Germany). The mass spectra of the components were obtained using the EI-MS mode at 70 eV with a Finnigan/Thermo Quest MAT mass spectrometer (Scientific Instrument Services, Inc., Ringoes, NJ). The temperature was raised by steps of 128 °C/min from 100 to 300 °C.

The nuclear magnetic resonance (NMR) spectra were measured in CDCl₃ and methanol-*d*₄ with a Varian VXR-600S FT-NMR spectrometer (Varian Inc., Walnut Creek, CA) operating at 599.95 MHz for ¹H NMR and 75.43 MHz for ¹³C NMR with complete proton decoupling. The sweep width, pulse angle, repetition delay, and acquisition time for ¹H NMR were 5500.0 Hz, 7.0 μs, and 0 s, respectively, and for ¹³C NMR were 22 000.0 Hz, 7.0 μs, and 2.0 s, respectively. The chemical

shifts are reported in parts per million (ppm) from tetramethylsilane. Melting points were determined on a Buchi 535 apparatus (Haverhill, MA).

The IR spectrum of compound A showed absorptions at 1493.9, 1243.9, 1038.9, and 927.8 cm⁻¹. The FAB-MS of compound A gave an M⁺ ion peak at 354, suggesting a molecular weight of 354.4. The ¹H NMR spectrum showed the following chemical shifts: δ 6.78 (1H, d, *J* = 7.8 Hz), 6.84 (1H, dd, *J* = 1.8 and 7.8 Hz), and 6.88 (1H, d, *J* = 1.8 Hz); δ 3.09 (m), 3.84 (dd, *J* = 4.0 and 9.0 Hz), 4.22 (m), and 4.07 (d, *J* = 4.0 Hz). The ¹³C NMR spectrum revealed 15 peaks with the following chemical shifts: δ 93.70 (C-8), 99.82 (C-6), 106.09 (C-10), 115.91 (C-50), 116.12 (C-30), 123.02 (C-10), 123.98 (C-3), 129.00 (C-60), 131.34 (C-20), 154.25 (C-2), 158.38 (C-9), 159.02 (C-4), 163.87 (C-5), 165.07 (C-7), and 181.59 (C-4).

The IR spectrum of compound B showed absorptions at 1486.0, 1245.6, 1038.1, and 927.6 cm⁻¹. The FAB-MS of compound B gave an M⁺ ion peak at 370, suggesting a molecular weight of 370.3. HRMS, *m/z* 370.355 (calculated for C₂₀H₁₈O₇: 370.358). The ¹H NMR spectrum showed the following chemical shifts: δ 6.51 (1H, dd, *J* = 2.4 and 8.4 Hz), 6.62 (1H, dd, *J* = 2.4 Hz), and 6.70 (1H, d, *J* = 8.4 Hz); δ 6.79 (1H, d, *J* = 7.8 Hz), 6.86 (1H, dd, *J* = 1.2 and 7.8 Hz), and 6.90 (1H, d, *J* = 1.2 Hz); δ 2.95 (1H, m), 3.62 (1H, dd, *J* = 7.8 and 9.0 Hz), and 4.408 (1H, d, *J* = 9.0 Hz); δ 3.30 (1H, m), 3.98 (1H, d, *J* = 9.0 Hz), and 4.10 (1H, dd, *J* = 6.6 and 9.0 Hz). The ¹³C NMR spectrum revealed 14 peaks with the following chemical shifts: δ 53.98 (C-5), 54.47 (C-1), 70.80 (C-4), 72.25 (C-6), 88.58 (C-2), 101.14 (C-2'), 102.42 (C-7'), 102.52 (C-7''), 107.51 (C-2'), 108.23 (C-8), 108.84 (C-5''), 108.99 (C-5'), 110.39 (C-6'), 120.82 (C-6').

Assay for Hemolysis of Red Blood Cells (RBCs). The method described by Zhu et al. (12) was used in this study. Blood was obtained from healthy volunteers and collected into heparinized tubes. Erythrocytes were separated from plasma and the buffy coat and washed three times with 5 volumes of phosphate buffered saline, pH 7.4. During each wash, the erythrocytes were centrifuged at 3000g for 10 min to obtain the cell pellet. After the last wash, the packed erythrocytes were suspended in 4 volumes of PBS solution. Oxidative hemolysis of erythrocytes was induced by AAPH, a peroxy radical initiator. Addition of AAPH to the suspension of diluted erythrocytes induces the oxidation of membrane lipids and proteins, resulting in hemolysis. Two milliliters of the erythrocyte suspension was mixed with 2 mL of PBS solution containing different concentrations of compounds from tea seed oil. Two milliliters of 200 μM AAPH in PBS was then added to the mixture. The reaction mixture was shaken gently while being incubated at 37 °C for 3 h. After incubation, the reaction mixture was removed, diluted with 8 volumes of PBS, and centrifuged at 3000g for 5 min. The absorbance of the supernatant fraction at 540 nm was recorded in a Hitachi U-3000 spectrophotometer. Percent inhibition was calculated by the equation in ref 13.

Intracellular ROS Measurement. Intracellular ROS was estimated by the method of Amer et al. (14) using 2',7'-dichlorofluorescein diacetate (DCFH-DA) as a fluorescent probe. Erythrocytes were prepared as described above. Cells (10⁶ cells/mL) were collected and resuspended with PBS and seeded in endpdroff and incubated with various concentrations of compounds from tea seed oil for 30 min at 37 °C. Erythrocytes were washed with phosphate buffered saline three times again. Finally, the cell pellets were resuspended with PBS and co-incubated with 25 μM DCFH-DA for 30 min. After reaction, cells were collected and washed once with PBS, and the cells were incubated with or without freshly prepared 50 μM H₂O₂ for 30 min. Erythrocytes were analyzed by a FACScan flow cytometric (Becton-Dickinson Immunocytometry Systems USA, San Jose, CA). Erythrocytes were passed at a rate of about 1000 cells per second, using saline as the sheath fluid. A 488 nm argon laser beam was used for excitation. Green fluorescence of 10 000 gated erythrocytes was then measured using linear amplification.

LDL Isolation from Plasma. Blood from a single healthy volunteer was collected in tubes containing EDTA (1 mg/mL). Plasma was isolated by centrifugation at 3000g for 10 min. LDL was isolated from the plasma by density gradient ultracentrifugation (*d* = 1.006–1.063) at 10 °C with a Hitachi (Himac CS 120 GX) centrifuge working at 100 000g. After its isolation, LDL was dialyzed against 0.012 M

phosphate buffered saline (10 mM, pH 7.4) and freshened with N₂ at 4 °C. A Bio Rad protein assay solution was used for protein determinations, employing bovine serum albumin as the standard.

Detection of Conjugated Dienes. The formation of conjugated dienes was measured by determining the absorbance increase at 234 nm of the solution of LDL (100 µg protein/mL) in PBS incubated with 2.4 mM CuSO₄ in the absence or presence of compounds from tea seed oil. The absorbance was measured every 5 min for 250 min using a Hitachi U-3000 UV-vis spectrophotometer, and the results were expressed as relative absorbance at 234 nm. The duration of the lag phase was defined as time (min) to the intercept of the tangent of the absorbance curve in the propagation phase with baseline.

Effect of Isolated Compounds from Tea Seed Oil on H₂O₂-Induced DNA Damage. To evaluate the effect of isolated compounds from tea seed oil on modulating DNA damage in lymphocytes induced by H₂O₂, the DNA damage was determined using comet assay as described by Yen and Hsieh (15). Human lymphocytes were isolated from fresh whole blood by adding blood to the RPMI 1640 medium, before underlaying it with Histopaque 1077 and centrifuging at 1600 rpm for 10–15 min. Lymphocytes were separated as a pink layer at the top of the Histopaque 1077. The lymphocytes were washed in the RPMI 1640 medium. The cell number and viability (Trypan blue exclusion) were determined with a Neubauer improved haemocytometer before treatment. Human lymphocytes were incubated at a density of 5 × 10⁵/mL, the viability being over 90%.

The cell suspension was incubated with compounds (10–50 µM) and H₂O₂ (50 µM) at 37 °C for 30 min. After treatment with compounds, the cells were centrifuged and resuspended in preheated 1% low melting point agarose. The cell suspension was then added to Dakin fully frosted microscope slides (Menzel-Glaser, Germany), precoated with 1% normal melting agarose. After application of a third layer of 1% low melting point agarose, the slides were immersed in cold lysing solution (2.5 M NaCl, 100 mM EDTA, 10 mM Tris, 1% sodium lauryl sarcosineate, 1% Triton X-100, 10% DMSO, pH 10) for at least 1 h at 4 °C. The microscope slides were then placed in an electrophoresis tank, and the DNA was allowed to unwind for 20 min in freshly prepared alkaline electrophoresis buffer (1 mM EDTA, 0.3 N NaOH, pH 13). Electrophoresis was conducted at 4 °C for 20 min at 25 V and 300 mA. The slides were then neutralized with Tris buffer (0.4 M, pH 7.5) and stained with ethidium bromide. All of the steps were performed under yellow light to prevent additional damage. The slides were observed using a fluorescent microscope attached to a CCD camera connected to a personal computer-based image analysis system (Komet 3.0; Kinetic Imaging Ltd., UK). For each analysis, 50 individual cells were calculated, and, in most cases, three separate experiments were conducted for each series. Single cells were analyzed under the fluorescent microscope as desired. The DNA damage was expressed as % Tail DNA, where % Tail DNA = Tail DNA/(Head DNA + Tail DNA) × 100. A higher % Tail DNA means a higher level of DNA damage.

Statistical Analysis. Data presented are given as mean ± SD. Statistical significance was examined through one-way analysis of variance and Duncan's multiple range tests. Significant differences were accepted at *p* < 0.05.

RESULTS AND DISCUSSION

Antioxidant Activity of Extracts from Tea Seed Oil.

Comparison of antioxidant activity of tea seed oil extracts from various solvents is shown in **Table 1**. The yields of solvents extracts from 100 g of oil were in the order of methanol (1.65%) > ethyl acetate (0.73%) > acetone (0.52%) > acetonitrile (0.16%). At the same concentration of 200 µg/mL, the scavenging effects of different extracts on DDPH radical were in the order of methanol (66.50%) > ethyl acetate (5.92%) > acetone (4.46%) > acetonitrile (0.40%). The similar antioxidant activity was also obtained by the TEAC assay. The TEAC value of methanol extract of tea seed oil at 200 µg/mL was stronger than other extracts; it was the equal of 59 µM Trolox. Of the four solvent extracts, the methanol extract exhibited the highest yield

Table 1. Antioxidant Activity and Yield of Extracts from Tea Seed Oil with Various Solvents^a

solvent	DPPH scavenging effect (%)	TEAC (µM)	yield (%)
methanol	66.50 ± 1.08 a	59.21 ± 4.72 a	1.65 ± 0.22 a
acetone	4.46 ± 1.17 b	10.42 ± 3.25 b	0.52 ± 0.12 b
ethyl acetic	5.92 ± 1.62 b	12.53 ± 3.24 b	0.73 ± 0.23 b
acetonitrile	0.4 ± 0.96 c	2.32 ± 0.75 c	0.16 ± 0.04 c

^a The concentration of different solvents extracts used in antioxidant activity assays was 200 µg/mL. Each value is expressed as mean ± SD (*n* = 3). Values within a column followed by a different letter represent a significant difference at *p* < 0.05.




Figure 1. Typical HPLC profile of methanol extract from tea seed oils. The retention times of peaks A and B are 66.08 and 69.52 min, respectively. Diode array detector set at 280 nm.

and the strongest antioxidant activity; a significant difference (*p* < 0.05) was found in antioxidant activity among the four extracts. Therefore, the methanol extract of tea seed oil was used for the following isolation and identification of active compounds. The result was similar to the other studies on olive oil; methanol was usually used to extract bioactive compounds of olive oil (16).

Isolation of Active Compounds in Tea Seed Oil. To isolate and identify the active compounds, the methanol extract was separated by means of HPLC. The results showed that there were two major peaks (A and B) in the HPLC profile of METSO with retention times of 66 and 69 min, respectively (**Figure 1**). The content of sesamin and compound B in 100 g of tea seed oil was 33.88 and 18.41 mg, respectively. These two samples were collected by means of preparative HPLC. The UV absorption of compounds showed spectra characteristic of lignans: λ_{max} 210 nm in methanol. Compound A was identified as sesamin on the basis of MS, IR, ¹H NMR, and ¹³C NMR data. The spectroscopic properties obtained from compound A were also consistent with those reported in the literature (17). The molecular formula of compound B was C₂₀H₁₈O₇ and named as 2,5-bis-benzo[1,3]dioxol-5-yl-tetrahydro-furo [3,4-*d*][1,3]-dioxine. The compound B is similar to sesamin in structure, only differing in the two unsymmetrical rings of the central skeleton (**Figure 2**).

Sesamin and sesamol are the two major components of sesame oil lignans (18), and both of them have antioxidant and health promoting activities. Nakai et al. (19) noted that the lignans were highly correlated to the antioxidant effect of sesame. According to the chemical structure, the new compound B isolated from tea seed oil is one of the lignans. Among sesamin, sesamol, and sesaminol, sesamin is the least effective antioxidant, but it can be converted into more antioxidative derivatives. During metabolism in liver, sesamin increased the




Figure 2. Structures of sesamin and compound B isolated from the methanol extract of tea seed oil.




Figure 3. Effects of sesamin and compound B on AAPH-induced hemolysis of human erythrocytes. Results are mean \pm SD for $n = 3$. *Significantly different from treated with AAPH alone ($p < 0.05$).

hydroxyl groups on the phenyl structure and increased its antioxidant activity (20). Sesamin could protect oxidation, lower cholesterol, and improve lipid metabolism both in vitro and in vivo tests (21). On the other hand, Miyahara et al. (22) found that the sesamin in unbaked sesame could suppress the growth of lymphoid leukemia Molt 4B cells and induce apoptosis.

Inhibition of Hemolysis of RBCs by Sesamin and Compound B. Red blood cells act as an oxygen carrier. They are always exposed to high oxygen tension; therefore, they are easily subjected to oxidation injury by superoxides, H_2O_2 , and hydroxyl free radicals in metabolites. Mammalian RBCs are often used as a model system to test the efficacy of antioxidants. **Figure 3** shows the inhibitory effect of sesamin and compound B on hemolysis of RBCs induced by AAPH. As the results in **Figure 3** show, the inhibitory effect of sesamin on hemolysis showed no dose-dependence at the tested concentrations. However, compound B showed significant ($p < 0.05$) inhibitory effect at concentrations from 10 to 50 μM . Results also indicated that sesamin and compound B were able to suppress the oxidative injury of RBCs by free radicals up to 49% and 48%, respectively. Manna et al. (23) reported that with olive oil, both in vivo and in vitro, not only the triglycerides but also the phenolic compounds could suppress the hemolysis caused by the oxidative stress. Lipophilic polyphenols, which easily penetrate the cytoplasm of erythrocytes, seem to react with hemoglobin (24). Sesame seeds lignans could increase plasma γ -tocopherol and alter plasma tocopherol ratios in humans and enhanced vitamin E bioactivity (25). Thus, sesamin and compound B isolated from tea seed oil may also bind at cell membranes and protect against damage from free radical attack.

Inhibition of the ROS Production in Intracellular RBCs by Sesamin and Compound B. **Figure 4** shows the inhibition of sesamin and compound B on H_2O_2 -induced ROS in RBCs.




Figure 4. Effect of sesamin and compound B on H_2O_2 -stimulated ROS in human erythrocytes. Results are mean \pm SD for $n = 3$.




Figure 5. Effect of compounds from tea seed oil on Cu^{2+} -mediated conjugated dienes formation in LDL.

Among the test ranges of 10–50 μM , sesamin showed no significant inhibition on ROS formation. In contrast, the inhibitory effect of compound B increased significantly ($p < 0.05$) from 14% to 86% with increasing concentration. Apparently, the effect of compound B is much higher than that of sesamin, even up to 16 times at 50 μM .

Suppressing ROS could reduce the oxidative damage in cells. Hou et al. (26) reported that sesamin could reduce injuries of the PC12 cell through the decrease of ROS production due to the inhibition of mitochondria activated protein kinases. Zhang and Zhou (3) also found that feeding the tea seed oil to Wistar rats could increase the defense effect by increasing the superoxide dismutase activity of serum and liver. ROS could attack membranes of RBCs and induce cell membrane oxidation. In the present study, compound B showed more inhibitory effect than sesamin on H_2O_2 -induced intercellular ROS formation. It is thought that compound B could potentially reduce the oxidative injury by the aerobic metabolism.

Inhibition of LDL Oxidation by Sesamin and Compound B. **Figure 5** shows that the formation of LDL conjugated dienes under the induction of copper ion was inhibited by the functional compounds of tea seed oil. Results indicated that the addition of either sesamin or compound B could retard the LDL oxidation. During pretreatment with the isolated compounds of tea seed oil, oxidation inducing time could increase from 38 to 51–89 min with the increasing concentrations of samples. There was no conjugated dienes formation in the group without copper ion. In this test, compound B was obviously more effective than that of the sesamin. The lag time of compound B was prolonged by 9 and 27 min as compared to those of sesamin at concentrations of 10 and 30 μM , respectively.

It is well known that the Mediterranean diet could reduce the risk of cancer and arteriosclerosis, a major reason being that




Figure 6. Effect of sesamin and compound B on H_2O_2 -induced DNA damage of lymphocytes. Results are presented as mean \pm SD from three separated experiments. *Indicates significant difference from treated with H_2O_2 alone ($p < 0.05$).

bioactive compounds of vegetables and olive oils provide antioxidant effects to protect against the lipid oxidation. In Oriental countries, sesame is considered a healthy food that is rich in sesamin. In the present study, sesamin and compound B could suppress LDL oxidative modification, which is an important marker of arteriosclerosis. Compound B had more protective efficacy than sesamin in the LDL oxidation system. Yamashita et al. (27) found that sesamin could increase the tocotrienol and tocopherol in rat liver, kidney, and blood plasma to increase the antioxidant defensive effect. Sesamin is highly related to the LDL tocopherol content and oxidation inducing time in the presence copper ion (21). In another study, sesamin in vivo could provide more antioxidant effect than in vitro (19). Sesamin and compound B have antioxidant effects in the in vitro system, and we speculate that they may exhibit potential biological and pharmacological activities.

Inhibition of Genotoxic Injury in Human Lymphocytes by Sesamin and Compound B. H_2O_2 could produce free radicals during metabolism and then attack cell DNA. Figure 6 shows the inhibitory effect of sesamin and compound B on H_2O_2 -induced DNA damage of lymphocytes. The oxidative DNA damage of lymphocytes induced by H_2O_2 was effectively suppressed by these two compounds from tea seed oil. At concentrations of 10–50 μM , the % tail DNA of sesamin and compound B treated groups was significantly different ($p < 0.05$) from the control group. Many studies showed that antioxidants could reduce cell and DNA damage from free radicals. Some antioxidants, such as quercetin and caffeic acid, showed efficacious protection using the comet assay for DNA damage (28). A vegetarian diet can prove to have an antioxidant effect, leading to a slight decrease in oxidative DNA damage in lymphocytes (29). Many studies indicated that sesamin did not provide a strong antioxidant effect, but sesamin could inhibit lipid peroxidation in the microsomal enzyme system due to its methylenedioxy groups. Sesamin could be metabolized to mono-catechol and di-catechol metabolites by liver enzymes, and changed into more antioxidant metabolites in liver (19). Sesamin can be absorbed from the intestines and reaches the liver via the portal vein where it is metabolized to catechol and enhances antioxidation. Both metabolites had antioxidant activity that means sesamin could be a pro-antioxidant (20). Sesamin also could be absorbed and affected hepatic fatty acid oxidation and synthesis (30). Furthermore, sesamin had antioxidant and anti-inflammatory effects on hepatic ischemia-reperfusion injury in rat animal testing (31). Compound B is similar to sesamin in catechol moiety structure; it may be metabolized to contain catechol structure by liver.

In the present study, our results demonstrated that methanol extract from tea seed oil showed strong DPPH scavenging effect and antioxidant ability. There are two major constituents of lignans, sesamin and compound B, isolated from tea seed oil. By using in vitro experiments, both of the compounds could reduce the oxidative DNA damage. Apart from the traditional pharmacological effects of *Camellia oleifera*, the oil of tea seed may also act as a prophylactic agent to prevent free radical related diseases.

LITERATURE CITED

- (1) Yu, Y. S.; Ren, S. X.; Tan, K. Y. Study on climatic regionalization and layer and belt distribution of *oiltea ccamellia* quality in china. *J. Nat. Res.* **1999**, *14*, 123–127.
- (2) Chen, L. F.; Qiu, S. H.; Peng, Z. H. Effects of sasanguasaponin on blood lipids and subgroups of high-density lipoprotein cholesterol in hyper lipidemia rat models. *Pharm. Clin. Chin. Mater. Med.* **1998**, *14*, 13–16.
- (3) Zhang, B.; Zhou, Y. Effects of tea seed oil and soybean oil on oxygen radical and activities of antioxidative enzymes in rat. *Acta Nutr. Sin.* **1995**, *17*, 199–203.
- (4) Saija, A.; Uccella, N. Olive biophenols functional effects on human wellbeing. *Trends Food Sci. Technol.* **2001**, *11*, 357–363.
- (5) Huang, M. T.; Ferraro, T. Phenolic compounds in food and cancer prevention. In *Phenolic Compounds in Food and their Effects on Health*; Ho, C. T., Lee, C. Y., Huang, M. T., Eds.; American Chemical Society Press: Washington, DC, 1992; pp 8–32.
- (6) Pirisi, F. M.; Cabras, P.; Cao, C. F.; Migliorini, M.; Muggelli, M. Phenolic compounds in virgin olive oil. 2. Reappraisal of the extraction, HPLC separation, and quantification procedures. *J. Agric. Food Chem.* **2000**, *48*, 1191–1196.
- (7) Vissers, M. N.; Zock, P. L.; Katan, M. B. Bioavailability and antioxidant effects of olive oil phenols in humans: a review. *Eur. J. Clin. Nutr.* **2004**, *58*, 955–965.
- (8) Ramirez-Tortosa, M. C.; Urbano, G.; Lopez-Jurado, M.; Nestares, T.; Gomez, M. C.; Mir, A.; Ros, E.; Mataix, J.; Gil, A. Extra-virgin olive oil increases the resistance of LDL to oxidation more than refined olive in free-living men with peripheral vascular disease. *J. Nutr.* **1999**, *129*, 2177–2183.
- (9) Shyu, S. L.; Huang, J. J.; Yen, G. C.; Chang, R. L. Study on properties and oxidative stability of teaseed oil. *Food Sci.* **1990**, *17*, 114–122.
- (10) Duh, P. D.; Yen, G. C.; Yen, W. J.; Chang, L. W. Antioxidant effects of water extracts from barley (*Hordeum vulgare* L.) prepared under different roasting temperatures. *J. Agric. Food Chem.* **2001**, *49*, 1455–1463.
- (11) Yeh, C. T.; Yen, G. C. Effects of phenolic acids on human phenolsulfotransferases in relation to their antioxidant activity. *J. Agric. Food Chem.* **2003**, *51*, 1474–1479.
- (12) Zhu, Q. Y.; Hackman, R.; Ensunsa, R. R.; Keen, C. L. Antioxidative activities of oolong tea. *J. Agric. Food Chem.* **2002**, *50*, 6929–6934.
- (13) Zou, C.-G.; Agar, N. S.; Jones, G. L. Oxidative insult to human red blood cells induced by free radical initiator AAPH and its inhibition by a commercial antioxidant mixture. *Life Sci.* **2001**, *69*, 75–86.
- (14) Amer, J.; Goldfarb, A.; Fibach, E. Flow cytometric measurement of reactive oxygen species production by normal and thalassemic red blood cells. *Eur. J. Haematol.* **2003**, *70*, 84–90.
- (15) Yen, G. C.; Hsieh, C. L. Inhibitory effect of *Eucommia ulmoides* Oliv. on oxidative DNA damage in lymphocytes induced by H_2O_2 . *Teratog., Carcinog., Mutagen.* **2003**, *1*, 23–34.
- (16) Owen, R. W.; Gisacosa, A.; Hull, W. E.; Haubner, R.; Spiegelhalder, B.; Bartsch, H. The antioxidant/anticancer potential of phenolic compounds isolated from olive oil. *Eur. J. Cancer* **2000**, *36*, 1235–1247.

- (17) Pelter, A.; Ward, R. S.; Watson, D. J.; Collins, P. Synthesis of 2,6-diary-4,8-dihydroxy-3,7-dioxabicyclo[3.3.0]octanes. *J. Chem. Soc., Perkin. Trans.* **1982**, *1*, 175–181.
- (18) Sirato-Yasumoto, S.; Katsuta, M.; Okuyama, Y.; Takahashi, Y.; Ide, T. Effect of sesame seeds rich in sesamin and sesamol in on fatty acid oxidation in rat liver. *J. Agric. Food Chem.* **2001**, *49*, 2647–2651.
- (19) Nakai, M.; Harada, M.; Nakahara, K.; Akimoto, K.; Shibata, H.; Miki, W.; Kiso, Y. Novel antioxidative metabolites in rat liver with ingested sesamin. *J. Agric. Food Chem.* **2003**, *51*, 1666–1670.
- (20) Kiso, Y. Antioxidative roles of sesamin, a functional lignan in sesame seed, and its effect on lipid- and alcohol-metabolism in the liver: a DNA microarray study. *Biofactors* **2004**, *21*, 191–196.
- (21) Hirata, F.; Fujita, K.; Ishikura, Y.; Hosoda, K.; Ishikawa, T.; Nakamura, H. Hypocholesterolemic effect of sesame lignan in humans. *Atherosclerosis* **1996**, *122*, 135–136.
- (22) Miyahara, Y.; Komiya, T.; Katsuzaki, H.; Imai, K.; Nakagawa, M.; Ishi, Y.; Hibasami, H. Sesamin and episesamin induce apoptosis in human lymphoid leukemia Molt 4B cells. *Int. J. Mol. Med.* **2000**, *6*, 43–46.
- (23) Manna, C.; Galletti, P.; Cucciolla, V.; Monttedoro, G.; Zappia, V. Olive oil hydroxytyrosol protects human erythrocytes against oxidative damages. *J. Nutr. Biochem.* **1999**, *10*, 159–165.
- (24) Kitagawa, S.; Sakamoto, H.; Tano, H. Inhibitory effects of flavonoids on free radical-induced hemolysis and their oxidative effects on hemoglobin. *Chem. Pharm. Bull.* **2004**, *52*, 999–1001.
- (25) Cooney, R. V.; Custer, L. J.; Okinaka, L.; Franke, A. A. Effects of dietary sesame seeds on plasma tocopherol levels. *Nutr. Cancer* **2001**, *39*, 66–71.
- (26) Hou, R. C. W.; Hunang, H. M.; Tzen, J. T. C.; Jeng, K. C. G. Protective effect of sesamin and sesamol on hypoxic neuronal and PC12 cells. *J. Neurosci. Res.* **2003**, *74*, 123–133.
- (27) Yamashita, K.; Ikeda, S.; Iizuka, Y.; Ikeda, I. Effect of sesaminol on plasma and tissue alpha-tocopherol and alpha-tocotrienol concentrations in rats fed a vitamin E concentrate rich in tocotrienols. *Lipids* **2002**, *37*, 351–358.
- (28) Serra-Majem, L.; Ribas, L.; Treserras, R.; Ngo, J.; Salleras, L. How could changes in diet explain changes in coronary heart disease mortality in Spain? The Spanish paradox. *Am. J. Clin. Nutr.* **1995**, *61*, 1351–1359.
- (29) Kažimírová, A.; Barančokova, M.; Volkovová, K.; Staruchová, M.; Krajčovičová-Kudláčková, M.; Wsolová, L.; Collins, A. R.; Dušinská, M. Does a vegetarian diet influence genomic stability? *Eur. J. Nutr.* **2004**, *43*, 32–38.
- (30) Kushiro, M.; Takahashi, Y.; Ide, T. Species differences in the physiological activity of dietary lignan (sesamin and episesamin) in affecting hepatic fatty acid metabolism. *Br. J. Nutr.* **2004**, *91*, 377–386.
- (31) Utsunomiya, T.; Shimada, M.; Rikimaru, T.; Hasegawa, H.; Yamashita, Y.; Hamathsu, T.; Yamasaki, M.; Kaku, S.; Yamada, K.; Sugimachi, K. Antioxidant and antiinflammatory effects of a diet supplemented with sesamin on hepatic ischemia-reperfusion injury in rat. *Hepatogastroenterology* **2003**, *50*, 1609–1613.

Received for review September 21, 2005. Revised manuscript received December 12, 2005. Accepted December 21, 2005. This research was partially supported by the Council of Agriculture (93AS-5.1.3-FD-Z1-(2)), Taiwan, Republic of China.

JF052325A