

CORRECTION

Oligomeric Procyanidins in Apple Polyphenol Are Main Active Components for Inhibition of Pancreatic Lipase and Triglyceride Absorption, by Hiroshi Sugiyama,* Yoko Akazome, Toshihiko Shoji, Atsuko Yamaguchi, Masaaki Yasue, Tomomasa Kanda, and Yasuyuki Ohtake, *J. Agric. Food Chem.* 2007, 55, 4604.

Table 3 was incomplete. The correct table is given below.

Table 3. Effect of Apple Polyphenol Extract on Triglyceride Absorption in Humans^a

| | 0 h | 1 h | 2 h | 3 h | 4 h | 6 h |
|--------------|---------|----------|----------|----------|----------|---------------------|
| placebo | 99 ± 10 | 117 ± 16 | 133 ± 29 | 142 ± 30 | 147 ± 36 | 107 ± 13 |
| AP (600 mg) | 75 ± 4 | 99 ± 5 | 110 ± 6 | 125 ± 15 | 130 ± 16 | 77 ± 7 ^b |
| AP (1500 mg) | 86 ± 13 | 102 ± 16 | 103 ± 20 | 114 ± 26 | 119 ± 23 | 108 ± 18 |

^a Values are mean ± SEM. ^b Significantly different from placebo, $P < 0.05$.

JF078004B

10.1021/jf078004b

Published on Web 06/14/2007