

CORRECTIONS

Identification of Cytotoxic Constituents of *Nartheicum ossifragum* Using Bioassay-Guided Fractionation, by Silvio Uhlig,** Helene Wisløff, and Dirk Petersen. *J. Agric. Food Chem.* **2007**, *55*, 6018.

Table 2 contained several errors, which are corrected below.

Table 2. ¹H and ¹³C NMR Assignments for the 5S Isomer of 4-Methoxyfuran-2(5H)-one 5-(β-D-Glucoside)

position	δ H ^a (pyridine-d ₅)	δ C ^b (pyridine-d ₅)
2		170.6
3	5.41 (1H, s)	90.3
4		178.3
5	6.47 (1H, s)	98.3
OCH ₃	3.67 (3H, s)	59.5
1'	5.29 (1H, d, <i>J</i> = 7.9 Hz)	103.5
2'	4.09 (1H, dd, <i>J</i> = 8.8, 7.9 Hz)	74.8
3'	4.21 (1H, dd, <i>J</i> = 8.8, 8.8 Hz)	78.2
4'	4.27 (1H, dd, <i>J</i> = 8.8, 8.8 Hz)	70.8
5'	3.91 (1H, ddd, <i>J</i> = 8.8, 5.0, 2.3 Hz)	79.2
6'a	4.48 (1H dd, <i>J</i> = 12.0, 2.3 Hz)	62.0
6'b	4.34 (1H dd, <i>J</i> = 12.0, 5.0 Hz)	62.0

^a ¹H NMR shifts relative to CD(CD)₂(CH)₂N = 8.71 ppm. ^b ¹³C NMR shifts relative to CD(CD)₂(CH)₂N = 149.8 ppm.

JF8004433

10.1021/jf8004433

Published on Web 02/26/2008