

Differential Analysis of Mycoalexins in Confrontation Zones of Grapevine Fungal Pathogens by Ultrahigh Pressure Liquid Chromatography/Time-of-Flight Mass Spectrometry and Capillary Nuclear Magnetic Resonance

GAETAN GLAUSER,^{†,‡} KATIA GINDRO,^{*,§} JULIANE FRINGELI,[†]
 JEAN-PIERRE DE JOFFREY,[§] SERGE RUDAZ,[‡] AND JEAN-LUC WOLFENDER[†]

Laboratory of Pharmacognosy and Phytochemistry, School of Pharmaceutical Sciences, EPGL, University of Geneva, University of Lausanne, 30 quai Ernest-Ansermet, CH-1211 Geneva 4, Switzerland, Laboratory of Pharmaceutical Analytical Chemistry, School of Pharmaceutical Sciences, EPGL, University of Geneva, University of Lausanne, 20, Bd d'Yvoy, CH-1211 Geneva 4, Switzerland, and Swiss Federal Research Station Agroscope Changins-Wädenswil, Route de Duillier, P.O. Box 1012, CH-1260 Nyon, Switzerland

An original approach was developed for the chemical and biological investigation of zone lines formed by the confrontation of fungi growing in confined spaces. Two wood-decaying fungi involved in esca disease, *Eutypa lata* and *Botryosphaeria obtusa*, were grown in Petri dishes. Metabolic profiles of pure fungal strains and confrontation zones were differentially analyzed by ultrahigh pressure liquid chromatography coupled to time-of-flight mass spectrometry (UHPLC/TOFMS). Selected metabolites induced by the confrontation were isolated and characterized by capillary NMR (CapNMR) at the submilligram level. Fungitoxic and phytotoxic assays were applied to the crude extracts and isolated molecules. While the extracts of pure strains were inactive, the extract from confrontation zones exhibited significant activities. A very strongly induced compound, O-methylmellein, may explain these toxic properties. The developed approach demonstrates the use of fungal confrontations as an original source of bioactive molecules and gives new insights into the study of esca disease.

KEYWORDS: UHPLC/TOFMS; CapNMR; esca; fungi; *Eutypa lata*; *Botryosphaeria obtusa*; mycelial interaction; zone lines; mycoalexins

INTRODUCTION

Esca apoplectic disease of grapevine is due to a complex of fungi, whose species composition and colonization processes are currently not fully understood. The disease is present worldwide and causes important economic losses to growers who must replace infested vines. Currently, although the defense mechanisms against pathogenic microorganisms have been extensively studied (1, 2), no efficient treatment exists against esca. The cause of the disease has been thought to be the conjunction of physiological imbalance of the plant with the presence of fungi. As reported by Graniti et al. (3), esca is not due to a single pathogen but the combination of different fungi. The mycological complexity of esca has led to numerous studies

of the epidemiological aspects of the disease. Previous studies on different grapevine cultivars and from different areas reported the presence of 11 most frequent fungal species in France (4), 5 in Italy (5, 6), and 4 in Germany (7). The six most frequent fungi isolated in Switzerland (8) are the same as those in other countries, namely, *Phaeoconiella chlamydospora*, *Phaeoacremonium angustius*, *Eutypa lata*, *Botryosphaeria obtusa*, *Phomopsis viticola*, and *Fomitiporia mediterranea*. Most of these species are able to colonize and degrade wood constituents, including lignin, leading to the active degradation of sieve elements. Besides this, fungi in close proximity to each other can interact in different ways, namely, in mutualistic, neutralistic, or competitive interactions (9). Competitive mycelial interactions are very important in the overall development of fungal communities in wood. It has been shown that secondary metabolites of some wood-decaying fungi can act as total inhibitors or stimulators of the growth of other wood-decaying fungi (10). In grapevine, fungi that are in competition can form zone lines, also defined as confrontation or barrage zones (Figure 1), which are morphologically different from pure

* Corresponding author. Tel: +41 22 363 43 74. Fax: +41 22 363 43 94. E-mail: katia.gindro@acw.admin.ch.

[†] Laboratory of Pharmacognosy and Phytochemistry, University of Geneva.

[‡] Laboratory of Pharmaceutical Analytical Chemistry, University of Geneva.

[§] Swiss Federal Research Station Agroscope Changins-Wädenswil.




Figure 1. Interactions between fungal phytopathogens. (A) Zone lines in wood. (B) Aspect of the agar culture and confrontation zone formation (arrows) between *Botryosphaeria obtusa* (Bo) and *Eutypa lata* (El) on the upper side of the plate under a magnifying glass.

cultures. The fungi involved mutually inhibit each other and cannot overgrow their opponent (11). Peiris et al. (12) have recently demonstrated the interest of studying metabolite profiles of mycelial fronts between various oak decomposer basidiomycetes using a GC-MS metabolomic-based approach. However, to date, very little attention has been paid to competition phenomena between fungi for a substrate, as a source of phytotoxic or antifungal substances, or disease markers (13). Their chemical analysis in wood samples presents many constraints, such as the difficulty of ensuring the purity of the confronted strains or the precise excision of the zone lines produced. Furthermore, interference from wood constituents can complicate the preliminary analyses, and the amount of trunk necessary to produce enough fungi is prohibitive.

For these reasons, in this study, fungi were grown in solid media using Petri dishes to investigate the production of mycoalexins (i.e., defense compounds synthesized by fungi under attack, analogous with phytoalexins) (14) resulting from competitive interactions. Two of the fungi mentioned above, namely, *Eutypa lata* and *Botryosphaeria obtusa*, were confronted, and a global strategy of investigation including characterization of the mycoalexins and preliminary assessment of their biological activity was developed. Numerous secondary metabolites produced by *Eutypa lata* alone and implicated in grapevine disease have been reported (15–20). Some phytotoxic compounds have also been described in *Botryosphaeria obtusa* (21).

Because of the restricted amounts of material that are available from Petri dishes, very sensitive analytical techniques had to be selected for metabolite profiling and complete de novo characterization of mycoalexins. In this respect, ultrahigh pressure liquid chromatography (22) coupled to time-of-flight mass spectrometry (UHPLC/TOFMS) is known to provide high resolution profiling of complex biological matrices (23, 24). Additionally, the at-line structural determination of natural products at the microgram level (25–27) can be performed by

micro NMR methods designed for mass limited samples such as capillary NMR (CapNMR) (28). A complete integration of these methods in complement to rapid bioassays of original mycoalexins is presented.

MATERIALS AND METHODS

Chemicals. Solvents used for extraction were methanol ACS-ISO for analysis (Carlo Erba SA, Val de Reuil, France), chloroform GR for analysis (Merck, Darmstadt, Germany), and nanopure water (Millipore). ULC/MS grade acetonitrile (CH_3CN) and water from Biosolve (Valkenswaard, The Netherlands) were used for the UHPLC/TOFMS analyses. For the HPLC isolation step, solvents were HPLC grade CH_3CN Chromanorm from VWR (Leuven, Belgium) and nanopure water. For the NMR experiments, methanol- d_4 (CD_3OD ; 99.8% atom deuterium) was obtained from Armar Chemicals (Buchs, Switzerland).

Fungal Growth. Fungi were grown on potato dextrose agar (PDA, Difco) under alternating light and dark periods (12 h/12 h) at 21 °C. The strains of *Botryosphaeria obtusa* and *Eutypa lata* used in this study were isolated from the experimental vineyards of ACW Changins (Switzerland). Growth speed of each fungus was measured each day on PDA for two weeks under the same conditions. Fully developed colonies were used to inoculate 44 Petri dishes (145 mm diam.) using 3 mm diameter fungal chips of the different species on each side of the plate according to the growth speed of each fungus. Cultures were grown under identical conditions until formation of the fungal confrontation zone. Additionally, each fungus was grown separately on PDA in four replicates under the same conditions.

Extraction Procedure. Confrontation zones of 1.5 mm wide were cut and excised using a razor blade, then placed in a vessel with distilled water (100 mL water/100 g extracted agar). Agar media from uninoculated PDA plates and from plates inoculated with each one of both fungal species were used as controls. This initial material was crushed in a Primax blender (Müller & Krempel AG, Bülach, Switzerland) for 5 min and centrifuged (3500g, 15 min, 4 °C). The pellet was washed with water and centrifuged (3500g, 30 min, 4 °C). The extraction was performed in chloroform/methanol/water (64:36:8, v/v) in the dark under agitation (2 h, 4 °C), followed by filtration through LS 14 1/2 filter paper of 500 mm diameter (Schleicher & Schuell GmbH, Dassel, Germany). The filtrate was then dried under vacuum with a rotary evaporator, and the resulting residue was stored at –27 °C until use. Before analysis, a purification procedure using a Sep Pak Vac SPE C18 cartridge (12 cc, 2 g) was performed. MeOH/ H_2O (90:10, v/v) was used for loading and elution to retain very apolar compounds that would be incompatible with reverse-phase chromatography.

UHPLC/TOFMS Experiments. UHPLC/TOFMS analyses were performed on a Micromass-LCT Premier Time of Flight mass spectrometer from Waters (Milford, MA, USA) with an electrospray (ESI) interface coupled with an Acquity UPLC system from Waters. Detection was performed in positive (PI) and negative ion (NI) modes during the same analysis in the range m/z 100–1000 in centered mode with a scan time of 0.2 s and an interscan delay of 0.3 s for polarity switching. ESI conditions in PI and NI modes were capillary voltage 2800 V, cone voltage 40 V, source temperature 120 °C, desolvation temperature 250 °C, cone gas flow 20 L/h, and desolvation gas flow 800 L/h. For internal calibration, a solution of leucine–enkephalin from Sigma-Aldrich (Steinheim, Germany) at 5 $\mu\text{g}/\text{mL}$ was infused through the lockmass probe at a flow rate of 5 $\mu\text{L}/\text{min}$, using a second Shimadzu LC-10ADvp LC pump (Duisburg, Germany). The separation was performed on a 150 mm \times 2.1 mm i.d., 1.7 μm , Acquity BEH C₁₈ UPLC column (Waters) in the gradient mode at a flow rate of 0.3 mL/min with the following solvent system: A = 0.1 vol % formic acid–water, B = 0.1 vol % formic acid–acetonitrile; 2–98% B in 30.0 min. The temperature was set at 30 °C. The injected volume was 5 μL .

LC-MS-Monitored Semipreparative Isolation. The analyses were performed on a Finnigan MAT (San Jose, CA, USA) triple quadrupole mass spectrometer (TSQ700) equipped with a Finnigan ESI interface and coupled to a Varian modular HPLC system (Palo Alto, CA, USA)

with a Varian 9012 pump. The following ESI conditions were used: capillary temperature, 180 °C; source voltage, 4.5 kV; and sheath gas nitrogen, 45 psi. The acquisitions were performed in both PI and NI modes using a full scan mode over an m/z range of 100–1000 and a scan time of 1 s. The separation was performed on a 250 mm \times 10 mm i.d., 5 μ m, Xbridge BEH C₁₈ column (Waters) in the gradient mode at 4.0 mL/min under the following conditions: 2% B for 0.9 min and 2–98% B in 84.1 min. The temperature was set at 30 °C. The injected volume was 200 μ L. Fractions of 4 mL were collected every minute with a fraction collector in 44-well plates from Ratiolab (Dreieich, Germany). A Quicksplit adjustable flow splitter, from ASI (El Sobrante, CA, USA), was used to split 1/200 of the flow to the MS detector.

CapNMR Measurements. CapNMR analyses were carried out on a Varian Unity Inova 500 MHz NMR instrument (Palo Alto, CA, USA) equipped with a 5 μ L microflow CapNMR probe from Protasis/MRM (Savoy, IL, USA) having an active volume of 1.5 μ L. The samples (estimated to 200 μ g) were dissolved in 7 μ L of CD₃OD and parked in the probe with a push volume of 10 μ L through 100 cm of a 100 μ m i.d. capillary tubing. The signal of CD₃OD at 3.31 ppm was used as a reference (temperature 30 °C). For each molecule analyzed, ¹H (16 transients, acquisition time 1 min), gCOSY (8 \times 256 transients \times increments, 21 min), TOCSY (8 \times 256, 93 min), NOESY (32 \times 256, 7 h), gHSQC (16 \times 256, 5 h), and gHMBC (128 \times 256, 23 h) experiments were from the Varian pulse sequence library (vnmr 6.1C, 2003).

Biological Assays. *Antifungal Activity: Effect on Fungal Spore Germination and Development.* *Botrytis cinerea* Pers.: Fr. strain P69, isolated in 1988 on berries of *Vitis vinifera* cv. Pinot noir in the experimental fields of ACW Changins (Switzerland), was grown in Petri dishes on PDA at 21 °C under alternating 12 h light and dark periods. Conidia were collected by vacuum aspiration according to Pezet and Pont (29), sealed in plastic tubes, and stored dry at –80 °C, according to Gindro and Pezet (30) until use. The inhibitory activity on the germination of *Botrytis cinerea* conidia of purified extracts or pure compounds was evaluated on PDA plates. For this purpose, 5 μ L of methanolic solutions (25 μ g/5 μ L and 200 μ g/5 μ L) were spotted on the agar medium (PDA). After methanol evaporation, 4 μ L of an aqueous suspension of 7 \times 10³ conidia/mL were deposited on the dry spots. Blank tests were performed using methanol only. Positive controls were carried out using a 0.1% methanolic solution of the fungicide Switch (Cyprodinil and Fludioxonil) (Syngenta Agro AG, Dielsdorf, Switzerland). Germination rate was determined microscopically by counting 3 \times 50 conidia after 24 h at 21 °C under alternating 12 h light and dark periods. Experiments were done in triplicate. The inhibitory activity on the development of *Botryosphaeria obtusa* and *Eutypa lata* was assessed with slight modifications. Methanolic solutions (25 μ g/5 μ L and 200 μ g/5 μ L) were spotted on a virtual circle at 2 cm of the center, where a 3 mm diameter fungal chip was deposited. Blank tests were performed using methanol only. Positive controls were carried out using a 0.2% methanolic solution of the fungicide Quadrismax (Azoxystrobin and Folpet) (Syngenta Agro AG, Dielsdorf, Switzerland). Growth inhibition was evaluated 10 days after inoculation. Experiments were done in triplicate.

Phytotoxic Activity. The phytotoxic activity of purified extracts or pure compounds was evaluated with respect to the germination rate and further development of seeds of garden cress (*Lepidium sativum* L.). Stock solutions of each purified extract were prepared at 0.1% in dimethylsulfoxide (DMSO)/water (1:19 v/v) and diluted to obtain four solutions (0.1, 0.05, 0.025, and 0.0125%). In the case of pure compounds, a stock solution was prepared at 0.006% in DMSO/water (1:19 v/v) and diluted to obtain two solutions (0.006 and 0.003%). All these solutions were diluted twice with sterilized American Bacteriological agar medium (Laboratorios Conda, Madrid, Spain) and distributed in tips of 200 μ L pipettes, with the apex heat-sealed. A seed of garden cress was then deposited on the surface of the firm medium, and tips were incubated in a humid chamber at 21 °C, with a photoperiod of 16 h daylight and 8 h of darkness. Germination rate, aerial part development, and root length were measured after 3 days. Results are given in length of the aerial part measured in centimeters. Negative controls were performed by replacing extracts with water. A positive control was performed according to the same procedure using




Figure 2. Overview of the general strategy used for chemical and biological characterization of mycoalexins at the Petri dish level.

a known phytotoxic molecule, artemisinin, at a final concentration of 0.033% (Roth, Karlsruhe, Germany) (31). The experiment was repeated 10 times.

RESULTS AND DISCUSSION

The global strategy used in this study for the chemical and biological investigation of confrontation zones between fungi is presented in **Figure 2**.

Fungal Growth and Sample Preparation. Among all the fungal species identified in esca apoplectic grape woods, those isolated most frequently were selected for confrontation tests. The final choice was based on the ability of the selected fungi (*Eutypa lata* and *Botryosphaeria obtusa*) to have a rapid growth rate and to form well-defined confrontation zones. Growth speeds of *Botryosphaeria obtusa* (0.8 mm/h) and *Eutypa lata* (0.2 mm/h) were measured to determine which of the two fungi had the slowest growth and needed to be inoculated first. This point was crucial to obtain broad and well-defined confrontation zones (**Figure 1**). In the case of this specific confrontation, *Botryosphaeria obtusa* was inoculated 5 days after *Eutypa lata*. The whole culture medium including fungi and agar, representing 44 Petri dishes for a total mass of 750 g of material, was extracted in quadruplicate (4 \times 11 dishes). The yield was thereby calculated from the ratio between the initial mass of agar medium and the mass of final extract residue. Sample preparation was kept to a minimum in view of the analysis of the widest possible range of compounds; a simple SPE procedure using a C18-based cartridge was undertaken to retain apolar compounds that could contaminate the UPLC column and instrumentation. This step was determined to be mandatory to avoid carryover and cross-contamination effects. The four replicates represented a total of 170 mg of enriched extract corresponding to a yield of 0.022 \pm 0.01%.

UHPLC/TOFMS Profiling of Fungal Confrontation Zones. Since fungal strains represent biological samples of complex metabolite composition, UHPLC was selected for the differential analysis of the confrontation zone and pure strain extracts. Separation conditions were chosen to maximize peak capacity




Figure 3. UHPLC/TOFMS base peak intensity (BPI) chromatograms of the extracts recorded in positive ion mode. (A) Pure strain of *Eutypa lata*. (B) Pure strain of *Botryosphaeria obtusa*. (C) Mycelial confrontation zone. Compounds 1–4 are highlighted in the different extracts when they are detectable. The amounts injected and the intensity thresholds are identical for all extracts.

(24): an average retention factor in gradient analysis (k_c) of 3 and a peak capacity around 300 were obtained. TOFMS was chosen as a detector for its very high sensitivity in scan mode and for its high mass accuracy, which could give some structural information by providing molecular formulas of the compounds of interest.

Samples were analyzed simultaneously in both PI and NI modes in quadruplicate to detect a maximal number of metabolites and evaluate the consistency of their occurrence. **Figure 3** presents the base peak intensity (BPI) chromatograms of *Eutypa lata*, *Botryosphaeria obtusa*, and confrontation zone extracts, recorded in PI mode. In all cases, the quantity injected was equivalent. In a first step, the data obtained for the *Eutypa lata* and *Botryosphaeria obtusa* chromatograms were processed for the dereplication of known secondary metabolites. The determination of their presence was based on literature data and exact mass measurements by TOFMS (<5 ppm) providing molecular formulas. Although an unambiguous confirmation of their occurrence was not possible in the absence of available standards, six reported vinyl-acetylene derivatives were tentatively identified in the *Eutypa lata* extract, namely, eutypine (MW 186.0681), eutypinol, and its isomer eulatinol (MW 188.0837), sicayne (MW 174.0681), methyleutypinol (MW 202.0994), eutypinic acid (MW 202.0630), and a vinyl-allene derivative (MW 194.0943) (20). It has to be noted that over 20 compounds have been described in *Eutypa lata*; however, metabolite patterns vary according to the origin of the strain (16). This might explain why only part of these compounds were detected. In the *Botryosphaeria obtusa* extract, only two putative isomers of hydroxymellein (MW 194.0579) could be identified.

All of the described compounds appeared in the chromatographic zone of high polarity (0–15 min) and were also present in the confrontation zone extract; however, for these metabolites, no significant induction was observed. As presented in **Figure 3** (PI mode), the confrontation zone extract was obviously richer in compounds than the pure strain extracts, particularly in the high polarity domain of the chromatogram (0–10 min). This observation was confirmed in the NI mode (data not shown). In the BPI chromatogram recorded in PI mode, about 150 peaks were detected in each pure strain, and over 200 peaks in the

confrontation zone. This indicated that numerous unidentified compounds were strongly induced in the region where the fungi came into competition. The differences between the various situations were highly significant. Approximately 60% of the induced metabolites were not detected in the pure strains; they were most likely synthesized de novo as a result of the confrontation. The others were already present in the pure strains but at a much lower level than in the confrontation zone. In particular, some polar substances of low molecular weight recorded in the PI mode (compounds 1–4) (**Figure 3C**) exhibited a very strong MS response in the confrontation extract. Since all of the substances toxic for grapevine reported to date in *Eutypa lata* or *Botryosphaeria obtusa* are small, polar molecules (MW 122–266 Da, e.g., vinyl-acetylene derivatives mentioned above), compounds 1–4 presented a particular interest and were thus selected for further experiments. Compound 1 had a $[M + H]^+$ ion at m/z 193.0870 corresponding to the molecular formula $C_{11}H_{13}O_3$ (error 2.6 ppm). The double bond equivalency (DBE) of 6 meant that the molecule contained 6 rings and/or double bonds. Compounds 2–4 all presented a $[M + H]^+$ ion at m/z 209.0822 ($C_{11}H_{13}O_4$, error 3.8 ppm). They only differed by one oxygen atom from 1 and could be potential hydroxylated derivatives of 1. However, it was not possible to determine the complete structure of these molecules on the basis of only TOFMS information. Therefore, their microisolation by semipreparative LC was undertaken to enable complete characterization by NMR.

MS-Monitored Semipreparative LC Isolation of Induced Metabolites. Because of the small amount of extract available (four extracts representing a total of 170 mg after pooling), the sample consumption during the optimization of the isolation method had to be as low as possible. Thus, a geometrical transfer from the analytical to semipreparative scales was performed to maintain identical selectivity and chromatographic profile (32). The stationary phase chemistry and mobile phase composition were kept identical, while the gradient time, flow rate, and injected volume were adapted to the column geometry. A splitter (for MS detection) and a fraction collector were used. About 160 mg of extract was separated in four successive injections. Two analyses were recorded in each ionization mode. The excellent repeatability of retention times enabled a precise




Figure 4. (A) LC-MS semipreparative chromatogram of the confrontation zone extract recorded in positive ion mode. (B) ¹H CapNMR spectrum of the mixture of 2 and 3. Inset in B: Magnification of the peaks corresponding to the methyl groups, illustrating the difference in signal intensities between both isomers. (C) ¹H CapNMR spectrum of 1. (D) gHMBC CapNMR spectrum of 1. Circles: residual HSQC correlations. * Impurities.

pooling of the fractions obtained from the different injections. Because of the MS monitoring and the selectivity identical to that achieved in UHPLC/TOFMS, the compounds of interest were easily localized. Compound 1 was collected in fractions 23–25 (RT 22–25 min). Compounds 2 and 3 were obtained together from fraction 13 (RT 12–13 min). Compound 4 was present in fraction 16 (RT 15–16 min). Their purity was controlled by UHPLC/TOFMS. Compound 1 was most likely the major product of the extract since the corresponding peak was the only one that saturated the semipreparative column (Figure 4A); about 1.5 mg was isolated. As it was approximately 100 times less abundant in the pure strains, this compound was likely to play a major role in the competition between both species studied. For 2, 3 and 4, about 300–400 μg was obtained. Such amounts were not sufficient to perform 1D and 2D NMR experiments with conventional instruments. Thus, capillary NMR (CapNMR) was selected for the structural elucidation of 1–4. Because of its low active volume, the

CapNMR probe is particularly appropriate for the analysis of limited sample amounts (28). The fractions containing 1–4 were dried and dissolved in 7 μL of CD₃OD before CapNMR experiments.

Structural Elucidation of the Isolated Mycoalexins by CapNMR. *Characterization of Compound 1.* Before performing CapNMR, a hydrogen/deuterium (H/D) exchange experiment was conducted on purified compound 1 (C₁₁H₁₂O₃) using TOFMS and indicated that no labile hydrogen atom was bound to one of the three oxygen atoms present. 1D and 2D CapNMR spectra revealed that 1 held a trisubstituted aromatic ring (3 adjacent protons between δ 6.91–7.55 ppm) with one of the substituents being an *O*-methoxy group (singlet at δ 3.91 ppm) and another an ester group (long-range HMBC correlation between H-7 and C-1 at δ 164.1 ppm; Figure 4D). Additionally, a CH₂-CH-CH₃ chain being part of a methylated pyran-2-one cycle attached to the aromatic ring was present. The protons of the methylene group were diastereotopic (δ 2.89 and 2.98 ppm). These NMR data were in agreement with those reported by




Figure 5. Structures of compounds 1–4. Absolute configuration in C-3 was assumed to be *R* based on previous work (34).

Dimitriadis et al. for *O*-methylmellein (33), and **1** was identified accordingly (**Figure 5**). The complete ¹H and ¹³C signal assignments for **1** are reported in **Table 1**, for comparison with **2–4** and because data were acquired in a different solvent (CD₃OD). **Figure 4C** presents the ¹H CapNMR spectrum and **Figure 4D** the gHMBC CapNMR spectrum of *O*-methylmellein. It clearly indicates the very high degree of purity obtained by the single semipreparative isolation step. The very small amount obtained in this study for **1** prevented the determination of its absolute configuration. Melleins and substituted melleins are known metabolites of many fungi, including *Botryosphaeria obtusa* (21) and *Eutypa lata* (34) with an *R* configuration at C-3. Thus, it was assumed that **1** was *R*-(-)-*O*-methylmellein and that the other derivatives identified shared the same core structure (**Figure 5**). No biological activity has previously been described for *O*-methylmellein, but its very strong induction in the confrontation zone suggests a putative role in the competition between the confronted fungi.

Characterization of Compounds 2, 3, and 4. Because of the mass difference corresponding to an oxygen atom, **2–4** were presumed to be hydroxylated isomers of **1**. H/D exchange experiments on **2–4** revealed one labile hydrogen atom and confirmed a possible hydroxylation of **1**.

With the generic gradient and fractionation time used for semipreparative isolation, **2** and **3** were obtained in the same fraction. **Figure 4B** presents the ¹H CapNMR spectra of the mixture of **2** and **3**. Both isomers exhibited very similar ¹H spectra, closely related to that of **1**. Nevertheless, the deconvolution of their respective signals was possible since slight differences in chemical shifts and an integration ratio of about 1/2 existed between them (**Figure 4B**, inset). The complete ¹H and ¹³C signal assignments for **2** and **3** are reported in **Table 1**. The only significant difference from **1** was the downfield shift of C-4 (δ 67.3 and 69.0 ppm for **2** and **3**, respectively, vs δ 35.4 ppm for **1**) and its attached proton (H-4 at δ 4.56 and 4.51 ppm for **2** and **3**, respectively, vs H-4 α/β at δ 2.89 and 2.98 ppm for **1**). This indicated the presence of a hydroxyl group at C-4 for both **2** and **3**, which was confirmed by a gCOSY correlation between H-3 and H-4. The distinction between **2** and **3** could be made from the different coupling constants between H-3 and H-4. For **3**, a H-3/H-4 coupling constant of 8.3 Hz was observed, indicative of an axial–axial coupling. For **2**, no coupling constant could be detected, suggesting either axial–equatorial or equatorial–equatorial couplings. As for **1**, the absolute configuration could not be determined by NMR.

Compound **4** was found to be a positional isomer of **2** and **3** since the hydroxyl group was bound to C-5 instead of C-4. This was shown by a strong downfield shift of C-5 at δ 147.1 ppm compared to **1**, **2**, and **3** and by the absence of the corresponding aromatic H-5 proton. The position of H-7 remained unchanged

in **1–4** on the basis of a NOESY correlation with the protons of the methoxy group. Since, in the case of **4**, H-7 exhibited an ortho coupling with H-6, the hydroxylation position was ascertained as C-5 (**Table 1**).

In Vitro Antifungal and Phytotoxicity Assays. The biological activity of crude extracts and pure compounds (*O*-methylmellein and its hydroxylated forms) was evaluated in representative fungi and plants, on the basis of developmental criteria of germination and growth. First, the ability to specifically inhibit the development of *Botryosphaeria obtusa* and *Eutypa lata* was considered by microscopic observation of solid media cultures. Remarkably, *O*-methylmellein (25 μ g spotted) showed a very high fungitoxic activity on the development of *Botryosphaeria obtusa*. The mycelial growth was completely stopped, to the same extent as the fungicide control. However, interestingly, no effect was detected on *Eutypa lata*. Hydroxylated forms of *O*-methylmellein at identical concentrations did not exhibit any effect on the growth of *Eutypa lata* or *Botryosphaeria obtusa*. These compounds are present in high concentration in confrontation zones and absent from pure strains. Their concomitant appearance in the mycelial front with the active *O*-methylmellein might indicate the metabolism of the latter by *Botryosphaeria obtusa*, as part of a putative detoxification process. Indeed it has been reported that, apart from metabolite induction, numerous metabolic changes occur during mycelial interactions (35). Therefore, different enzymes implicated in detoxification mechanisms might be activated, especially in the case of fungi involved in the degradation of wood constituents (36). Total extracts from pure fungal cultures and the confrontation zone did not show any effect on the growth speed of *Botryosphaeria obtusa* and *Eutypa lata* (25 and 200 μ g spotted). Due to the low amount of extract available, higher concentrations could not be tested.

In order to estimate the overall fungitoxic and phytotoxic activity of the extracts and compounds described above, the inhibitory activity on the germination of *Botrytis cinerea* conidia and on the germination and growth rate of garden cress (*Lepidium sativum*) was evaluated. In the case of *Botrytis cinerea*, the inhibition was assessed microscopically as was done for the tests on *Eutypa lata* and *Botryosphaeria obtusa*. Here also, the extracts from pure cultures (25 and 200 μ g spotted) did not show any activity. However, the confrontation zone extract significantly affected germination. The conidia germinated by producing very short (1/10 of the control) and swollen germ tubes. In the case of the isolated compounds (25 μ g spotted), *O*-methylmellein presented a high inhibitory activity on germination, showing only the formation of the initial germ tubes, while its hydroxylated metabolites were inefficient.

The phytotoxicity was evaluated using an original assay performed on garden cress on agar medium (**Figure 6**), partly derived from the work of Delabays et al. (31). Only *O*-methylmellein showed a strong antigerminative effect at very low concentration (0.001%), more pronounced than the positive control, artemisinin (0.033%). The hydroxylated forms did not show any effect, with the growth rate of the aerial part and roots identical to the negative water control. The phytotoxicity of the crude extracts was tested at different concentrations. Only the extract resulting from the confrontation area showed a significant effect on growth, resulting in the development of the aerial part six times lower than the water control at a concentration of 0.05% (**Figure 6**). The combined action of both fungi appears thus to increase the toxicity toward garden cress. The strong antigerminative effects observed suggest that the fight between fungi might affect the primary metabolism of the plant.

Table 1. CapNMR Signal Assignments for *O*-Methylmellein (**1**), Diastereoisomers of 4-Hydroxy-8-*O*-methylmellein (**2** and **3**), and 5-Hydroxy-8-*O*-methylmellein (**4**) in CD₃OD^a

position	1, δ H ^b (J)	1, δ C ^c	2, δ H ^b (J)	2, δ C ^c	3, δ H ^b (J)	3, δ C ^c	4, δ H ^b (J)	4, δ C ^c
1		164.1		163.6		163.6		164.6
2								
3	4.58m	74.9	4.59q (6.5)	77.0	4.37m	78.6	4.53m	75.6
3-Me	1.46d (6.3)	19.6	1.48d (6.5)	15.0	1.45d (6.5)	16.8	1.47d (6.5)	19.8
4 α	2.89dd (16.5, 11.2)	35.4	4.56s	67.3	4.51d (8.3)	69.0	2.57dd (16.6, 11.0)	29.2
4 β	2.98dd (16.5, 2.5)						3.18dd (16.6, 2.5)	
4a		142.7		144.0		144.0		128.4
5	6.91d (7.5)	119.4	7.09d (7.6)	119.8	7.23d (7.6)	117.1		147.1
6	7.55t (8.1)	135.1	7.65t (8.2)	135.4	7.67t (8.2)	135.4	7.07d (8.8)	121.2
7	7.07d (8.5)	111.0	7.20d (8.6)	112.8	7.16d (8.6)	112.0	6.91d (8.8)	111.7
8		161.6		160.9		160.9		154.5
8-OMe	3.91s	55.3	3.92s	55.4	3.92s	55.4	3.84s	55.4
8a		113.0		111.8		111.3		113.5

^as, singlet; d, doublet; t, triplet; q, quadruplet; m, multiplet. ^bReference for protons (methanol δ H 3.31 ppm). ^cReference for carbons (methanol δ C 49.0 ppm).




Figure 6. Phytotoxic effects of crude extracts and pure compounds issued from fungal cultures on germination and growth of garden cress: aerial part growth in centimeters is indicated for each case. (Lane A) Positive control: artemisinin (0.033%). (Lane B) *O*-Methylmellein (**1**) (0.001%). (Lane C) 4-Hydroxy-8-*O*-methylmellein (**2+3**) (0.003%). (Lane D) Negative control (water). (Lane E) Crude extract from pure strains of *Botryosphaeria obtusa* at 0.1%. (Lane F) Crude extract from pure strains of *Eutypa lata* at 0.1%. (Lane G–I) Crude extracts from the confrontation zone at 0.05, 0.025, and 0.0125%, respectively.

Consequently, toxicity on grapevine seems highly probable, potentially related to *O*-methylmellein. Further tests on the germination of vine and on wood of mature specimens should be conducted to confirm this hypothesis but will necessitate the production of higher amounts of bioactive extracts and related mycoalexins.

In conclusion, these results clearly indicate that some of the metabolic changes recorded within the confrontation zone play an important role in the specific growth inhibition of *Botryosphaeria obtusa* and more generally on the toxic activity toward other fungi or plants. The metabolite induction phenomena and related bioactivities evidenced shed light on the complex processes implicated in esca disease. Moreover, the herbicide potential of the mycoalexins studied may open new areas of applications in agronomy.

While the ability of fungi to produce active secondary metabolites has been extensively investigated (13), little attention has been paid to the induction of compounds in confrontation zones. The present work has demonstrated the complexity and richness of such zones and their potential for the discovery of new bioactive natural products. Other fungi should be investi-

gated and biological assays conducted to potentially identify, for instance, new lead compounds also active against human pathogens.

ACKNOWLEDGMENT

Dr. Yves Allenbach is thanked for useful comments on the NMR data.

LITERATURE CITED

- (1) Bruno, G.; Sparapano, L. Effects of three esca-associated fungi on *Vitis vinifera* L.: II. Characterization of biomolecules in xylem sap and leaves of healthy and diseased vines. *Physiol. Mol. Plant Pathol.* **2006**, *69*, 195–208.
- (2) Bruno, G.; Sparapano, L. Effects of three esca-associated fungi on *Vitis vinifera* L.: I. Characterization of secondary metabolites in culture media and host responses to the pathogens in calli. *Physiol. Mol. Plant Pathol.* **2006**, *69*, 209–223.
- (3) Graniti, A.; Surico, G.; Mugnai, L. Esca of grapevine: a disease complex or a complex of disease. *Phytopathol. Mediterr.* **2000**, *39*, 16–20.
- (4) Larignon, P.; Dubos, B. Fungi associated with esca disease in grapevine. *Eur. J. Plant Pathol.* **1997**, *103*, 147–157.
- (5) Mugnai, L.; Graniti, A.; Surico, G. Esca (Black measles) and brown wood-streaking: Two old and elusive diseases of grapevines. *Plant Dis.* **1999**, *83*, 404–418.
- (6) Mugnai, L.; Peduto, F.; Calamai, L.; Mattii, G.; Marchi, G.; Surico, G.; Zini, E. Recent developments of research on the factors influencing symptom appearance in esca diseased grapevines in Italy. *Phytopathol. Mediterr.* **2007**, *46*, 113–113.
- (7) Fischer, M.; Kassemeyer, H. H. Fungi associated with Esca disease of grapevine in Germany. *Vitis* **2003**, *42*, 109–116.
- (8) Richter, H.; Gindro, K.; Pezet, R.; Viret, O. Localization and quantification of fungi in esca diseased grapevine trunks. *Phytopathol. Mediterr.* **2007**, *46*, 105–106.
- (9) Rayner, A. D. M.; Todd, N. K. Population and community structure and dynamics of fungi in decaying wood. *Adv. Bot. Res.* **1979**, *7*, 333–420.
- (10) Heilmann-Clausen, J.; Boddy, L. Inhibition and stimulation effects in communities of wood decay fungi: Exudates from colonized wood influence growth by other species. *Microbial Ecol.* **2005**, *49*, 399–406.
- (11) Boddy, L. Interspecific combative interactions between wood-decaying basidiomycetes. *FEMS Microbiol. Ecol.* **2000**, *31*, 185–194.
- (12) Peiris, D.; Dunn, W. B.; Brown, M.; Kell, D. B.; Roy, I.; Hedger, J. N. Metabolite profiles of interacting mycelial fronts differ for pairings of the wood decay basidiomycete fungus, *Stereum hirsutum* with its competitors *Coprinus micaceus* and *Coprinus disseminatus*. *Metabolomics* **2008**, *4*, 52–62.

- (13) Schneider, P.; Misiek, M.; Hoffmeister, D. In vivo and in vitro production options for fungal secondary metabolites. *Mol. Pharmacuetics* **2008**, *5*, 234–242.
- (14) Tokimoto, K.; Fujita, T.; Takeda, Y.; Takaishi, Y. Increased or induced formation of antifungal substances in cultures of *Lentinus edodes* by the attack of *Trichoderma* spp. *Proc. Jpn. Acad., Ser. B* **1987**, *63*, 277–280.
- (15) Jimenez-Teja, D.; Hernandez-Galan, R.; Collado, I. G. Metabolites from *Eutypa* species that are pathogens on grapes. *Nat. Prod. Rep.* **2006**, *23*, 108–116.
- (16) Mahoney, N.; Lardner, R.; Molyneux, R. J.; Scott, E. S.; Smith, L. R.; Schoch, T. K. Phenolic and heterocyclic metabolite profiles of the grapevine pathogen *Eutypa lata*. *Phytochemistry* **2003**, *64*, 475–484.
- (17) Mahoney, N.; Molyneux, R. J.; Smith, L. R.; Schoch, T. K.; Rolshausen, P. E.; Gubler, W. D. Dying-arm disease in grapevines: Diagnosis of infection with *Eutypa lata* by metabolite analysis. *J. Agric. Food Chem.* **2005**, *53*, 8148–8155.
- (18) Molyneux, R. J.; Mahoney, N.; Bayman, P.; Wong, R. Y.; Meyer, K.; Irelan, N. *Eutypa* dieback in grapevines: Differential production of acetylenic phenol metabolites by strains of *Eutypa lata*. *J. Agric. Food Chem.* **2002**, *50*, 1393–1399.
- (19) Lardner, R.; Mahoney, N.; Zanker, T. P.; Molyneux, R. J.; Scott, E. S. Secondary metabolite production by the fungal pathogen *Eutypa lata*: Analysis of extracts from grapevine cultures and detection of those metabolites in planta. *Aust. J. Grape Wine Res.* **2006**, *12*, 107–114.
- (20) Tsoupras, G.; De Angelis, P.; Zesiger, T.; Renaud, J. M.; Tabacchi, R. Biologically-active natural acetylenes from the culture medium of *Eutypa lata*. *Bioact. Mol.* **1988**, *7*, 93–100.
- (21) Venkatasubbaiah, P.; Chilton, W. S. Phytotoxins of *Botryosphaeria obtusa*. *J. Nat. Prod.* **1990**, *53*, 1628–1630.
- (22) Nguyen, D. T. T.; Guillaume, D.; Rudaz, S.; Veuthey, J. L. Fast analysis in liquid chromatography using small particle size and high pressure. *J. Sep. Sci.* **2006**, *29*, 1836–1848.
- (23) Inagaki, S.; Noda, T.; Min, J. Z.; Toyooka, T. Metabolic profiling of rat hair and screening biomarkers using ultra performance liquid chromatography with electrospray ionization time-of-flight mass spectrometry. *J. Chromatogr., A* **2007**, *1176*, 94–99.
- (24) Grata, E.; Boccard, J.; Guillaume, D.; Glauser, G.; Carrupt, P.-A.; Farmer, E. E.; Wolfender, J.-L.; Rudaz, S. UPLC-TOF-MS for plant metabolomics: A sequential approach for wound marker analysis in *Arabidopsis thaliana*. *J. Chromatogr., B* **2008**, *871*, 261–270.
- (25) Wolfender, J. L.; Queiroz, E. F.; Hostettmann, K. Phytochemistry in the microgram domain - a LC-NMR perspective. *Magn. Reson. Chem.* **2005**, *43*, 697–709.
- (26) Hu, J. F.; Garo, E.; Yoo, H. D.; Cremin, P. A.; Zeng, L.; Goering, M. G.; O'Neil-Johnson, M.; Eldridge, G. R. Application of capillary-scale NMR for the structure determination of phytochemicals. *Phytochem. Anal.* **2005**, *16*, 127–133.
- (27) Lin, Y.; Schiavo, S.; Orjala, J.; Vouros, P.; Kautz, R. Microscale LC-MS-NMR platform applied to the identification of active cyanobacterial metabolites. *Anal. Chem.* **2008**, *80*, 8045–8054.
- (28) Olson, D. L.; Norcross, J. A.; O'Neil-Johnson, M.; Molitor, P. F.; Detlefsen, D. J.; Wilson, A. G.; Peck, T. L. Microflow NMR: Concepts and capabilities. *Anal. Chem.* **2004**, *76*, 2966–2974.
- (29) Pezet, R.; Pont, V. Ultrastructural Observations of Pterostilbene Fungitoxicity in Dormant Conidia of *Botrytis cinerea* Pers. *J. Phytopathol.* **1990**, *129*, 19–30.
- (30) Gindro, K.; Pezet, R. Effects of long-term storage at different temperatures on conidia of *Botrytis cinerea* Pers.: Fr. *FEMS Microbiol. Lett.* **2001**, *204*, 101–104.
- (31) Delabays, N.; Slacanin, I.; Bohren, C. Herbicidal potential of artemisinin and allelopathic properties of *Artemisia annua* L.: from the laboratory to the field. *J. Plant Dis. Prot.* **2008**, 317–322.
- (32) Glauser, G.; Guillaume, D.; Grata, E.; Boccard, J.; Thiocone, A.; Carrupt, P.-A.; Veuthey, J.-L.; Rudaz, S.; Wolfender, J.-L. Optimized liquid chromatography-mass spectrometry approach for the isolation of minor stress biomarkers in plant extracts and their identification by capillary nuclear magnetic resonance. *J. Chromatogr., A* **2008**, *1180*, 90–98.
- (33) Dimitriadis, C.; Gill, M.; Harte, M. F. The first stereospecific approach to both enantiomers of mellein. *Tetrahedron: Asymmetry* **1997**, *8*, 2153–2158.
- (34) De Angelis, P. Isolement et caractérisation de l'Eutypine et d'autres métabolites secondaires d'Eutypa Lata (Pers.: Fr.) Tul., parasite de la vigne. Thesis, Faculty of Chemistry, University of Neuchâtel, Switzerland, 1990.
- (35) Rayner, A. D. M.; Griffith, G. S.; Wildman, H. G. Induction of metabolic and morphogenetic changes during mycelial interactions among species of higher fungi. *Biochem. Soc. Trans.* **1994**, *22*, 389–394.
- (36) Cerniglia, C. E. Biodegradation of polycyclic aromatic hydrocarbons. *Biodegradation* **1992**, *3*, 351–368.

Received for review October 27, 2008. Revised manuscript received December 18, 2008. Accepted December 19, 2008.

JF8033539