

## In Situ Seasonal Study of the Volatile Production of Almonds (*Prunus dulcis*) Var. ‘Nonpareil’ and Relationship to Navel Orangeworm

JOHN J. BECK,<sup>\*,†</sup> GLORY B. MERRILL,<sup>†</sup> BRADLEY S. HIGBEE,<sup>§</sup> DOUGLAS M. LIGHT,<sup>†</sup> AND WAI S. GEE<sup>†</sup>

<sup>†</sup>Plant Mycotoxin Research, Western Regional Research Center, Agricultural Research Service, U.S. Department of Agriculture, 800 Buchanan Street, Albany, California 94710, and <sup>§</sup>Paramount Farming Company, 33141 East Lerdo Highway, Bakersfield, California 93308

Nonpareil almonds, *Prunus dulcis*, account for the largest percentage of almond varieties grown in the Central Valley of California. Several studies have investigated the various nonvolatile and volatile components of various plant parts; however, the volatile organic compound (VOC) emission of almonds from a single cultivar has not been studied over the course of a growing season. This aspect is particularly relevant to research concerning the navel orangeworm (NOW), a major insect pest of almonds and other tree nuts. Despite the continued presence of NOW, the identification of particular VOCs and their relationship to NOW have not been addressed. The VOC emission of Nonpareil almonds was collected in situ over the course of a growing season by solid-phase microextraction (SPME). The VOCs (*Z*)-hex-3-enyl acetate, (*Z*)-hex-3-enyl butyrate, undecan-2-ol,  $\beta$ -bourbonene, and tetradecane were present for the majority of the days investigated. Several VOCs exhibited positive electroantennographic signals from male and/or female NOW moths.

**KEYWORDS:** Almond; electroantennogram; in situ; navel orangeworm; Nonpareil; SPME; VOCs

### INTRODUCTION

The Nonpareil almond (*Prunus dulcis*) variety represents the most widely planted almond variety in the Sacramento and San Joaquin Valleys of California and comprises ca. 37% of the total acres of varieties grown (1, 2). California is the top producer of almonds, supplying 80% of the world's needs and 100% of the U.S. market's supply. Approximately 5% of California's cropland is committed to almond production (3). Moreover, the California almond industry is approaching \$2 billion/year (2, 4).

The navel orangeworm (NOW), *Amyelois transitella* (Walker) (Lepidoptera: Pyralidae), is an insect pest of California tree nuts. Its feeding damage lowers nut kernel quality, resulting in extensive monetary loss to growers, producers, and shippers. Moreover, NOW feeding damage contributes to aflatoxin contamination (2).

There are numerous reports in the literature on both the volatile and nonvolatile composition of various tree parts of almond cultivars: e.g., proteins (5) and fatty acids (6) from kernels; fatty acids (7) and triterpenoids (8) from hulls; volatile organic compounds (VOCs) extracted via steam distillation of dried almond hulls (9, 10); almond oil (11); and ex situ VOCs of whole damaged and undamaged almonds (12). Nonetheless, the VOC emission of almonds of any one single cultivar has *not* been studied over the course of an entire

growing season. This aspect is particularly relevant to research concerning NOW. Despite the presence of NOW throughout the almond growth period, the identification of particular VOCs, or their potential role in NOW behavior, has not been addressed.

The discovery of an efficacious attractant for NOW monitoring/trapping has remained elusive despite breakthroughs with the female sex pheromone (13), the pheromone blend (14), long-chain fatty acids (15), or the use of almond press cake or trapped virgin female NOW. The ability of an insect to locate the desired host plant is in part dependent upon its ability to detect a specific VOC (semiochemical). As with the complex blend of NOW pheromone noted by Leal et al. (14), a complex mixture of ubiquitous plant VOCs may be necessary to elicit an appropriate response from the insect to the host plant (16, 17).

This study investigated the VOC emission of Nonpareil almonds in a single growing season from bloom to hull split. The collected VOCs were subjected to electroantennogram (EAG) bioassays with both male and female NOW to determine the responses.

### MATERIALS AND METHODS

**Plant Material.** The VOCs of the fruits of *P. dulcis* (P. Mill) D. A. Webb var. Nonpareil, common name sweet almond, were collected in situ via an inert collection system similar to the reported method (18). A representative VOC collection experiment is illustrated in **Figure 1**. All VOC collections were performed within the

\*Corresponding author [telephone (510) 559-6154; fax (510) 559-6493; e-mail john.beck@ars.usda.gov].



**Figure 1.** In situ VOC collection system showing the inert bag with enclosed almonds and SPME cartridge in place.

same row of Nonpareil almonds between the hours of 8:00 and 11:00 a.m., at S&J Ranch, Madera, CA.

**Volatile Collection.** Three to five almonds were enclosed in a customized Teflon bag (manufactured by SKC West, Inc., Fullerton, CA), 11 × 21 cm, thermally sealed on three sides, with two aluminum portals for SPME (Supelco, Bellefonte, PA; 100  $\mu$ m, polydimethylsiloxane fiber) analysis. The bag was gently sealed onto the stem/branch of the plant by the use of a twist tie, and one SPME was inserted and attached to a portal by means of a large clip (**Figure 1**). VOC collections were kept consistent by applying the following PEST (permeation; exposure; storage; thermal desorption) parameters:  $P = 2$  min;  $E = 30$  min;  $S = 3-6$  h; and  $T = 15$  min (18). The long VOC storage times on the fiber ( $S = 3-6$  h) were due to long transportation times and multiples (up to eight samples) being run on two GC instruments. The SPME cartridges were sealed with Teflon tape, covered with a septum, sealed in a plastic bag, and transported in an ice chest at 0 °C. Background analyses were performed on an unexposed SPME fiber and ambient bag/air volatiles.

**Volatile Analyses.** After transportation from the field, all experiments utilized transfer of adsorbed volatiles onto either a J&W Scientific (Folsom, CA) DB-Wax column (60 m × 0.32 mm i.d. × 0.25  $\mu$ m) or a J&W Scientific DB-1 column (60 m × 0.32 mm i.d. × 0.25  $\mu$ m) installed on one of two HP-6890 gas chromatographs (GC) coupled to HP-5973 mass selective detectors (MS; Palo Alto, CA). Desorbed volatiles were analyzed with the following methods. For DB-Wax: injector temperature, 200 °C; splitless mode; inlet temperature, 200 °C; constant flow, 3.0 mL/min; oven settings, initial temperature, 40 °C; hold time, 0.0 min; ramp 1, 4 °C/min; final temperature, 200 °C; hold time, 40 min. For DB-1: injector temperature, 200 °C; splitless mode; inlet temperature, 200 °C; constant flow, 2.0 mL/min; oven settings, initial temperature, 40 °C; hold time, 0.0 min; ramp 1, 4 °C/min; final temperature, 250 °C; hold time, 30 min. MSD parameters: source temperature, 230 °C; MS source temperature, 150 °C; EI mode, 70 eV; solvent delay, 1 min; scan group 1, 40–300 amu; scan group 2 at 20 min, 40–450 amu. NIST, Wiley, and internally generated databases were used for fragmentation pattern identification. The retention indices (RIs) were calculated using a homologous series of *n*-alkanes on the DB-Wax and DB-1 columns. Volatile identifications were verified by injection of authentic samples and comparison to retention times of an internally generated list of volatiles on identical columns. Each experiment was performed in duplicate, but injected onto separate columns for RI comparison purposes.

Calculated RIs were used to assist in compound identification and to perform comparison of DB-1 to DB-Wax column results. Due to the equilibrium of adsorbed VOCs on the SPME fibers in conjunction with the long VOC storage times, a strict quantitative interpretation of the peak areas was untenable. The relative abundances (peak areas) were noted, and the inclusion of a VOC into **Table 1** was based upon presence in both GC analyses, as well as a minimum peak area based on a percentage of the largest VOC peak present in each run.

**Electroantennogram Bioassays.** The antennae of laboratory-reared, sexed navel orangeworm moths, *A. transitella* (Walker) (Lepidoptera: Pyralidae), were excised, positioned on a fork electrode using electrode gel, and connected to an IDAC-4 acquisition controller electroantennogram using Syntech's PC-based software (Syntech, Kirchzarten, Germany). The antennae were humidified with a stream of purified air bubbled through distilled water at a flow rate of 200 mL/min. The individual compounds for EAG analysis (50  $\mu$ g; 10  $\mu$ L of a 5  $\mu$ g/ $\mu$ L solution in pentane) were loaded onto oven-dried 0.25 in. assay disks, allowed to air-dry for 5 min, and inserted into 5.75 in. Pasteur pipets, and the ends were temporarily capped with parafilm. The sources of the VOCs for EAG analyses are noted in **Table 1**. Negative control (NegCtrl) disks were prepared using a similar method, but with 10  $\mu$ L of pentane. Positive control (PosCtrl) disks were prepared using the major pheromone component (*Z,Z*)-11,13-hexadecadienal (50  $\mu$ g; Suterra, Bend, OR). The pipets loaded with the individual compounds were attached via tubing to a stimulus controller unit (Syntech). The antennae were exposed to each compound by a 2 s puff of air, and the resulting response was recorded. The antennal response was duplicated for each VOC with a 1 min delay between puffs. Each antenna pair was exposed to seven duplicated puffs in the order PosCtrl, compound A, NegCtrl, compound B, compound C, compound D, and PosCtrl, with each run lasting no longer than 30 min from excision to completion of run on the antenna pair.

Male NOW responses ( $\mu$ V) to the individual VOCs were normalized to the averaged EAG responses ( $\mu$ V) of the male to the PosCtrl and less the response to the NegCtrl. The lowest response to the PosCtrl of 500  $\mu$ V was used as the normalization factor. The average male response to the PosCtrl was 1100  $\mu$ V over the course of all VOCs tested. The crude male EAG responses (MaleEAG) were normalized using the equation  $\text{MaleEAG} \times (500/\text{PosCtrl}) - \text{NegCtrl}$ . Therefore, any VOC demonstrating an EAG value of 500  $\mu$ V or more would be judged as good as or better than the PosCtrl. Because there are no current "standards" to judge the female response, and females do not typically provide strong or consistent response to the pheromone, the female responses were stringently normalized by subtraction of response to both the PosCtrl and the NegCtrl. The crude female EAG responses (FemaleEAG) were normalized using the equation  $\text{FemaleEAG} - \text{PosCtrl} - \text{NegCtrl}$ . The values shown in **Table 1** are the average of the duplicate responses. Any normalized response of the female > 500  $\mu$ V would be considered to be significant. Normalized responses of ca. 250  $\mu$ V were considered to be moderate responses.

## RESULTS AND DISCUSSION

The in situ collection and analysis of Nonpareil almonds provided 24 VOCs in various amounts and frequency. One interesting result was the small number, yet diverse class, of compounds emitted over the majority (> 55%) of the growing season: (*Z*)-hex-3-enyl acetate, **4**; (*Z*)-hex-3-enyl butyrate, **10**; undecan-2-ol, **12**;  $\beta$ -bourbonene, **16**; and, tetradecane, **18**. Of these, only  $\beta$ -bourbonene, **16**, and tetradecane, **18**, were present in the earlier months, both of which are common floral scent VOCs of numerous plants, including several genera from the family Rosaceae (19). Neither of these persistent VOCs has demonstrated semiochemical characteristics for Lepidoptera in the literature, and tetradecane, **18**, elicited only weak EAG activity from the female in this study (**Table 1**).  $\beta$ -Bourbonene, **16**, has been reported to demonstrate a slight increase in relative amount during a study of damaged ex situ almonds (12), however, was not available in isolable amounts for EAG bioassay.

The absence of the  $C_6$  and  $C_9$  compounds early in the almond growth stages supports the idea of the enzymatic breakdown of the fatty acids in almonds (20) to  $C_6$  green leaf volatiles, in this case  $C_6$  esterified derivatives, (*Z*)-hex-3-enyl acetate, **4**, and (*Z*)-hex-3-enyl butyrate, **10**, and the  $C_9$

Table 1. Volatile Emission of Nonpareil Almonds from Bloom to Hull Split, March–August 2007, Madera, CA<sup>a</sup>

compd library/ID	source <sup>d</sup> lit. <sup>e</sup> calcd <sup>f</sup> lit. <sup>e</sup> calcd <sup>f</sup>	DB-Wax RI		VOC occurrence, m/ddd 2007 (tree no.) <sup>c</sup>																			EAG response (μV)							
		DB-1 RI	DB-2 RI	3/21 (8)	3/28 (3)	4/04 (7)	4/12 (11)	4/18 (14)	4/25 (3)	5/02 (7)	5/07 (8)	5/14 (14)	5/18 (15)	5/21 (16)	6/01 (20)	6/04 (21)	6/11 (22)	6/18 (24)	6/25 (25)	7/02 (26)	7/09 (28)	7/16 (31)	7/23 (34)	7/30 (34)	8/06 (35)	8/13 (37)	M	F		
1 (Z)-hex-3-en-1-ol	Iso	834	838	1381	1388	+		+		+		+		+		tr	+		+		+		+		+		+		0	0
2 butyrolactone	Ald	855	859	1623	1627	+	tr	++																				+	0	25
3 styrene	Ald	871	872	1252	1253							tr																	0	0
4 (Z)-hex-3-enyl acetate	Bed	986	987	1312	1317						++++	++++	++++	+	+							++	+					+	0	0
5 hexyl acetate	Ald	995	993	1268	1272																								0	0
6 nonanal	Ald	1082	1081	1389	1392																								0	50
7 2-phenylethanol	Ald	1081	1082	1910	1912						tr																		54	320
8 linalool	Iso	1083	1084	1546	1549						tr																		58	200
9 non-2-enal	Ald	1134	1132	1532	1533																								183	200
10 (Z)-hex-3-enyl butyrate	Bed	1167	1168	1458	1461						tr																		238	75
11 undecan-2-one	Iso	1273	1271	1596	1597																								387	275
12 undecan-2-ol	Ald	1286	1284	1721	1724																								0	25
13 methyl anthranilate	Iso	1302	1301	2238	2232						tr																		33	65
14 α-copaene	Iso	1374	1374	1490	1489						+																		30	20
15 farnesane	Iso	1379	1378	1358	1354																								45	0
16 β-bourbonene	Tent	1382	1381	1516	1515																								-	-
17 decyl acetate	Iso	1393	1392	1679	1678																								264	75
18 tetradecane	Poly	1400	1398	1400	1397																								0	25
19 caryophyllene	Iso	1415	1413	1594	1593																								0	190
20 geranylacetone	Ald	1429	1428	1853	1855						tr																		88	200
21 α-humulene	Iso	1449	1455	1666	1666																								187	250
22 germacrene D	Iso	1474	1473	1707	1713																								20	115
23 pentadecane	Poly	1500	1499	1500	1489																								0	150
24 hexadecane	Poly	1600	1596	1600	1597																								74	40

<sup>a</sup> Volatile relative peak area ( $\times 10^6$ ): tr, trace, <0.5; +, 0.5–20; ++, 20–50; +++, 50–150; +++++, 150–400. <sup>b</sup> Compound identification by RI relative to *n*-alkanes on DB-1 and DB-Wax columns, retention times, mass fragment libraries, and comparison to authentic samples. <sup>c</sup> Volatiles sampled on days noted between 8:00 a.m. and 11:00 a.m. Trees analyzed were from the same orchard. <sup>d</sup> Source of compound for authentication: Ald, Aldrich or Fluka; Bed, Bedoukian; Iso, previously isolated by a researcher in this laboratory (Buttery, Flath, or Binder); Poly, Polyscience Corp.; Tent, tentative. <sup>e</sup> Compound RI of literature value. <sup>f</sup> Calculated compound RI based upon retention time of respective column.



compounds, nonanal, **6**, and non-2-enal, **9** (21). The generation of the C<sub>6</sub> and C<sub>9</sub> compounds also correlates with the phenological production of fatty acid oils in the kernels, which generally increases at about 60 days after fertilization (22). The origin of undecan-2-ol, **12**, is speculative at this juncture because it has been detected from both plant and microbial sources. Undecan-2-ol, **12**, and undecan-2-one, **11**, have been reported from trees (23) and fruit (24), among others. Moreover, both of these compounds have been detected as volatiles from a myxobacterium (25), although one not found on almonds. Alternatively, it is possible that the undecan-2-ol, **12**, may ultimately be derived from a fatty acid in addition to the C<sub>6</sub> and C<sub>9</sub> compounds described above. The oxidized form of lauric acid is also formed during the enzymatic breakdown of fatty acids (21), and it is known that undecan-2-one, **11**, can be generated from lauric acid (26). Therefore, it could be hypothesized that undecan-2-one, **11**, and undecan-2-ol, **12**, can be generated in this manner, thus explaining the similarities to the kernel phenological growth pattern. This hypothesis is being explored for merit.

The electroantennogram (EAG) data are very interesting with respect to a potential relationship between almond VOCs and NOW. The acetylated green leaf volatile (*Z*)-hex-3-enyl acetate, **4**, and the alcohol undecan-2-ol, **12**, did not show any notable EAG response from either male or female; however, the other corresponding VOCs did demonstrate sufficient EAG responses to warrant interest as prospective background signaling volatiles (BSVs) (17). For the purpose of this study, BSVs are defined as ubiquitous volatiles from almonds that may act as obligatory cues to direct NOW toward key attractant(s). Hence, the BSVs need not demonstrate an EAG response greater than a specified attractant, such as the major aldehyde component of the female NOW pheromone (13), but rather a reasonable EAG response that suggests a basal interest in the individual VOC or bouquet.

The VOCs (*Z*)-hex-3-enyl butyrate, **10**,  $\beta$ -caryophyllene, **19**, geranylacetone, **20**, and  $\alpha$ -humulene, **21**, elicited modest to moderate EAG responses from the male and/or female NOW and were present for > 40% of the growing season. The VOC  $\alpha$ -humulene, **21**, synonymous with  $\alpha$ -caryophyllene, provided similar normalized responses for both the male and female antennae; it is a ubiquitous plant volatile and is known to possess semiochemical behavior, albeit none is reported for NOW (27). (*Z*)-Hex-3-enyl butyrate, **10**,  $\beta$ -caryophyllene, **19**, and geranylacetone, **20**, elicited preferential male/female EAG signals: 238/75, 0/190, and 88/200  $\mu$ V responses, respectively. Geranylacetone, **20**, was not present for > 55% of the evaluation period; however, it did show a modest yet persistent presence toward the later collections and tapered off toward hull-split. Geranylacetone, **20**, (*Z*)-hex-3-enyl butyrate, **10**, and  $\beta$ -caryophyllene, **19**, are ubiquitous plant VOCs; however, none demonstrate notable semiochemical behavior, although  $\beta$ -caryophyllene, **19**, has shown attractant behavior toward one Lepidoptera, Tortricidae (28). Finally, undecan-2-one, **11**, elicited the highest normalized combined male/female responses from NOW antennae, although present only during the later months of collection.

Other compounds, albeit infrequent and in trace amounts, that elicited notable EAG responses included 2-phenylethanol, **7**, linalool, **8**, non-2-enal, **9**, and decyl acetate, **17**. One observation can be made regarding these "transient" VOCs—they all have documented origins from Orchidaceae plant fatty acids (29). In fact, 12 of the 24 compounds (**5–9**, **11**, **17–21**, and **23**) listed in **Table 1** were listed as having plant fatty

acid origins, and of those 12, 8 VOCs demonstrated modest to moderate responses from male and/or female NOW antennae. Moreover, it is interesting to note the presence of long-chain alkanes tetradecane, **18**, and pentadecane, **23**, at the early stages of almond growth.

The VOCs from a single cultivar of almonds were collected in situ over the 2007 San Joaquin Valley growing season. The cultivar Nonpareil was chosen primarily for its agronomic and organoleptic characteristics, which are compulsory to uniformity. The VOCs 2-phenylethanol, **7**, (*E*)-non-2-enal, **9**, undecan-2-one, **11**, and  $\alpha$ -humulene, **21**, produced the highest combined EAG responses from the male and female antennae. In addition to the VOCs with notable NOW EAG activity, several VOCs demonstrated promising BSV characteristics. This is the first report of VOC emissions collected in situ from a single almond cultivar over the course of a growing season; however, the VOCs provided in **Table 1** provide only a sampling of BSV candidates and/or VOC emissions. Further replications of tree VOCs are required to confidently assign a pattern between VOC emission and phenological development. Investigations into the possible fatty acid origin of several of the almond VOCs are ongoing.

#### ABBREVIATIONS USED

BSV, background signaling volatile; EAG, electroantennogram; GC-MS, gas chromatography–mass spectroscopy; NOW, navel orangeworm; PEST, permeation/exposure/storage/thermal desorption; RIs, retention indices; SPME, solid-phase microextraction; VOC, volatile organic compound.

#### ACKNOWLEDGMENT

We thank James Bettiga (S&J Ranch) and James L. Baker (USDA-ARS) for their valuable contributions and the anonymous reviewers for their helpful suggestions.

#### LITERATURE CITED

- (1) USDA NASS 2007 California Almond Acreage Report, [http://www.nass.usda.gov/Statistics\\_by\\_State/California/Publications/recent\\_reports.asp](http://www.nass.usda.gov/Statistics_by_State/California/Publications/recent_reports.asp) (accessed Oct 28, 2008).
- (2) Almond Board of California, California Almond facts and The High Cost of Aflatoxins, <http://www.almondboard.com> (accessed Oct 28, 2008).
- (3) Epstein, L.; Bassein, S.; Zalom, F. G.; Wilhoit, L. R. Changes in pest management practice in almond orchards during rainy season in California, USA. *Agric. Ecosyst. Environ.* **2001**, *83*, 111–120.
- (4) Rowhani, O.; Sumner, D. A. California International Agricultural Exports in 2006. *AIC Issues Brief* 2007, *32* (<http://aic.ucdavis.edu/pub/briefs/brief32.pdf>, accessed Oct 28, 2008).
- (5) Ahrens, S.; Venkatachalam, M.; Mistry, A. M.; Lapsley, K.; Sathe, S. K. Almond (*Prunus dulcis* L.) protein quality. *Plant Foods Hum. Nutr.* **2005**, *60*, 123–128.
- (6) Kodad, O.; Socias I Company, R. Variability of oil content and of major fatty acid composition in almond (*Prunus amygdalus* Batsch) and its relationship with kernel quality. *J. Agric. Food Chem.* **2008**, *56*, 4096–4101.
- (7) Takeoka, G. R.; Dao, L. T. Antioxidant constituents of almond [*Prunus dulcis* (Mill.) D.A. Webb] hulls. *J. Agric. Food Chem.* **2003**, *51*, 496–501.
- (8) Takeoka, G.; Dao, L.; Teranishi, R.; Wong, R.; Flessa, S.; Harden, L.; Edwards, R. Identification of three triterpenoids in almond hulls. *J. Agric. Food Chem.* **2000**, *48*, 3437–3439.
- (9) Buttery, R. G.; Soderstrom, E. L.; Seifert, R. M.; Ling, L. C.; Haddon, W. F. Components of almond hulls: possible navel orangeworm attractants and growth inhibitors. *J. Agric. Food Chem.* **1980**, *28*, 353–356.

- (10) Buttery, R. G.; Seifert, R. M.; Haddon, W. F.; Lundin, R. E. 2-Hexyl-3-methylmaleic anhydride: an unusual volatile component of raisins and almonds hulls. *J. Agric. Food Chem.* **1980**, *28*, 1336–1338.
- (11) Miraliakbari, H.; Shahidi, F. Oxidative stability of tree nut oils. *J. Agric. Food Chem.* **2008**, *56*, 4751–4759.
- (12) Beck, J. J.; Higbee, B. S.; Merrill, G. B.; Roitman, J. N. Comparison of volatile emissions from undamaged and mechanically damaged almonds. *J. Sci. Food Agric.* **2008**, *88*, 1363–1368.
- (13) Coffelt, J. A.; Vick, K. W.; Sonnet, P. E.; Doolittle, R. E. Isolation, identification, and synthesis of a female sex pheromone of the navel orangeworm, *Amyelois transitella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.* **1979**, *5*, 955–966.
- (14) Leal, W. S.; Parra-Pedrazzoli, A. L.; Kaissling, K.-E.; Morgan, T. I.; Zalom, F. G.; Pesak, D. J.; Dundulis, E. A.; Burks, C. S.; Higbee, B. S. Unusual pheromone chemistry in the navel orangeworm: novel sex attractants and a behavioral antagonist. *Naturwissenschaften* **2005**, *92*, 139–146.
- (15) Phelan, P. L.; Roelofs, C. J.; Youngman, R. R.; Baker, T. C. Characterization of chemicals mediating ovipositional host-plant finding by *Amyelois transitella* females. *J. Chem. Ecol.* **1991**, *17*, 599–613.
- (16) Bruce, J. J. A.; Wadhams, L. J.; Woodcock, C. M. Insect host location: a volatile situation. *Trends Plant Sci.* **2005**, *10*, 269–274.
- (17) Schroder, R.; Hilker, M. The relevance of background odor in resource location by insects: a behavioral approach. *Bioscience* **2008**, *58*, 308–316.
- (18) Beck, J. J.; Smith, L.; Merrill, G. B. In situ volatile collection, analysis, and comparison to three *Centaurea* species and their relationship to biocontrol with herbivorous insects. *J. Agric. Food Chem.* **2008**, *56*, 2759–2764.
- (19) Knudsen, J. T.; Eriksson, R.; Gershenzon, J.; Stahl, B. Diversity and distribution of floral scent. *Bot. Rev.* **2006**, *72*, 1–20.
- (20) Sathe, S. K.; Seeram, N. P.; Kshirsagar, H. H.; Heber, D.; Lapsley, K. A. Fatty acid composition of California grown almonds. *J. Food Sci.* **2008**, *73*, C607–C614.
- (21) Noordermeer, M. A.; Veldink, D. A.; Vliegthart, J. F. G. Fatty acid hydroperoxide lyase: a plant cytochrome P450 enzyme involved in wound healing and pest resistance. *ChemBioChem* **2001**, *2*, 494–504.
- (22) Kumar, K.; Sharma, S. D.; Goyal, R. K. Enzymatic changes and oil accumulation during almond kernel development. *Acta Hort. (ISHS)* **2000**, *515*, 287–295.
- (23) Tayoub, G.; Schwob, I.; Bessiere, J.-M.; Masotti, V.; Rabier, J.; Ruzzier, M.; Viano, J. Composition of volatile oils of *Styrax officinalis* L. leaves at different phenological stages. *Biochem. Syst. Ecol.* **2006**, *34*, 705–709.
- (24) Zabetakis, I.; Holden, M. A. Strawberry flavour: analysis and biosynthesis. *J. Sci. Food Agric.* **1997**, *74*, 421–434.
- (25) Dickschat, J. S.; Bode, H. B.; Wenzel, S. C.; Muller, R.; Schulz, S. Biosynthesis and identification of volatiles released by the myxobacterium *Stigmatella aurantiaca*. *ChemBioChem* **2005**, *6*, 2023–2033.
- (26) Dartey, C. K.; Kinsella, J. E. Metabolism of [U-<sup>14</sup>C]lauric acid to methyl ketones by the spores of *Penicillium roqueforti*. *J. Agric. Food Chem.* **1973**, *21*, 933–936.
- (27) Bedard, C.; Gries, R.; Gries, G.; Bennet, R. *Cydia strobilella* (Lepidoptera: Tortricidae): antennal and behavioral responses to host and nonhost volatiles. *Can. Entomol.* **2002**, *134*, 793–804.
- (28) El-Sayed, A. M. The Pherobase: Database of Insect Pheromones and Semiochemicals, 2007; <http://www.pherobase.com> (accessed Jan 26, 2009).
- (29) Johnson, S. D.; Steiner, K. E.; Kaiser, R. Deceptive pollination in two subspecies of *Disa spathulata* (Orchidaceae) differing in morphology and floral fragrance. *Plant Syst. Evol.* **2005**, *255*, 87–98.

---

Received for Review January 28, 2009. Revised manuscript received March 10, 2009. Accepted March 14, 2009. This research was conducted under USDA-ARS CRIS Project 5325-42000-036-00 and CRADA 58-3K95-7-1198.