

Discrimination of Animal Species Using Polymorphisms of the Nuclear Gene *Zinc Finger Protein 238*

WON KIM,[†] SUNMI KIM,[†] HOJUN CHOI,[†] NGUYEN DINH TRUONG,[†] LE MINH THONG,[†]
JIN-HOI KIM,[†] RUI XIAO,[†] KEUN-KYU PARK,[§] KUNHO SEO,[#] HANG LEE,[‡]
BO-SOOK KIM,[⊗] MI-HYUN YOO,[⊗] AND CHANKYU PARK^{*,†,§}

[†]Department of Animal Biotechnology, [§]Animal Resources Research Center, and [#]Colleges of Veterinary Medicine, Konkuk University, Seoul 143-701, South Korea, [‡]Conservation Genome Resource Bank for Korean Wildlife (CGRB), College of Veterinary Medicine, Seoul National University, Seoul 151-742, South Korea, and [⊗]Division of Animal Research, Seoul Zoo, Seoul 427-702, South Korea

We screened 3750 single exonic genes listed in the intronless genes in the eukaryotes (SEGE) database and performed bioinformatic analyses to identify candidate genes for new species-specific markers. A set of PCR primers for the conserved regions of *ZNF238* was developed and used to amplify the 823 bp DNA fragment. We compared nucleotide variations of the PCR products among 20 species plus two subspecies of animals, which led to the identification of interspecies nucleotide variations. To establish a simple method for the analysis of species-specific DNA polymorphisms using *ZNF238*, we developed a PCR-RFLP method using *HhaI* and *HpyCH4IV* restriction enzymes for 13 species. For the remaining species, the direct sequencing of PCR products provided additional SNPs, enabling precise species classification. As a result, we report here that a new nuclear DNA marker, *ZNF238*, can be used to increase the accuracy of species identification among euteleostomi (bony vertebrates).

KEYWORDS: Species discrimination; PCR-RFLP; *ZNF238*; nuclear DNA

INTRODUCTION

Accurate species identity determination may be important in forensic science, food safety, and other fields. The idea of using species-specific DNA sequences for species determination was introduced as early as the 1970s (1, 2). Due to the availability of genome-level DNA sequence information from diverse species, many studies have attempted to establish simple and accurate methods for classifying species information from unknown origins (3).

The mitochondrial genome has advantages in species identification and phylogenetic analysis, including a high copy number within the cell (4) and higher polymorphism generation than the nuclear genome, due to its faster evolution (5). Therefore, the most frequently studied genes in animal species identification have belonged to the mitochondrial genome regions, including *Cytochrome b* (6), *subunits of cytochrome c oxidase* (7), *D-loop* (8), and *12S* (9) and *16S rRNA* (10). In nuclear genes, *internal transcribed spacers 1-rDNA* (11), *recombination activation gene* (12), *introns of actin gene* (13), *myoglobin* (14), and *β-fibrinogen genes* (15) have been tested for species identification. However, nuclear genes are used less frequently in actual application, because the use of any single nuclear gene alone has proven to be insufficient for the classification of vertebrate animal species. This

is due to the difficulty in finding consistent interspecies DNA polymorphisms.

In this study, we sought to identify a nuclear gene that can be used to determine the species identity more efficiently as a single locus system from nuclear DNA, even for closely related vertebrate species. We identified a region from *ZNF238* that is highly conserved among different animal species and yet contains species-specific SNPs. The nuclear gene *ZNF238* is a common gene among euteleostomi (<http://www.ncbi.nlm.nih.gov/homologene>) and encodes zinc finger protein 238, which acts as a transcriptional activator or repressor and is involved in chromatin assembly (16). We show that *ZNF238* can be used as a novel nuclear DNA marker system for the species identification of bony vertebrates.

MATERIALS AND METHODS

DNA Extraction. To test the specificity of the technique, the following numbers of unrelated specimens were tested: 10 dogs (*Canis lupus familiaris*), 6 wolves (*Canis lupus chanco*), 4 raccoons (*Nyctereutes procyonoides*), 10 pigs (*Sus scrofa domestica*, 4 breeds), 3 wild boars (*Sus scrofa coreanus*), 8 mice (*Mus musculus*, 4 strains), 10 horses (*Equus caballus*), 5 rabbits (*Lepus brachyurus*), 10 cattle (*Bos taurus*, 2 breeds), 4 Chinese water deer (*Hydropotes inermis*), 2 goats (*Capra aegagrus hircus*), 4 western roe deer (*Capreolus capreolus*), 8 Formosan deer (*Cervus nippon taiouanus*), 10 sheep (*Ovis aries*), 10 kangaroos (*Macropus rufus*), 10 ducks (*Anas platyrhynchos*), 10 chickens (*Gallus gallus*), 5 pheasants (*Phasianus colchicus*), 3 vultures (*Aegypius monachus*), 4 falcons (*Falco peregrinus japonensis*), 3 elephants (*Loxodonta africana*), and 3 hippopotamuses (*Hippopotamus amphibius*). Animal tissues were obtained either from the

*Address correspondence to this author at the Department of Animal Biotechnology, Konkuk University, Hwayang-dong, Kwangjin-gu, Seoul 143-701, Korea (telephone 82-2-450-3697; fax 82-2-457-8488; e-mail chankyu@konkuk.ac.kr).

conservation genome resource bank for Korean wildlife (<http://www.cgrb.org>), Seoul Zoo (<http://grandpark.seoul.go.kr>), local farms, local meat stores, and other researchers. Genomic DNA was prepared from tissue samples by a simple lysis method (17).

PCR Primers and Amplification. The nucleotide sequence information of *ZNF238* was obtained from Genbank for six species (*Homo sapiens*, NM_205768.2; *Bos taurus*, NM_001078016.1; *Equus caballus*, XM_00149-2205.2; *Mus musculus*, NM_001012330.1; *Canis lupus familiaris*, XM_547-497.2; *Gallus gallus*, XM_426137.2). Nucleotide sequence alignments were performed with the ClustalW2 program (<http://www.ebi.ac.uk/Tools/clustalw2>), and PCR primers were designed from a highly conserved region of *ZNF238* using the primer designer V 2.0 (Scientific & Educational Software, USA). The primer sequences were 5'-ACC TGT CTG TCA AGT CCA GCC T-3' and 5'-TCT TCT AAG GTC CAG TCT CTG A-3' for the forward and reverse primers, respectively. PCR consisted of an initial denaturation step at 94 °C for 5 min, followed by 35 cycles of denaturation at 94 °C for 45 s, primer annealing at 66 °C for 45 s, and product extension at 74 °C for 45 s in a Thermocycler 3000 (Biometra, Germany). A final extension step was performed at 94 °C for 7 min. PCR amplifications were performed in a total volume of 15 μ L, containing 50 ng of genomic DNA, 0.3 μ M of each primer, 200 μ M of dNTPs, and 0.5 unit of Supertherm DNA polymerase (LPI, U.K.) in a PCR reaction buffer (1.5 mM MgCl₂).

PCR-RFLP. Restriction enzymes for RFLP analysis were selected on the basis of the analysis results of restriction enzyme sites using webcutter V2.0 (<http://rna.lundberg.gu.se/cutter2>) or NEBcutter V2.0 (<http://tools.neb.com/NEBcutter2>). The PCR amplicon of *ZNF238* was digested with 10 units of *HhaI* (New England Biolabs, USA) or *HpyCH4IV* (New England Biolabs, USA) in a final volume of 20 μ L for 3 h at 37 °C. The resulting fragments were separated by gel electrophoresis on 3% agarose gels in 1 \times TAE running buffer for 60 min at 100 V, stained with ethidium bromide (Sigma-Aldrich, USA), and visualized under UV light. The size of the restriction fragments was estimated using a DNA size marker (G&P Life Science, Korea).

Direct Sequencing of PCR Products. The PCR products were treated with ExoSAP-IT (USB Corporation, USA) for 15 min at 37 °C, followed by incubation at 80 °C for 15 min. One microliter of reaction mix was subjected to sequencing reactions performed in a total volume of 10 μ L with 0.5 μ M primer (5'-ACC TGT CTG TCA AGT CCA GCC T-3' [forward] and 5'-TCT TCT AAG GTC CAG TCT CTG A-3' [reverse]) using a BigDye Terminator v3.1 cycle sequencing kit (Applied Biosystems, USA) according to the manufacturer's protocols. Purification of the sequencing reactions was performed using Magnesil (Promega, USA) in 25 μ L, and sequencing runs were performed on an ABI 3730 DNA analyzer (Applied Biosystems, USA). The results were analyzed by a CLC Main workbench 5.0.2 (CLC bio, Denmark).

RESULTS AND DISCUSSION

In contrast to mitochondrial DNA-based markers, no single nuclear DNA marker alone has proven to be discriminative in diverse animal species. Therefore, we sought to develop a simple nuclear DNA-based method for species identification of vertebrates. In this study, we evaluated the possible use of *ZNF238* as a successful nuclear DNA marker that can be applied to most euteleostomi for species identification. We examined the polymorphisms of *ZNF238* from as many vertebrate species ($n = 20$) as we could to evaluate the marker applicability of *ZNF238* against a wide variety of vertebrates, including major domestic animals and birds.

***ZNF238* Was Identified as a Candidate Species-Specific Marker Gene from the Bioinformatical Analysis of Single Exonic Genes.** The analysis of large numbers of genes from multiple species to identify polymorphisms and their subsequent confirmation through either genotyping or sequencing require extensive effort. Use of single exonic or intronless genes could minimize the effort in designing PCR primers for gene amplification. They could also minimize the effort in comparing sequence variations for genes with species-variable exonic structures or lengths.

To expedite the screening process of candidate genes for identifying species-specific genetic markers in vertebrates, we retrieved the DNA sequences of 3750 single exonic genes listed in the SEGE database (18), a database for intronless or single exonic genes in eukaryotes. Next, nucleotide sequences were bioinformatically analyzed using NCBI blast and a multiple sequence alignment program. This was done to find genes with sequence information available from several vertebrate species and well-conserved regions containing possible species-specific SNPs. From our analysis, only *ZNF238* met the criteria (data not shown). However, from further analysis on *ZNF238*, we found that *ZNF238* consisted of two exons, one large exon for the entire protein coding region and one small exon for part of the 5' untranslated region, suggesting that the bioinformatic prediction of gene structures on a genome scale is challenging.

Determination of *ZNF238* DNA Sequences from 15 Previously Unreported Species. Because the depth of the existing sequence information for *ZNF238* in the database was shallow, we designed a specific primer set to amplify the conserved regions of *ZNF238* to obtain more sequence information from other species. The nucleotide alignment of five different species in the NCBI database allowed us to identify highly conserved regions to design PCR primers. The PCR reaction yielded the expected 823 bp products from 20 different species. PCR products from 2 to 13 individuals per species depending on the availability of samples were subjected to DNA sequence determination. The results were identical to the previously reported sequences in Genbank, except for previously unreported sequences of 15 species, *N. procyonoides*, *S. scrofa*, *L. brachyurus*, *H. inermis*, *C. hircus*, *C. capreolus*, *C. taiouanus*, *O. aries*, *M. rufus*, *A. platyrhynchos*, *P. colchicus*, *A. monachus*, *F. japonensis*, *L. africana*, and *H. amphibius* (Supporting Information), which were deposited into Genbank with accession numbers FJ799906–FJ799909, GU045454–GU045465, and GU082393–GU082394. Interestingly, the DNA sequence information from wild boars (*Sus scrofa coreanus*) and wolves (*Canis lupus chanco*) was identical to that of subspecies pigs (*Sus scrofa domesticus*) and dogs (*Canis lupus familiaris*), respectively.


In general, the usefulness of DNA markers in forensic science, food safety, and investigation of illegal wildlife traffic requires the successful analysis of the markers in low-copy and low-quality DNA samples. Therefore, further methodological adjustments for the use of *ZNF238* may be necessary to increase the usefulness of this marker in applications with field samples.

Species Discrimination by PCR-RFLP Using *HhaI* and *HpyCH4IV*. To evaluate the presence of species-specific polymorphisms of *ZNF238* and to identify possible restriction enzyme sites for PCR-RFLP, we performed in silico restriction mapping analysis for the 823 bp region of *ZNF238* against all possible restriction enzyme sites. To confirm the in silico results and to evaluate the possible presence of intraspecies variation, we performed PCR-RFLP analysis on 10 to two unrelated animals for each species. When the PCR amplicons were digested with *HhaI* restriction enzyme, eight RFLP patterns were distinctly discriminated (Table 1; Figure 1). *HhaI* generated the restriction patterns of type a in *C. familiaris* and *N. procyonoides*; b in *S. scrofa*; c in *M. musculus*; d in *E. caballus*; e in *L. brachyurus* and *L. africana*; f in *B. taurus*, *H. inermis*, *C. hircus*, *C. capreolus*, *C. taiouanus*, and *O. aries*; g in *M. rufus*, *A. platyrhynchos*, *A. monachus*, and *F. japonensis*; h in *G. gallus* and *P. colchicus*; and i in *H. amphibius*.

To distinguish the unidentifiable species by *HhaI*, *HpyCH4IV* was used (Table 2; Figure 2). *HpyCH4IV* yielded fragment patterns of type e in *C. familiaris*, *N. procyonoides*, and *E. caballus*; b in *M. musculus*; c in *L. brachyurus*; a in *S. scrofa*, *B. taurus*,


Table 1. *HhaI* Restriction Patterns Predicted for the Nucleotide Sequence of the *ZNF238* Conserved Region for 20 Vertebrate Species

RFLP type	<i>HhaI</i> restriction patterns (bp)
a	249 165 126 105 97 48 33
b	310 168 105 88 81 38 33
c	559 138 88 38
d	559 105 88 38 33
e	559 138 126
f	478 105 88 81 38 33
g	823
h	697 126
i	478 186 88 38 33

**Figure 1.** PCR-RFLP patterns for *ZNF238* obtained by digestion with *HhaI*. The symbols "a" to "i" on the top of the picture correspond to the PCR-RFLP types of *HhaI* in **Table 1**. DNA fragments with sizes of <100 bp were not clearly shown. M, size marker.**Table 2.** *HpyCH4IV* Restriction Patterns Predicted for the Nucleotide Sequence of the *ZNF238* Conserved Region for 20 Vertebrate Species

RFLP type	<i>HpyCH4IV</i> restriction patterns (bp)
a	285 230 144 66 63 29 6
b	285 230 158 84 66
c	285 230 150 84 66 8
d	429 230 92 66 6
e	285 230 144 92 66 6
f	429 285 230 144 92 66 6
g	628 129 66
h	556 267
i	351 267 205
j	435 322 66

H. inermis, *C. hircus*, *C. capreolus*, *C. taiouanus*, and *H. amphibius*; g in *M. rufus*; h in *A. platyrhynchos*; i in *G. gallus*, *P. colchicus*, *A. monachus*, and *F. japonensis*; and j in *L. africana*. For *O. aries*, we identified a C-to-T polymorphism at cDNA nucleotide position 1101 (Supporting Information) through the process of experimental validation, resulting in three different *HpyCH4IV* RFLP patterns. The patterns of homozygotes for alleles A and B were types d and e, respectively. Heterozygotes for alleles A and B showed a banding pattern of type f (**Table 2**). We also find allelic variation for *N. procyonoides* and *F. japonensis* at nucleotide positions 1122 and 1257, respectively (Supporting Information). However, these polymorphisms were not associated with any species specificity. The results from PCR-RFLP

**Figure 2.** PCR-RFLP patterns for *ZNF238* obtained by digestion with *HpyCH4IV*. The symbols "a" to "j" on the top of the picture correspond to the PCR-RFLP types of *HpyCH4IV* in **Table 2**. DNA fragments with sizes of <100 bp were not clearly shown. M, size marker.**Table 3.** *HhaI* and *HpyCH4IV* Restriction Patterns for *ZNF238* DNA Sequence Analysis for 20 Vertebrate Species

species	<i>HhaI</i>	<i>HpyCH4IV</i>
<i>Canis lupus familiaris</i>	a	e
<i>Nyctereutes procyonoides</i>	a	e
<i>Sus scrofa</i>	b	a
<i>Mus musculus</i>	c	b
<i>Equus caballus</i>	d	e
<i>Lepus brachyurus</i>	e	c
<i>Bos taurus</i>	f	a
<i>Hydropotes inermis</i>	f	a
<i>Capra aegagrus hircus</i>	f	a
<i>Capreolus capreolus</i>	f	a
<i>Cervus nippon taiouanus</i>	f	a
<i>Ovis aries A</i>	f	d
<i>Ovis aries B</i>	f	e
<i>Ovis aries A/B</i>	f	f
<i>Macropus rufus</i>	g	g
<i>Anas platyrhynchos</i>	g	h
<i>Gallus gallus</i>	h	i
<i>Phasianus colchicus</i>	h	i
<i>Aegypius monachus</i>	g	i
<i>Falco peregrinus japonensis</i>	g	i
<i>Loxodonta africana</i>	e	j
<i>Hippopotamus amphibius</i>	i	a

experiments and the predictions from in silico analyses were identical.

However, both *HhaI* and *HpyCH4IV* RFLP patterns were identical among the closely related species such as dogs and raccoons; cattle, deer (*H. inermis*, *C. capreolus*, and *C. taiouanus*) and goats; chickens and pheasants; and vultures and falcons. Therefore, we were able to distinguish 13 animal species groups, including major domestic animals, using two restriction enzymes against the conserved region of *ZNF238* (**Table 3**).

Complete Species Discrimination Using Single Nucleotide Polymorphisms (SNPs) from the Conserved Region of *ZNF238*. To differentiate the undistinguishable species by *HhaI* and *HpyCH4IV* RFLP patterns, we analyzed SNPs for the conserved region of *ZNF238* (nt771–1549) among 21 different species, including the 20 species in this study as well as *H. sapiens* from a public database (Supporting Information). The dog and raccoon

Table 4. Differentiation of Closely Related Species Using Single Nucleotide Polymorphisms

species	(A)				
	nucleotide position				
	1032		1326		
<i>Canis lupus familiaris</i>	C		G		
<i>Nyctereutes procyonoides</i>	T		A		

species	(B)				
	nucleotide position				
	1014	1051	1275	1440	1479
<i>Bos taurus</i>	A	A	G	C	T
<i>Hydropotes inermis</i>	C	G	G	T	C
<i>Capreolus capreolus</i>	C	G	G	C	C
<i>Cervus nippon taiouanus</i>	A	A	C	C	C
<i>Capra aegagrus hircus</i>	A	A	G	T	C

species	(C)			
	nucleotide position			
	1116	1167		1311
<i>Gallus gallus</i>	T	C		C
<i>Phasianus colchicus</i>	C	T		T

species	(D)				
	nucleotide position				
	921	954	1020	1455	1542
<i>Aegypius monachus</i>	A	T	G	G	G
<i>Falco peregrinus japonensis</i>	G	C	A	A	A

were distinguished by nucleotide positions 1032 and 1326 (**Table 4A**). The cattle, deer, and goat group was separated into each species on the basis of five SNPs at nucleotide positions 1014, 1051, 1275, 1440, and 1479 (**Table 4B**). Chickens and pheasants had three nucleotide differences at positions 1116, 1167, and 1311 (**Table 4C**). Vultures and falcons showed five nucleotide differences at positions 921, 954, 1020, 1455, and 1542 (**Table 4D**). Therefore, we were able to distinguish 21 vertebrate species compared in this study by analyzing the DNA sequence polymorphisms of the conserved region of *ZNF238*.

We also analyzed the average sequence similarity of the conserved region of *ZNF238* (nt771–1549) among the species used in this study. The amino acid and nucleotide sequence similarities from pairwise comparisons between humans and 20 other species for the conserved region of *ZNF238* (excluding *X. laevis*, which is distantly related) were >96.8 and >87.9%, respectively (**Table 5**). Even the amino acid sequence similarity between humans and distantly related *X. laevis* was 82.2%. These results indicate that sequence conservation of the region selected for species discrimination may be very important for *ZNF238* function. This result may suggest that the region can be applied to other euteleostomi species that were not tested in this study. Accordingly, we did not find any inconsistent DNA polymorphisms among different breeds of animals within a species or even between subspecies such as *Sus scrofa domestica* and *Sus scrofa coreanus* and *Canis lupus familiaris* and *Canis lupus chanco*, indicating the usefulness of *ZNF238* for species discrimination (Supporting Information).


Species Identification from DNA Mixtures of Multiple Species.

In addition to species identification using single species samples, we evaluated species determination from DNA mixtures of multiple species at different concentrations. DNA of *G. gallus*, *E. caballus*, and *B. taurus* was combined at ratios of 1:1:1, 2:1:1, 10:1:1, and 50:1:1, and the accuracy of species determination was

Table 5. Amino Acid and Nucleotide Sequence Similarities for the Conserved Regions of *ZNF238* Used for Species Discrimination between Humans and Other Species^a

species	accession no.	ID (aa, %)	ID (nt, %)
<i>Mus musculus</i>	NM_001012330.1	99.6	95.4
<i>Lepus brachyurus</i>	GU045456	99.2	96.8
<i>Bos taurus</i>	NM_001078016.1	98.8	95
<i>Ovis aries</i>	FJ799908	98.8	95
<i>Cervus nippon taiouanus</i>	GU045460	98.8	95
<i>Capra aegagrus hircus</i>	GU045458	98.8	95
<i>Hippopotamus amphibius</i>	GU082394	98.8	95
<i>Equus caballus</i>	XM_001492205.2	98.1	95.6
<i>Capreolus capreolus</i>	GU045459	98.1	94.9
<i>Hydropotes inermis</i>	GU045457	98.1	94.7
<i>Sus scrofa</i>	GU045465	97.7	94.9
<i>Loxodonta africana</i>	GU082393	97.7	93.8
<i>Macropus rufus</i>	FJ799907	96.9	88.5
<i>Aegypius monachus</i>	GU045462	96.9	88.3
<i>Falco peregrinus japonensis</i>	GU045463–4	96.9	87.9
<i>Gallus gallus</i>	XM_426137.2	96.5	88.6
<i>Phasianus colchicus</i>	GU045461	96.5	88.3
<i>Anas platyrhynchos</i>	FJ799906	96.5	88.2
<i>Canis lupus familiaris</i>	XM_547497.2	95.8	93.5
<i>Nyctereutes procyonoides</i>	GU045454–5	95.8	93.5
<i>Xenopus laevis</i>	BC048019.1	82.2	75.8

^a The region from amino acid 258 to 516 (nucleotides 771–1549, the numbering starts from the translation start codon of NM_205768.2) of the conserved region of *ZNF238* was used for comparisons after removal of the sequences included in primers.

**Figure 3.** PCR-RFLP patterns of DNA from multiple species (*G. gallus*, *B. taurus*, *E. caballus*) with various amounts of DNA: lanes 1 and 2, 1:1:1; lanes 3 and 4, 2:1:1; lanes 5 and 6, 10:1:1; lanes 7 and 8, 50:1:1. DNA fragments with sizes of <100 bp were not clearly shown. M, size marker.

evaluated by both PCR-RFLP and direct sequencing methods. The band intensity for *G. gallus* was slightly stronger than that of other species, possibly due to a higher PCR amplification efficiency. We identified DNA banding patterns representing each species on the gel after PCR-RFLP, even at the 50:1:1 ratio (**Figure 3**).

We also attempted direct sequencing of the PCR products to more informatively present the species identification results using the same DNA mixtures. However, these results were clear only at the 10:1:1 ratio in multiple species DNA. DNA peaks from species at <10% of the total DNA amount (such as at 50:1:1) were not clearly identifiable (**Figure 4**). Although direct




Figure 4. Sequencing chromatogram of the *ZNF238* conserved region from DNA mixtures consisting of three species (*G. gallus*, *B. taurus*, *E. caballus*) with various amounts of DNA. The total amount of template DNA per each reaction was about 50 ng. Numbers at the top of the figure indicate nucleotide positions in cDNA. Polymorphic regions have either double or triple peaks. The nucleotide position 1005 distinguishes all three species.

sequencing-based species identification using *ZNF238* was affected more significantly by differences in DNA concentration in the case of mixed DNA, it may be the more accurate method for interpreting experimental results, considering the presence of several species-specific SNPs that do not form any restriction sites (Supporting Information).

In this paper, we developed PCR-RFLP and direct sequencing methods for species identification of bony vertebrates using a new nuclear DNA marker, *ZNF238*. We showed that the conserved region of the gene can be used specifically as a marker for species identification. Because we only experimentally tested 20 species plus 2 subspecies for multiple animals, additional species should be experimentally evaluated to expand the usability of *ZNF238* as a simple and efficient nuclear DNA marker for species identification. The use of *ZNF238* together with mitochondrial analysis can provide more information and higher accuracy in species identification than relying on the results using mtDNA alone.

ACKNOWLEDGMENT

We thank Dr. Sakharkar at the National University of Singapore for providing SEGE data.

Supporting Information Available: Result of multiple sequence alignment for the *ZNF238* conserved region among 21 species plus 2 subspecies of animals. The human sequence was used as a reference sequence. Nucleotides identical with the reference are indicated with a dot. Numbers indicate the nucleotide positions in cDNA (NM_205768.2). Two alleles were identified in *Nyctereutes procyonoides*, *Ovis aries*, and *Falco peregrinus japonensis*. This material is available free of charge via the Internet at <http://pubs.acs.org>.

LITERATURE CITED

- (1) De Ley, J.; Cattoir, H.; Reynaerts, A. The quantitative measurement of DNA hybridization from renaturation rates. *Eur. J. Biochem.* **1970**, *12*, 133–142.

- (2) Johnson, J. L. Use of nucleic-acid homologies in the taxonomy of anaerobic bacteria. *Int. J. Syst. Bacteriol.* **1973**, *23*, 308–315.
- (3) Shneyer, V. S. On the species-specificity of DNA: fifty years later. *Biochemistry (Moscow)* **2007**, *72*, 1377–1384.
- (4) Robin, E. D.; Wong, R. Mitochondrial DNA molecules and virtual number of mitochondria per cell in mammalian cells. *J. Cell Physiol.* **1988**, *136*, 507–513.
- (5) Brown, W. M.; George, M., Jr.; Wilson, A. C. Rapid evolution of animal mitochondrial DNA. *Proc. Natl. Acad. Sci. U.S.A.* **1979**, *76*, 1967–1971.
- (6) Bataille, M.; Crainic, K.; Leterreux, M.; Durigon, M.; de Mazancourt, P. Multiplex amplification of mitochondrial DNA for human and species identification in forensic evaluation. *Forensic Sci. Int.* **1999**, *99*, 165–170.
- (7) Dawney, N.; Ogden, R.; McEwing, R.; Carvalho, G. R.; Thorpe, R. S. Validation of the barcoding gene COI for use in forensic genetic species identification. *Forensic Sci. Int.* **2007**, *173*, 1–6.
- (8) Montiel-Sosa, J. F.; Ruiz-Pesini, E.; Montoya, J.; Roncalés, P.; López-Pérez, M. J.; Pérez-Martos, A. Direct and highly species-specific detection of pork meat and fat in meat products by PCR amplification of mitochondrial DNA. *J. Agric. Food Chem.* **2000**, *48*, 2829–2832.
- (9) Rodríguez, M. A.; García, T.; González, I.; Asensio, L.; Mayoral, B.; López-Calleja, I.; Hernández, P. E.; Martín, R. Identification of goose, mule duck, chicken, turkey, and swine in foie gras by species-specific polymerase chain reaction. *J. Agric. Food Chem.* **2003**, *51*, 1524–1529.
- (10) Ono, K.; Satoh, M.; Yoshida, T.; Ozawa, Y.; Kohara, A.; Takeuchi, M.; Mizusawa, H.; Sawada, H. Species identification of animal cells by nested PCR targeted to mitochondrial DNA. *In Vitro Cell Dev. Biol. Anim.* **2007**, *43*, 168–175.
- (11) Pérez, M.; Vieites, J. M.; Presa, P. ITS1-rDNA-based methodology to identify world-wide hake species of the genus *Merluccius*. *J. Agric. Food Chem.* **2005**, *53*, 5239–5247.
- (12) Groth, J. G.; Barrowclough, G. F. Basal divergences in birds and the phylogenetic utility of the nuclear RAG-1 gene. *Mol. Phylogenet. Evol.* **1999**, *12*, 115–123.
- (13) Hopwood, A. J.; Fairbrother, K. S.; Lockley, A. K.; Bardsley, R. G. An actin gene-related polymerase chain reaction (PCR) test for identification of chicken in meat mixtures. *Meat Sci.* **1999**, *53*, 227–231.
- (14) Ericson, P. G.; Envall, I.; Irestedt, M.; Norman, J. A. Inter-familial relationships of the shorebirds (Aves: Charadriiformes) based on nuclear DNA sequence data. *BMC Evol. Biol.* **2003**, *3*, 16.
- (15) Prychitko, T. M.; Moore, W. S. The utility of DNA sequences of an intron from the beta-fibrinogen gene in phylogenetic analysis of woodpeckers (Aves: Picidae). *Mol. Phylogenet. Evol.* **1997**, *8*, 193–204.
- (16) Aoki, K.; Meng, G.; Suzuki, K. RP58 associates with condensed chromatin and mediates a sequence-specific transcriptional repression. *J. Biol. Chem.* **1998**, *273*, 26698–26704.
- (17) Miller, S. A.; Dykes, D. D.; Polesky, H. F. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res.* **1988**, *16*, 1215.
- (18) Sakharkar, M. K.; Kanguane, P.; Petrov, D. A.; Kolaskar, A. S.; Subbiah, S. SEGE: a database on 'intron less/single exonic' genes from eukaryotes. *Bioinformatics* **2002**, *18*, 1266–1267.

Received for review October 22, 2009. Revised manuscript received December 17, 2009. Accepted December 31, 2009. This work was supported by grants from the BioGreen 21 program (20070401-034-029-01), Rural Development Administration, Republic of Korea and Korea Institute of Planning and Evaluation for Technology of Food, Agriculture, Forestry and Fisheries. Rui Xiao gratefully acknowledges the support of the KU-Brain pool program of Konkuk University, Republic of Korea.