

## Using Neural Networks to Estimate the Losses of Ascorbic Acid, Total Phenols, Flavonoid, and Antioxidant Activity in Asparagus during Thermal Treatments

HONGFEI LU,<sup>\*,†</sup> HONG ZHENG,<sup>†</sup> HEQIANG LOU, LINGLING JIANG,  
 YONG CHEN, AND SHUANGSHUANG FANG

College of Chemistry and Life Science, Zhejiang Normal University, Jinhua, China 321004.

<sup>†</sup>These authors contributed equally to this work.

Artificial neural networks (ANNs) with back-propagation algorithm were developed to predict the percentage loss of ascorbic acid, total phenols, flavonoid, and antioxidant activity in different segments of asparagus during water blanching at temperatures ranging from 65 to 95 °C as a function of blanching time and temperature. In this study, the one-hidden-layer ANNs are used, and the number of neurons in the hidden layer were chosen by trial and error. Optimized ANN models were developed for predicting nutrient losses in bud, upper, middle, and butt segments of asparagus. ANN models were then tested against an independent data set. Our results showed that the predicted values of the correlation coefficients between experimental and ANNs ranged from 0.8166 to 0.9868. Therefore, ANNs could be potential tools for the prediction of nutrient losses in vegetables during thermal treatments.

**KEYWORDS:** *Asparagus officinalis* L.; ANN; ascorbic acid; total phenols; flavonoid; scavenging activity; thermal treatments

### INTRODUCTION

Asparagus (*Asparagus officinalis* L.) is a green vegetable with high antioxidant activity among the commonly consumed vegetables (1). The antioxidants in asparagus include ascorbic acid, flavones, phenols, glutathione, etc. (2). Antioxidants can scavenge free radicals and protect the human body from oxidative stress, which is the main cause of some cancers and heart diseases (3). However, asparagus is an easily perishable vegetable characterized by a limited postharvest life, mainly because of its high respiratory activity, which continues after harvesting. Therefore, it is preserved by canning, pickling, freezing, and drying. Blanching, a thermal process preceding freezing or canning of vegetables, is necessary to inactivate shelf-life-limiting enzymes and to exhaust gas from the plant tissue (4). Hot water blanching is by far the most popular and commercially adopted process as it is the simplest and most economical technique. Blanching, however, has some adverse effects, such as pigment modifications, tissue softening, and nutrient losses. Many researchers have observed the dramatic blanching effect on the degradation of vegetables nutrient content and antioxidant properties (5–8). As far as we know, by taking into account these recent works, information regarding the mathematical model to predict the nutrient content and antioxidant properties of vegetables during thermal treatments is still scarce.

The artificial neural network (ANN) is a set of mathematical methods, often encompassed with artificial intelligence, which in

some way attempts to mimic the functioning of the human brain (9). In the last two decades, ANN has revealed its huge potential in many areas of science and engineering (10–13) because of its exceptional function of self-organizing, self-study, fault tolerance, and high robustness. Recently, interest in using neural networks as a modeling tool in agriculture and food technology is also increasing. It has been successfully used in several food-processing technologies such as drying technologies (14, 15), postharvest technology (16), food rheology (17), microbial predictions (18), fermentation (19), and thermal processing (20, 21).

Therefore, the aim of this study was to evaluate the percentage loss of ascorbic acid, total phenols, flavonoid, and antioxidant activity of different segments of green asparagus during blanching in water through mathematical modeling based on the ANN. The results will help to define optimal blanching conditions for maximum quality retention, and the success of this research will provide the food industries with modeling and simulation for nutrient loss control in vegetables during thermal treatments.

### MATERIALS AND METHODS

**Sample Preparation and Blanching Process.** Fresh asparagus (*Asparagus officinalis* L. var. Grande) was harvested from a local farm in Jinhua (Zhejiang, P.R. China) and transported by refrigeration at 8 °C for 30 min to the laboratory. Spears of the same diameter (0.8–1.0 cm) at the base and length (20 cm) were used in our experiment. The spears, after being sorted for size and length, were washed with tap water and drained. Sixty asparagus spears were placed in distilled water baths set at 60 °C, 65 °C, 70 °C, 75 °C, 80 °C, 85 °C, 90 °C, and 95 °C. Nine spears were removed at the time intervals listed in **Table 1** and immediately cooled in


\*To whom correspondence should be addressed. Tel: +86-0579-8228-2284. E-mail: luhongfei63@yahoo.com.cn.

running water at 15 °C. After cooling, nine spears were randomly divided into three groups (three spears per group). Prior to ascorbic acid, total phenols, flavonoid, and scavenging capacity measurements, asparagus spears at room temperature were cut into four segments (bud, upper, middle, and butt segments) as shown in **Figure 1**. All experiments were repeated three times (one time per group), and the results were reported as average. In addition, the samples were kept at below 0 °C until analysis was done to prevent nutrient losses.

**Chemicals and Reagent.** 2,2'-Diphenyl-1-picrylhydrazyl (DPPH) free radical (purity ≥90%) and rutin (purity ≥95%) were of analytical grade and purchased from Sigma-Aldrich (St. Louis, MO, USA). Gallic acid (purity ≥98%), ascorbic acid (purity ≥99%), Folin–Ciocalteu reagent (purity ≥99%), and 2,6-dichlorophenol indophenol (purity ≥99%) were purchased from Shanghai Sangon Biological Engineering Technology & Services Co., Ltd. (Shanghai, China).

**Ascorbic Acid (AA) Determination.** The AA content in different segments of asparagus was determined upon the basis of quantitative discoloration of the 2,6-dichlorophenol indophenol titrimetric method as described in AOAC (22). Results of AA content were expressed as milligram ascorbic acid per 100 mL juice. The AA content was measured in triplicate.

**Total Flavonoid (TF) Content Determination.** The TF content was evaluated by a colorimetric assay according to the method of Bonvehi et al. (23), with modifications. Briefly, 0.5 mL solution of each plant extract



**Figure 1.** Specimens for measuring ascorbic acid, total phenols, flavonoid, and DPPH radical scavenging activity.

**Table 1.** Blanching Time and Temperature for Fresh Asparagus

temperature (°C)	blanching time (min)			
60	40	80	120	160
65	35	70	105	140
70	30	60	90	120
75	25	50	75	100
80	20	40	60	80
85	15	30	45	60
90	10	20	30	40
95	5	10	15	20

in methanol was separately mixed with 1.5 mL of methanol, 0.1 mL of 2% aluminum chloride, 0.1 mL of 1 M potassium acetate, and 2.8 mL of distilled water, and left at room temperature for 30 min. The absorbance of the reaction mixture was measured at 415 nm using a spectrophotometer. A standard curve ( $y = 0.4564x$ ;  $R^2 = 0.9942$ ;  $SD = 0.0046$ ) was prepared with rutin. The TF content was measured in triplicate and expressed as rutin equivalents in  $\text{mg g}^{-1}$  dry weight of asparagus.


**Total Phenolic (TP) Content Determination.** The TP content was determined colorimetrically using Folin–Ciocalteu reagent, as described by Emmons, Peterson, and Paul (24), with modifications. Total phenolic assay was conducted by mixing 8.3 mL of deionized water, 0.5 mL of extracts, 0.7 mL of 20%  $\text{Na}_2\text{CO}_3$ , and 0.5 mL of Folin–Ciocalteu reagent. After 40 min of reaction in a water bath at 40 °C, the absorbance at 755 nm was measured using a spectrophotometer. A standard curve ( $y = 0.0903x - 0.0011$ ;  $R^2 = 0.9969$ ;  $SD = 0.0015$ ) was prepared with gallic acid. Final results were expressed as gallic acid equivalents (GAE)  $\text{mg/g}$  of dry asparagus. The TP content was measured in triplicate.

**DPPH Radical Scavenging Activity (SA) Determination.** The free radical scavenging activity of plant extracts was evaluated using the stable radical DPPH as described by Maisuthisakul et al. (25). DPPH radical in methanol (5 mM) was prepared, and this solution (100  $\mu\text{L}$ ) was added to sample solutions in methanol (4.9 mL). After 30 min in the dark, absorbance was measured at 517 nm. SA was measured in triplicate samples. The percentage of DPPH radical scavenging activity of each plant extract was calculated using the following equation:


$$\text{Scavenging activity (\%)} = \left[ \frac{A_0 - (A_1 - A_s)}{A_0} \right] \times 100 \quad (1)$$

where  $A_0$  is the absorbance of the control solution (containing only DPPH);  $A_1$  is the absorbance of the DPPH solution containing plant extract; and  $A_s$  is the absorbance of the sample extract solution without DPPH.

**ANN Analysis.** The ANN is a mathematical algorithm inspired by studies of the brain and nervous systems in biological organisms, which has the capability of relating the input and output parameters, learning from examples through iteration, without requiring prior knowledge of the relationships between the process variables (26). A feed forward network structure with input, output, and hidden layers was used in this study as shown in **Figure 2**. The input layer consisted of two neurons which corresponded to blanching time and temperature. The output layer had one neuron representing the percentage loss of nutrient (eg, ascorbic acid, total phenols, flavonoid, or antioxidant activity). The neural network can have more than one hidden layer; however, theoretical works have shown that a single hidden layer is sufficient for the neural network to approximate any complex nonlinear function (27, 28). Hence, the one-hidden-layer ANNs are used in this study. The number of neurons within each of these layers varied from 2 to 10. The back-propagation algorithm was utilized in the training of ANN models (27). A hyperbolic tangent sigmoid was used as the transfer function in the hidden layer and output layer.



**Figure 2.** Schematic diagram of artificial neural networks for the prediction of the percentage loss of ascorbic acid, total phenols, flavonoid, and DPPH radical scavenging activity used in this study.



**Figure 3.** Water blanching effect on ascorbic acid (A), total phenols (B), flavonoid (C), and DPPH radical scavenging activity (D) of the bud segment of asparagus.

Minimization of error was accomplished using the Levenberg–Marquardt (LM) algorithm (29, 30). The numerical values of the input and output variables used by the ANN were normalized values in the range 0–1.

Training was based on a supervised method with back-propagation strategy and was finished when the mean square error (MSE) converged and was less than 0.1. If the MSE did not go below 0.1, training was completed after 1,000 epochs, where an epoch represents one complete sweep through all of the data in the training set. The ANN modeling program was designed and programmed under Matlab software, version 6.1.

The prediction performances of the various ANN models were compared using mean squared error (MSE). The MSE was calculated using the following equation:

$$MSE = \frac{1}{N} \sum_{i=1}^N (k_E - k_P)^2 \quad (2)$$

where  $N$  is the total number of data;  $k_P$  represents the predicted output from the neural network model for a given input, while  $k_E$  is the experimental value.

## RESULTS AND DISCUSSION

Blanching is a heat treatment preferably in a wet medium either by steam or hot water which provides uniform heating and high heat transfer rate (4). Water blanching is by far the most popular and commercially adopted process because it is the simplest and most economical method. In addition, the household practice of cooking vegetables is usually in the presence of water. However, water blanching may have some negative effect on vegetable quality, such as excessive loss of texture, unwanted changes in color, and nutritional losses (8, 31–33). In our study, the AA, TP, TF, and SA in green asparagus markedly decreased, depending on the blanching time and temperature. A representative graph for nutritional losses in the bud segment of asparagus is given in

**Table 2.** Learning Data Set<sup>a</sup> Used in the Development of ANNs

temperature (°C)	time (min)	percentage loss (%)			
		AA	TP	TF	SA
60	80	26.32	3.93	13.34	33.41
60	160	38.66	14.25	23.02	51.08
65	35	29.48	4.13	12.52	28.30
65	105	33.68	10.88	15.30	53.41
65	70	32.88	5.52	13.64	39.66
70	30	35.02	5.15	11.97	35.55
70	60	38.72	7.56	14.45	45.50
70	120	51.73	22.91	26.51	65.83
75	25	38.06	11.82	11.45	33.10
75	75	53.47	26.95	24.20	59.57
80	20	36.79	12.43	13.50	27.24
80	40	38.68	18.13	14.27	33.48
80	80	60.74	30.89	34.13	49.66
85	15	37.58	13.96	13.46	25.20
85	45	53.55	28.82	29.21	43.93
85	60	60.42	38.63	38.46	50.33
90	10	41.63	14.90	14.91	29.81
90	30	56.98	27.21	30.95	40.88
90	40	61.72	37.45	35.58	49.17
95	5	46.62	15.02	19.08	33.74
95	15	50.90	20.90	22.11	37.20
95	20	59.40	32.37	31.24	38.14

<sup>a</sup> These data were collected from bud segments of asparagus as representatives.

Figure 3, and the percentage losses of AA, TP, TF, and SA were calculated in the bud segment of asparagus and given in Tables 2–4. In recent years, many authors have developed kinetic models to describe the quality changes in vegetables during blanching (34–38). However, to our knowledge no previous study has been reported on using ANN to predict the percentage

loss of ascorbic acid, total phenols, flavonoid, and antioxidant activity of different segments of green asparagus during blanching in water. So far, the application of neural networks in the food science area has broad coverage: sensory analysis, classifications, microbial predictions, thermal control, etc. (17, 18, 21, 39, 40). The purpose of this study was to prove that the ANN is a convenient and potential tool for the prediction of the quality changes of different segments of green asparagus during water

**Table 3.** Validating Data Set<sup>a</sup> Used in the Development of ANNs

temperature (°C)	time (min)	percentage loss (%)			
		AA	TP	TF	SA
60	40	18.10	3.78	7.14	22.18
65	140	36.01	12.35	16.61	56.84
75	100	54.78	30.79	23.12	61.93
80	60	50.12	21.65	24.47	46.52
90	20	45.10	22.41	20.36	35.68

<sup>a</sup>These data were collected from bud segments of asparagus as representatives.

**Table 4.** Testing Data Set<sup>a</sup> Used in the Development of ANNs


temperature (°C)	time (min)	percentage loss (%)			
		AA	TP	TF	SA
60	120	30.25	8.11	14.72	47.98
70	90	49.12	21.41	25.04	64.90
75	50	36.42	17.56	9.47	47.06
85	30	40.89	22.63	15.44	35.62
95	10	61.01	41.29	33.06	42.65

<sup>a</sup>These data were collected from bud segments of asparagus as representatives.


blanching as a function of blanching time and temperature. The development of the ANN model involves three basic steps: training, validation, and prediction phases.

**Training of the Neural Network.** In our study, 75% of the data were used to generate the model, and validation was performed with 12.5% of the total data. In addition, the remaining 12.5% of the data were used in the prediction phase. **Table 2** shows the training set, which was presented to the network, and a back-propagation algorithm automatically adjusted the weights until the output response to input vector was as close as possible to the desired response (39). The one-hidden-layer ANNs are used in this study because many researchers have shown that a single hidden layer is sufficient to approximate any complex nonlinear function (27, 28). The optimal number of nodes in the hidden layer was selected by using a trial and error method. **Figure 4** shows the changes of MSE in the prediction of the percentage loss of AA, TP, TF, and SA in the bud, upper, middle, and butt segments of asparagus with different numbers of neurons in the hidden layer. Our results showed that the optimal number of nodes in the hidden layer was 5, 5, 7, and 5 in the bud segment of asparagus, 5, 7, 5, and 7 in the upper segment, 8, 4, 6, and 7 in the middle segment, and 5, 5, 5, and 7 in the butt segment, for predicting the percentage loss of AA, TP, TF, and SA, respectively. The above models were developed for asparagus diameters varying from 0.8 to 1.0 cm and temperatures from 65 to 95 °C. These ANN models, however, did not consider temperatures below 65 °C and above 95 °C or diameters of asparagus below 0.8 cm and above 1.0 cm.

**Validation and Prediction Phases.** In validation and prediction phases, we used the selected topology of ANN models with



**Figure 4.** Mean squared error (MSE) in the prediction of the percentage loss of ascorbic acid (□), total phenols (○), flavonoid (▽), and DPPH radical scavenging activity (△) with different numbers of neurons in the hidden layer for bud, upper, middle, and butt segments of asparagus during blanching in water. The arrows signify the optimal number of nodes in the hidden layer.



**Figure 5.** Correlation of experimental and predicted percentage loss of ascorbic acid (A), total phenols (B), flavonoid (C), and DPPH radical scavenging activity (D) with training and validation data sets, as well as the prediction data set for the bud segment of asparagus during water blanching using the optimal ANN.

**Table 5.** Mean Squared Errors (MSE) and the Correlation Coefficients ( $R^2$ ) in the Prediction of the Percentage Loss of AA, TP, TF, and SA in Different (Bud, Upper, Middle, and Butt) Segments of Asparagus with the Optimal ANNs

	AA		TP		TF		SA	
	MSE (%)	$R^2$	MSE (%)	$R^2$	MSE (%)	$R^2$	MSE (%)	$R^2$
bud segment	10.6745	0.9868	51.2112	0.8166	7.6963	0.9088	32.8099	0.9620
upper segment	62.1037	0.8318	30.7957	0.8767	12.7976	0.8234	33.6940	0.8716
middle segment	28.2051	0.9205	44.4763	0.8568	4.6623	0.9780	75.3507	0.8665
butt segment	24.1384	0.9654	14.8777	0.8597	7.3191	0.9862	26.9330	0.9129

the previously adjusted weights. Representative graphs of the correlations between the predicted and experimental values in respect to the training set, validation set, and prediction set are shown in **Figure 5**, together with the full description of equations of the linear regression model. In addition, **Table 5** shows the MSE and the correlation coefficients ( $R^2$ ) between experimental and ANN predicted values in the prediction of the percentage loss of AA, TP, TF, and SA in different (bud, upper, middle, and butt) segments of asparagus with the optimal ANNs. The optimal ANN could predict the percentage loss of AA, TP, TF, and SA in asparagus with the MSE of 10.6745%, 51.2112%, 7.6963%, and 32.8099% for the bud segment, 62.1037%, 30.7957%, 12.7976%, and 33.6940% for the upper segment, 28.2051%, 44.4763%, 4.6623%, and 75.3507% for the middle segment, and 24.1384%, 14.8777%, 7.3191%, and 26.9330% for the butt segment, respectively. In addition, the correlation coefficients between experimental and ANN predicted values ranged from 0.8166 to 0.9868, as shown in **Table 5**. Therefore, the ANN model as a potential tool can be used to determine the percentage loss of AA, TP, TF, and SA during water blanching.

In conclusion, the ANN, a convenient and potential tool, can be a promising method for modeling the nutrient losses of vegetables during water blanching. Therefore, the ANN seems to find application in the quality analysis of functional properties of food of plant origin for the prediction of not only the ascorbic acid, total phenols, flavonoid, and DPPH radical scavenging activity but also for the control of food quality during processing.

#### ABBREVIATIONS USED

AA, ascorbic acid; TP, total phenols; TF, total flavonoid; SA, DPPH radical scavenging activity.

#### ACKNOWLEDGMENT

We sincerely thank Professor Changjiang Zhang in the College of Information Science and Engineering at Zhejiang Normal University (Jinhua, China) for his assistance with the revision of this manuscript.

#### LITERATURE CITED

- (1) Vinson, J. A.; Hao, Y.; Su, X.; Zubik, L. Phenol antioxidant quantity and quality in foods: Vegetables. *J. Agric. Food Chem.* **1998**, *46*, 3630–3634.
- (2) Shao, Y.; Poobrasert, O.; Kennelly, E. J.; Chin, C. K.; Ho, C. T.; Huang, M. T.; Garrison, S. A.; Cordell, G. A. Steroidal saponins from *Asparagus officinalis* and their cytotoxic activity. *Planta Med.* **1997**, *63* (3), 258–262.
- (3) Verlangieri, A. J.; Kapeghian, J. C.; el-Dean, S.; Bush, M. Fruit and vegetable consumption and cardiovascular mortality. *Med. Hypotheses* **1985**, *16* (1), 7–15.
- (4) Lee, F. A. The blanching process. *Adv. Food Res.* **1958**, *8*, 63–109.
- (5) Selman, J. D. Vitamin retention during blanching of vegetables. *Food Chem.* **1994**, *49* (2), 137–147.
- (6) Howard, L. A.; Wong, A. D.; Perry, A. K.; Klein, B. P.  $\beta$ -Carotene and ascorbic acid retention in fresh and processed vegetables. *J. Food Sci.* **1999**, *64* (5), 929–936.
- (7) Murcia, M. A.; López-Ayerra, B.; Garcia-Carmona, F. Effect of processing methods and different blanching times on broccoli: proximate composition and fatty acids. *LWT-Food Sci. Technol.* **1999**, *32* (4), 238–243.
- (8) Oboh, G. Effect of blanching on the antioxidant properties of some tropical green leafy vegetables. *LWT-Food Sci. Technol.* **2005**, *38* (5), 513–517.
- (9) Zupan, J.; Gasteiger, J. Neural networks: a new method for solving chemical problems or just a passing phase? *Anal. Chim. Acta* **1991**, *248*, 1–30.

- (10) Fang, Y. C.; Wu, B. W. Neural network application for thermal image recognition on flow-resolution objects. *J. Opt. A: Pure Appl. Opt.* **2007**, *9* (2), 134–144.
- (11) Parlak, A.; Islamoglu, Y.; Yasar, H.; Egrisogut, A. Application of artificial neural network to predict specific fuel consumption and exhaust temperature for a diesel engine. *Appl. Therm. Eng.* **2006**, *26* (8–9), 824–828.
- (12) Maier, H. R.; Dandy, G. C. Neural networks for the prediction and forecasting water resources variables: a review of modeling issues and applications. *Environ. Model. Soft.* **2000**, *15*, 101–124.
- (13) Yuanyou, X.; Yanming, X.; Ruigeng, Z. An engineering geology evaluation method based on an artificial neural network and its application. *Eng. Geol.* **1997**, *47* (1–2), 149–156.
- (14) Kaminski, W.; Strumillo, P.; Romczak, E. Neurocomputing approaches to modeling of drying process dynamics. *Drying Technol.* **1998**, *16* (6), 967–992.
- (15) Chen, C. R.; Ramaswamy, H. S.; Alli, I. Neural Network Based Optimization of Quality of Osmo-Convective Dried Blueberries. In *FoodSIM' 2000, 1st International Conference on Simulation in Food and Bio Industries*; Nantes, France, June 26–27, 2000; pp 33–35.
- (16) Morimoto, T.; Baerdemaeker, J. De.; Hashimoto, Y. An intelligent approach for optimal control of fruit storage process using neural networks and genetic algorithms. *Comput. Electron. Agric.* **1997**, *18*, 205–224.
- (17) Ruan, R.; Almaer, S.; Zhang, J. Prediction of dough rheological properties using neural networks. *Cereal Chem.* **1995**, *72* (3), 308–311.
- (18) Geeraerd, A. H.; Herremans, C. H.; Cenens, C.; Van Impe, J. G. Application of artificial neural networks as a non-linear modular modeling technique to describe bacterial growth in chilled food products. *Int. J. Food Microbiol.* **1998**, *44*, 49–68.
- (19) Teissier, P.; Perret, B.; Latrille, E.; Barillere, J. M.; Corrieu, G. Hybrid recurrent neural network model for yeast production monitoring and control in a wine base medium. *J. Biotechnol.* **1997**, *55* (2), 157–169.
- (20) Chen, C. R.; Ramaswamy, H. S. A neuro-computing approach for modeling of residence time distribution (RTD) of carrot cubes in a vertical scraped surface heat exchanger (SSHE). *Food Res. Int.* **2000**, *33* (7), 549–556.
- (21) Afaghi, M.; Ramaswamy, H. S.; Prasher, S. O. Thermal process calculations using artificial neural network models. *Food Res. Int.* **2001**, *34*, 55–65.
- (22) AOAC Vitamins and Other Nutrients (Chapter 45). In *Official Methods of Analysis*, 17th ed.; AOAC International: Washington, DC, 2000; pp 16–20.
- (23) Bonvehí, J. S.; Torrent, M. S.; Lorente, E. C. Evaluation of polyphenolic and flavonoid compounds in honey-bee collected pollen produced in Spain. *J. Agric. Food Chem.* **2001**, *49*, 1848–1853.
- (24) Emmons, C. L.; Peterson, D. M.; Paul, G. L. Antioxidant capacity of oat (*Avena sativa* L.) extracts. 2. In vitro antioxidant activity and contents of phenolic and tocol antioxidants. *J. Agric. Food Chem.* **1999**, *47*, 4894–4898.
- (25) Maisuthisakul, P.; Pongsawatmanit, R.; Gordon, M. H. Assessment of phenolic content and free-radical scavenging capacity of some Thai indigenous plants. *Food Chem.* **2007**, *100*, 1409–1418.
- (26) Palancar, M. C.; Aragón, J. M.; Torrecilla, J. S. pH-control system based on artificial neural networks. *Ind. Eng. Chem. Res.* **1998**, *37* (7), 2729–2740.
- (27) Hornik, K.; Stinchcombe, M.; White, H. Multilayer feed forward network are universal approximator. *Neural Network* **1989**, *2*, 359–366.
- (28) Hagan, M. T.; Menhaj, M. B. Training feed forward networks with the Marquardt algorithm. *IEEE Trans. Neural Networks* **1994**, *6*, 861–867.
- (29) Levenberg, K. A method for the solution of certain non-linear problems in least squares. *Quart. J. Appl. Math.* **1944**, *2* (2), 164–168.
- (30) Marquardt, D. W. An algorithm for least-squares estimation of non-linear parameters. *J. Soc. Ind. Appl. Math.* **1963**, *2* (2), 431–441.

- (31) Arroqui, C.; Rumsey, T. R.; Lopez, A.; Virseda, P. Effect of different soluble solids in the water on the ascorbic acid losses during water blanching of potato tissue. *J. Food Eng.* **2001**, *47*, 123–126.
- (32) Volden, J.; Borge, G. I. A.; Bengtsson, G. B.; Hansen, M.; Thygesen, I. E.; Wicklund, T. Effect of thermal treatment on glucosinolates and antioxidant-related parameters in red cabbage (*Brassica oleracea* L. ssp. *capitata* f. *rubra*). *Food Chem.* **2008**, *109*, 595–605.
- (33) Volden, J.; Borge, G. I. A.; Hansen, M.; Wicklund, T.; Bengtsson, G. B. Processing (blanching, boiling, steaming) effects on the content of glucosinolates and antioxidant-related parameters in cauliflower (*Brassica oleracea* L. ssp. *botrytis*). *LWT-Food Sci. Technol.* **2009**, *42*, 63–73.
- (34) Lau, M. H.; Tang, J.; Swanson, B. G. Kinetics of textural and color changes in green asparagus during thermal treatments. *J. Food Eng.* **2000**, *45*, 231–236.
- (35) Tijsskens, L. M. M.; Schijvens, E. P. H. M.; Biekman, E. S. A. Modelling the change in colour of broccoli and green beans during blanching. *Innov. Food Sci. Emerg. Technol.* **2001**, *2*, 303–313.
- (36) Gonçalves, E. M.; Pinheiro, J.; Abreu, M.; Brandão, T. R. S.; Silva, C. L. M. Modelling the kinetics of peroxidase inactivation, colour and texture changes of pumpkin (*Cucurbita maxima* L.) during blanching. *J. Food Eng.* **2007**, *81*, 693–701.
- (37) Liu, E. Z.; Scanlon, M. G. Modeling the effect of blanching conditions on the texture of potato strips. *J. Food Eng.* **2007**, *81*, 292–297.
- (38) Pedreschi, F.; Travisany, X.; Reyes, C.; Troncoso, E.; Pedreschi, R. Kinetics of extraction of reducing sugar during blanching of potato slices. *J. Food Eng.* **2009**, *91*, 443–447.
- (39) Ni, H.; Gunasekaran, S. Food quality prediction with neural networks. *Food Technol.* **1998**, *52* (10), 60–65.
- (40) Guyer, D.; Yang, X. Use of genetic artificial neural networks and spectral imaging for defect detection on cherries. *Comput. Electron. Agric.* **2000**, *29* (3), 179–194.

---

Received for review October 19, 2009. Revised manuscript received January 29, 2010. Accepted January 29, 2010. This work was carried out with financial support from the Postgraduate Innovation Research Projects of Zhejiang Province (YK2008038).