

**Correction to Antihypertensive Effect of Peptide-Enriched Soy Sauce-Like Seasoning and Identification of Its Angiotensin I-Converting Enzyme Inhibitory Substances** [*J. Agric. Food Chem.* 2010, 58, 821. DOI: 10.1021/jf903261h]. Takeharu Nakahara,\* Atsushi Sano, Hitomi Yamaguchi, Katsutoshi Sugimoto, Hiroyuki Chikata, Emiko Kinoshita, and Riichiro Uchida

The unit of ACE inhibitory activity of identified substances in the Abstract, Results and Discussion, and Table 3 should be corrected to ( $IC_{50}$ ,  $\mu M$ ). The correct ACE inhibitory activities are as follows: Ala-Trp,  $IC_{50} = 10 \mu M$ ; Gly-Trp,  $IC_{50} = 30 \mu M$ ; Ala-Tyr,  $IC_{50} = 48 \mu M$ ; Ser-Tyr,  $IC_{50} = 67 \mu M$ ; Gly-Tyr,  $IC_{50} = 97 \mu M$ ; Ala-Phe,  $IC_{50} = 190 \mu M$ ; Val-Pro,  $IC_{50} = 480 \mu M$ ; Ala-Ile,  $IC_{50} = 690 \mu M$ ; Val-Gly,  $IC_{50} = 1100 \mu M$ ; and nicotianamine,  $IC_{50} = 0.26 \mu M$ .

DOI: 10.1021/jf101333w  
Published on Web 04/14/2010

**Correction to Calculating Radiation Exposures during Use of  $^{14}C$ -Labeled Nutrients, Food Components, and Biopharmaceuticals To Quantify Metabolic Behavior in Humans** [*J. Agric. Food Chem.* 2010, 58, 4632. DOI: 10.1021/jf100113c]. Seung-Hyun Kim, Peter B. Kelly, and Andrew J. Clifford\*

The final sentence under the heading Sample Collections (in the Materials and Methods section) should read as follows: When 100 nCi of  $^{14}C$  was dosed to the 70 kg human subject, collected samples had 0.0014 nCi of  $^{14}C/g$ , a level below the U.S. Code of Federal Regulations, Title 10, Section 20.2005, 1991 (cutoff for nonradioactive waste from  $^{14}C$  animal studies,  $\leq 50$  nCi/g or  $1 \mu Ci/year$ ).

DOI: 10.1021/jf101389w  
Published on Web 04/16/2010