

## Galacto-oligosaccharide Synthesis from Lactose Solution or Skim Milk Using the $\beta$ -Galactosidase from *Bacillus circulans*

Barbara Rodriguez-Colinas,<sup>†</sup> Ana Poveda,<sup>§</sup> Jesus Jimenez-Barbero,<sup>‡</sup> Antonio O. Ballesteros,<sup>†</sup> and Francisco J. Plou<sup>\*,†</sup>

<sup>†</sup>Instituto de Catálisis y Petroleoquímica, CSIC, 28049 Madrid, Spain

<sup>§</sup>Servicio Interdepartamental de Investigación, Universidad Autónoma de Madrid, 28049 Madrid, Spain

<sup>‡</sup>Centro de Investigaciones Biológicas, CSIC, 28040 Madrid, Spain

### **S** Supporting Information

**ABSTRACT:** The synthesis of galacto-oligosaccharides (GOS) catalyzed by a novel commercial preparation of  $\beta$ -galactosidase from *Bacillus circulans* (Biolactase) was studied, and the products were characterized by MS and NMR. Using 400 g/L lactose and 1.5 enzyme units per milliliter, the maximum GOS yield, measured by HPAEC-PAD analysis, was 165 g/L (41% w/w of total carbohydrates in the mixture). The major transgalactosylation products were the trisaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc and the tetrasaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc. The GOS yield increased to 198 g/L (49.4% w/w of total carbohydrates) using a higher enzyme concentration (15 U/mL), which minimized the enzyme inactivation under reaction conditions. Using skim milk (with a lactose concentration of 46 g/L), the enzyme also displayed transgalactosylation activity: maximum GOS yield accounted for 15.4% (7.1 g/L), which was obtained at 50% lactose conversion.

**KEYWORDS:** glycosidase, galacto-oligosaccharides, prebiotics, transglycosylation,  $\beta$ -galactosidase, oligosaccharides

### ■ INTRODUCTION

$\beta$ -Galactosidases ( $\beta$ -D-galactoside galactohydrolases, EC 3.2.1.23) catalyze the hydrolysis of the galactosyl moiety from the nonreducing end of various oligosaccharides. These enzymes have attracted attention from dairy industries due to their ability to remove lactose from milk.<sup>1</sup> In addition,  $\beta$ -galactosidases catalyze transgalactosylation reactions in which lactose (as well as the released glucose and galactose) serve as galactosyl acceptors, yielding a series of di, tri-, and higher oligosaccharides called galacto-oligosaccharides (GOS).<sup>2,3</sup> GOS are noncariogenic, reduce the level of cholesterol in serum, prevent colon cancer, and exhibit prebiotic properties. In fact, GOS constitute the major part of oligosaccharides in human milk.<sup>4–6</sup> The properties of GOS depend significantly on their chemical composition, structure, and degree of polymerization.<sup>7</sup> Depending on the origin of  $\beta$ -galactosidase, the yield and composition of GOS vary notably.<sup>8–11</sup> The most studied  $\beta$ -galactosidases are those from *Kluyveromyces lactis*,<sup>12–14</sup> *Rhizopus oryzae*,<sup>15</sup> *Bifidobacterium* sp.,<sup>16</sup> and *Bacillus circulans*.<sup>17</sup>

With regard to the  $\beta$ -galactosidase from *Bacillus circulans*, different isoforms have been reported in the commercial preparation Biolacta (Daiwa Kasei). At least three isoforms with different behaviors in GOS production were characterized:  $\beta$ -galactosidase-1 showed very low transglycosylation activity,<sup>18</sup>  $\beta$ -galactosidase-2 contributed most significantly to GOS synthesis,<sup>18,19</sup> and  $\beta$ -galactosidase-3 was able to produce GOS with  $\beta$ (1 $\rightarrow$ 3) bonds.<sup>20</sup> More recently, Song et al.<sup>21</sup> described four isoforms with different molecular sizes in Biolacta:  $\beta$ -gal-A (189 kDa),  $\beta$ -gal-B (154 kDa),  $\beta$ -gal-C (134 kDa), and  $\beta$ -gal-D (91 kDa). The transferase activity of  $\beta$ -galactosidase from *Bacillus circulans* has been applied to the synthesis of lactosucrose,<sup>22</sup> N-acetyl-lactosamine,<sup>23</sup> and other galactosylated derivatives.<sup>24</sup>

Interestingly, the enzyme is able to catalyze the galactosylation of different acceptors in the presence of organic cosolvents up to 50% v/v.<sup>25</sup> The *Bacillus circulans*  $\beta$ -galactosidase has been also immobilized on different supports.<sup>19,26</sup> However, only partial analysis of the GOS formed in the transglycosylation reaction with lactose has been performed,<sup>19</sup> probably due to the complexity of the reaction mixture derived from the presence of several isoforms with different regiospecificities.

In this work, we have studied the transgalactosylation activity of a novel commercial  $\beta$ -galactosidase preparation from *Bacillus circulans* (Biolactase). A detailed kinetic study of the reaction with lactose was performed, including the structural characterization of the synthesized GOS. It is well reported that oligosaccharides formed by the same monosaccharides with the same anomeric configuration, but differing in the glycosidic bonds between them, exhibit different fermentation patterns.<sup>7</sup>

In addition, skim milk as lactose source was further investigated to assess the effect of lactose concentration on transglycosylation and to explore the in situ formation of GOS in dairy products.

### ■ EXPERIMENTAL PROCEDURES

**Materials.** Biolactase (batch MB-878) is a liquid  $\beta$ -galactosidase preparation from *Bacillus circulans* produced by Kerry Ingredients and Flavours (<http://www.kerry.com>) that was supplied by Biocon (Spain). Glucose, galactose, lactose monohydrate, and *o*-nitrophenyl- $\beta$ -D-galactopyranoside (ONPG) were from Sigma-Aldrich. 3-Galactobiose, 4-galactobiose, 6-galactobiose, 6-*O*- $\beta$ -galactosyl-glucose (allo-

Received: March 20, 2012

Revised: June 7, 2012

Accepted: June 7, 2012

Published: June 7, 2012

lactose), and 4-*O*- $\beta$ -galactosyl-lactose were from Carbosynth (Berkshire, U.K.). Skim milk "Hacendado" was purchased from Mercadono supermarket (Spain). All other reagents and solvents were of the highest available purity and used as purchased.

**Activity Assay.** The enzymatic activity toward ONPG was measured at 40 °C following *o*-nitrophenol release at 405 nm using a microplate reader (Versamax, Molecular Devices). The reaction was started by adding 10  $\mu$ L of the enzyme (conveniently diluted) to 190  $\mu$ L of 15 mM ONPG in 0.1 M sodium acetate buffer (pH 5.5). The increase of absorbance at 405 nm was followed in continuous mode during 5 min. The extinction molar coefficient of *o*-nitrophenol at pH 5.5 was determined (537 M<sup>-1</sup> cm<sup>-1</sup>). One unit (U) of activity was defined as that corresponding to the hydrolysis of 1  $\mu$ mol of ONPG per minute.

**SDS-PAGE.** SDS-PAGE was performed on 8% polyacrylamide gels, and the proteins were stained with colloidal Coomassie Blue (Protoblue Safe, National Diagnostics) diluted with ethanol. The HMW-SDS calibration kit (53–220 kDa) was from GE Healthcare Bio-Sciences. The LMW-SDS calibration kit (15–150 kDa) was from Novagen.

**Thermostability of *Bacillus circulans*  $\beta$ -Galactosidase.** The enzyme (approximately 5.5 U/mL) was incubated at different temperatures (40–60 °C) in 0.1 M sodium acetate buffer (pH 5.5). Aliquots were harvested at different times, and the remaining activity toward ONPG was determined as described above (after convenient dilution of the enzyme).

**Production of Galacto-oligosaccharides from Lactose Solution.** The reaction mixture (20 mL) contained 400 g/L lactose (34.7% w/w) in 0.1 M sodium acetate buffer (pH 5.5). The biocatalyst (Biolactase) was then added to adjust the  $\beta$ -galactosidase activity in the reaction mixture to 1.5 or 15 U/mL. The mixture was incubated at 40 °C in an orbital shaker (Vortemp 1550) at 200 rpm. At different times, 200  $\mu$ L aliquots were harvested from the reaction vessel and mixed with 800  $\mu$ L of 0.4 M Na<sub>2</sub>CO<sub>3</sub> (pH 11.0) to stop the reaction. Samples were filtered using 0.45  $\mu$ m cellulose filters (National Scientific) coupled to eppendorf tubes by centrifugation during 5 min at 6000 rpm. For each sample, two dilutions with water (1:400 and 1:4000) were done for HPAEC-PAD analysis.

**Production of Galacto-oligosaccharides from Skim Milk.** Biolactase was added to skim milk (20 mL) to adjust the  $\beta$ -galactosidase activity in the reaction vessel to 1.5 U/mL. The mixture was then incubated at 40 °C in an orbital shaker (Vortemp 1550) at 200 rpm. At different times, 200  $\mu$ L aliquots were harvested from the reaction vessel and mixed with 800  $\mu$ L of 0.4 M Na<sub>2</sub>CO<sub>3</sub> (pH 11.0) to stop the reaction. Samples were filtered, conveniently diluted, and analyzed as described elsewhere.

**HPAEC-PAD Analysis.** Product analysis was carried out by high-performance anion-exchange chromatography coupled with pulsed amperometric detection (HPAEC-PAD) on an ICS3000 Dionex system consisting of an SP gradient pump, an electrochemical detector with a gold working electrode and Ag/AgCl as reference electrode, and an autosampler (model AS-HV). All eluents were degassed by flushing with helium. A pellicular anion-exchange 4  $\times$  250 mm Carbo-Pack PA-1 column (Dionex) connected to a 4  $\times$  50 CarboPac PA-1 guard column was used at 30 °C. Eluent preparation was performed with Milli-Q water and 50% (w/v) NaOH (Sigma-Aldrich). The initial mobile phase was 15 mM NaOH at 1.0 mL/min for 28 min. A gradient from 15 to 200 mM NaOH was performed in 7 min at 1.0 mL/min, and 200 mM NaOH was maintained for 25 min. To increase the sensitivity of the detector at low NaOH concentrations, a Dionex PC10 postcolumn delivery system with 0.2 M NaOH was used at 96 psi. The peaks were analyzed using Chromeleon software. Identification of the different carbohydrates was done on the basis of commercially available standards or purified in our laboratory as described in a previous paper.<sup>11</sup>

**Purification of GOS by Semipreparative HILIC.** For the isolation of unknown GOS in the mixture, the reaction was stopped when GOS yield reached the maximum value. The biocatalyst (Biolactase) was inactivated by boiling the solution for 10 min. The reaction mixture was filtered, and the solution was purified by


semipreparative hydrophilic interaction chromatography (HPLC-HILIC). A quaternary pump (Delta 600, Waters) coupled to a LiChrospher-NH<sub>2</sub> column (5  $\mu$ m, 10  $\times$  250 mm, Merck) was used. The column temperature was kept constant at 30 °C. Acetonitrile/water 90:10 (v/v), degassed with helium, was used as mobile phase (flow rate = 6.25 mL/min) for 8 min. Then, a gradient to acetonitrile/water 80:20 (v/v) was performed in 3 min, and this eluent was maintained during 6 min. Finally, a gradient from the latter mobile phase to acetonitrile/water 75:25 (v/v) was performed in 3 min and maintained for 15 min. Peaks were detected using an evaporative light-scattering detector DDL-31 (Eurosep) equilibrated at 60 °C. A three-way flow splitter (model Accurate, Dionex) and a fraction collector II (Waters) were employed. The fractions containing the main peaks were pooled, and the solvent was eliminated by rotary evaporation.

**Mass Spectrometry.** Samples were analyzed by MALDI-TOF mass spectrometry (Bruker, model Ultraflex III TOF-TOF) using 2,5-dihydroxybenzoic acid doped with sodium iodide as matrix, in positive reflector mode.

**Nuclear Magnetic Resonance (NMR).** The structure of the oligosaccharides was elucidated using a combination of 1D (<sup>1</sup>H, <sup>13</sup>C) and 2D (COSY, TOCSY, NOESY, HSQC) NMR techniques (see the Supporting Information). The spectra of the samples, dissolved in deuterated water (ca. 10 mM), were recorded on a Bruker Avance DRX500 spectrometer equipped with a tunable broadband <sup>1</sup>H/X probe with a gradient in the Z axis, at a temperature of 298 K. Chemical shifts were expressed in parts per million with respect to the 0 ppm point of DSS, used as internal standard. COSY, TOCSY, NOESY, and HSQC standard pulse sequences were provided by Bruker. COSY, TOCSY (80 ms mixing time), and NOESY (500 ms mixing time) experiments were performed with a minimum of 8, 16, and 48 scans, respectively, with 256 increments in the indirect dimension and with 1024 points in the acquisition dimension. The spectral widths were 9 ppm in both dimensions. The HSQC experiment (16 scans) also used 256 increments in the indirect dimension with 1024 points in the acquisition dimension. The spectral width was 120 ppm in the indirect dimension and 9 ppm in the acquisition one.


## RESULTS AND DISCUSSION

### GOS Specificity of *Bacillus circulans* $\beta$ -Galactosidase. A novel commercial preparation of $\beta$ -galactosidase from *Bacillus*




**Figure 1.** SDS-PAGE of  $\beta$ -galactosidase from *B. circulans* (Biolactase). Lanes: 1, marker proteins (53–220 kDa); 2, Biolactase diluted 1:200; 3, Biolactase diluted 1:1000; 4, marker proteins (15–150 kDa).

*circulans* (Biolactase) was studied. SDS-PAGE gel (Figure 1) showed the presence in Biolactase of different proteins with molecular masses in the range of 75–200 kDa, which according to Song et al.<sup>21</sup> makes probable the existence of various  $\beta$ -galactosidase isoforms in the enzyme preparation. This was also confirmed by native PAGE with 4-methylumbelliferyl- $\beta$ -D-galactopyranoside, which showed various active bands in that range (data not shown). The volumetric activity of Biolactase



**Figure 2.** HPAEC-PAD analysis of the reaction of lactose with *Bacillus circulans*  $\beta$ -galactosidase (Biolactase). Peaks: 1, galactose; 2, glucose; 3, allolactose; 4, lactose; 5, 4-galactobiose; 6, 6-galactosyl-lactose; 7, 3-galactosyl-glucose; 10, 4-galactosyl-lactose; 14, Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 3)-Glc; 16, Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc; (8, 9, 11–13, 15) other GOS (unknown). The chromatogram corresponds to the reaction mixture after 77.5 h with Biolactase.



**Figure 3.** 2D-NMR heteronuclear single-quantum coherence (HSQC) analysis of two galacto-oligosaccharides obtained in the reaction of lactose with *Bacillus circulans*  $\beta$ -galactosidase: (A) tetrasaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc; (B) trisaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 3)-Glc. Only the most relevant signals are assigned and labeled.

toward ONPG was 2740 U/mL. Protein concentration was 15.4 mg/mL, and the specific activity accounted for 18.3 U/mg protein. We studied in detail the synthesis of GOS catalyzed by Biolactase using 400 g/L lactose and 1.5 U/mL ( $\beta$ -galactosidase activity toward ONPG), in particular, the selectivity of the bonds formed. It is well reported that the


**Table 1.** Composition of the Reaction Mixture (Weight Percentage Referred to the Total Amount of Carbohydrates) Using 400 g/L Lactose and 1.5 U/mL  $\beta$ -Galactosidase from *Bacillus circulans* (Biolactase)<sup>a</sup>

reaction time (h)	Gal (%)	Glc (%)	Lact (%)	Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc (%)	Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc (%)	other GOS (%)	total GOS (%)
0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0
1.0	0.2	2.9	92.1	4.5	0.1	0.2	4.8
2.0	0.4	5.7	85.6	7.6	0.4	0.3	8.3
3.0	0.4	6.2	79.1	13.2	0.7	0.4	14.3
6.0	0.6	8.7	67.4	20.2	2.1	1.0	23.3
9.0	0.8	8.5	65.6	19.7	3.2	2.2	25.1
12.0	1.3	16.1	58.9	17.1	3.6	3.0	23.7
22.0	2.4	22.4	45.9	17.9	4.7	6.7	29.3
32.5	2.8	24.5	38.4	20.0	4.9	9.4	34.3
47.0	3.5	26.8	34.3	18.1	4.8	12.5	35.4
56.5	3.0	26.2	32.3	20.3	4.7	13.5	38.5
71.0	3.3	26.4	30.3	20.4	4.5	15.2	40.1
77.5	3.8	25.6	29.3	22.1	2.7	16.5	41.3
94.5	4.2	27.4	29.3	18.4	2.8	17.9	39.1
103.0	4.3	28.0	28.7	17.5	2.5	19.0	39.0
277.0	4.9	29.8	26.7	14.2	2.0	22.4	38.6
319.0	5.7	30.0	23.0	14.8	3.3	23.2	41.3
418.0	7.9	29.6	22.6	12.4	3.0	24.5	39.9


<sup>a</sup>Experimental conditions: 0.1 M sodium acetate buffer (pH 5.5), 40 °C.

chemical structure of the obtained oligosaccharides (composition, number of hexose units, and types of linkages between them) may affect their fermentation pattern by probiotic bacteria in the gut.<sup>7,12</sup>

Figure 2 shows the HPAEC-PAD chromatogram of the reaction mixture at the point of maximum GOS concentration. Peaks 1, 2, and 4 correspond to galactose, glucose, and lactose, respectively. As illustrated in the chromatogram, the two main products present in the reaction mixture were peaks 10 and 16. Using a commercial standard, peak 10 was identified as the trisaccharide 4-galactosyl-lactose [Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc]. Peak 16 was purified by semipreparative HILIC, as the sugar concentration in samples for HPAEC-PAD analysis was too low for an efficient scaling-up. The mass spectrum of peak 16 showed that it was a tetrasaccharide (data not shown). Analysis of the NMR spectra permitted assessment of the presence of five anomeric signals. Further inspection of the spectra allowed us to assign three of them as coming from the Gal residues, whereas the other two arose from the Glc moiety ( $\alpha$  and  $\beta$  anomers). From the combination of the information derived from COSY, TOCSY, NOESY, and HSQC (Figure 3A), most of the relevant <sup>1</sup>H and <sup>13</sup>C resonance signals belonging to the different residues could be assigned. The key information on the substitution pattern was extracted from the analysis of the HSQC experiment. The three cross peaks for the three Gal H4/C4 atom pairs were first identified. Their distinct chemical shifts permitted two of them to be distinguished as belonging to glycosylated Gal O-4 atoms, whereas the third one was nonsubstituted at O-4. From the cross-peak pattern in the TOCSY experiment it was possible to identify the corresponding intrasidual Gal H-1 anomeric signals for each Gal H-4. Then, the sequential connectivity of the sugar chain moieties




**Figure 4.** Kinetics of GOS formation at 1.5 U/mL using 400 g/L lactose catalyzed by  $\beta$ -galactosidase from *Bacillus circulans* (Bio-lactase): (A) formation of total GOS; (B) GOS distribution as a function of polymerization degree. Reaction conditions: 0.1 M sodium acetate buffer (pH 5.5), 40 °C. Dashed line: addition of a fresh batch of enzyme after 440 h.



**Figure 5.** Thermoinactivation of *Bacillus circulans*  $\beta$ -galactosidase at different temperatures in 0.1 M sodium acetate buffer (pH 5.5). Residual activity was determined at the indicated times using the ONPG assay.

was derived in a straightforward manner from the inter-residual cross peaks in the NOESY experiment. On this basis, the NMR



**Figure 6.** Kinetics of GOS formation at 15 U/mL using 400 g/L lactose catalyzed by  $\beta$ -galactosidase from *Bacillus circulans* (Bio-lactase): (A) formation of total GOS; (B) GOS distribution as a function of polymerization degree. Reaction conditions: 0.1 M sodium acetate buffer (pH 5.5), 40 °C.

data for peak 16 are consistent with a molecule that shows two galactosyl moieties  $\beta$ -(1 $\rightarrow$ 4)-linked to O-4 at the galactose unit of lactose, resulting in the tetrasaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc.

Commercially available standards and other GOS purified in our previous work with the  $\beta$ -galactosidase from *Kluyveromyces fragilis*<sup>11</sup> allowed us to identify in the chromatograms the disaccharides allolactose [Gal- $\beta$ (1 $\rightarrow$ 6)-Glc] (peak 3), 4-galactobiose [Gal- $\beta$ (1 $\rightarrow$ 4)-Gal] (peak 5), and Gal- $\beta$ (1 $\rightarrow$ 3)-Glc (peak 7), as well as the trisaccharide 6-galactosyl-lactose [Gal- $\beta$ (1 $\rightarrow$ 6)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc] (peak 6). We also purified by semipreparative HILIC peak 14, the mass spectrum indicating that it was a trisaccharide. In this case, the NMR spectra displayed four anomeric signals: two of them from the Gal residues and the two other from the Glc moiety ( $\alpha$  and  $\beta$ ). Following the same methodology described above, the combined analysis of the HSQC (Figure 3B) and TOCSY spectra permitted the existence of one terminal nonreducing Gal residue and one O-4 substituted Gal moiety, along with a terminal reducing Glc unit, substituted at O-3, to be distinguished. Thus, the NOESY spectrum permitted assessment of the sequential connectivity, indicating that peak 14 indeed corresponded to the trisaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 3)-Glc. Peaks 8, 9, 11–13, and 15 remained unknown.

**Table 2. Composition of the Reaction Mixture (Weight Percentage Referred to the Total Amount of Carbohydrates) Using 400 g/L Lactose and 15 U/mL  $\beta$ -Galactosidase from *Bacillus circulans* (Biolactase)<sup>a</sup>**


reaction time (h)	Gal (%)	Glc (%)	Lact (%)	Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc (%)	Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc (%)	other GOS (%)	total GOS (%)
0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0
0.5	0.9	12.5	56.7	25.5	3.2	1.2	29.9
1.0	2.0	16.2	47.6	26.4	5.1	2.7	34.2
2.0	4.0	19.1	39.9	23.1	5.6	8.3	37.0
6.5	5.9	18.0	26.7	13.8	3.3	32.2	49.4
10.5	9.4	31.5	17.2	7.4	1.7	32.9	41.9
26.0	17.0	37.1	6.9	4.4	0.2	34.5	39.1
30.5	16.2	33.2	6.8	4.6	0.6	38.6	43.9
46.0	15.1	35.4	5.0	4.7	0.6	39.1	44.5
54.5	15.9	37.7	4.5	4.1	0.0	37.8	42.0
70.5	18.2	40.3	3.3	3.3	0.0	34.9	38.2
78.0	18.4	39.3	4.6	2.0	0.0	35.7	37.7
143.0	18.3	41.3	2.4	1.3	0.0	36.6	37.9
151.0	21.7	39.4	4.8	1.8	0.0	32.4	34.1
175.0	23.9	42.6	2.4	1.3	0.0	29.8	31.1

<sup>a</sup>Experimental conditions: 0.1 M sodium acetate buffer (pH 5.5), 40 °C.

It is worth emphasizing that the two major products synthesized by *Bacillus circulans*  $\beta$ -galactosidase [the trisaccharide 4-galactosyl-lactose and the tetrasaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc] contained only  $\beta$ -(1 $\rightarrow$ 4) bonds. This result goes further in the control of regioselectivity by an appropriate selection of the enzyme. Yanahira et al. were the first to perform structural analysis of the GOS formed by *Bacillus circulans*  $\beta$ -galactosidase (Biolacta from Daiwa Kasei); they reported that the main product was 4-galactosyl-lactose, but the formation of tetrasaccharides was not mentioned.<sup>27</sup> Several disaccharides and trisaccharides were purified and characterized; the authors reported the presence of various GOS with  $\beta$ -(1 $\rightarrow$ 2) bonds,<sup>27</sup> which may correspond to some of the unknown peaks in our study. Recently, Song et al. analyzed the GOS production by the different isoforms of *Bacillus circulans*  $\beta$ -galactosidase.<sup>28</sup> Although the authors found significant differences in total GOS yield, the structural analysis of the synthesized compounds was not reported.


The product specificity of *Bacillus circulans*  $\beta$ -galactosidase contrasts with that of the *K. lactis* counterpart. The latter exhibits a tendency to synthesize  $\beta$ -(1 $\rightarrow$ 6) bonds: the main products in the reaction mixture are the disaccharides 6-galactobiose [Gal- $\beta$ (1 $\rightarrow$ 6)-Gal] and allolactose [Gal- $\beta$ (1 $\rightarrow$ 6)-Glc], as well as the trisaccharide 6-galactosyl-lactose [Gal- $\beta$ (1 $\rightarrow$ 6)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc].<sup>11,12</sup> Another difference between both enzymes is in the formation of disaccharides, because *Bacillus circulans*  $\beta$ -galactosidase yields a moderate amount of allolactose (peak 3), 4-galactobiose (peak 5), and 3-galactosyl-glucose (peak 7), whereas the *K. lactis* enzyme is able to use efficiently free galactose and glucose as acceptors yielding 6-galactobiose and allolactose, respectively, in notable yields.<sup>11</sup>

**Kinetics of GOS Synthesis.** Table 1 summarizes the evolution of carbohydrate composition of the reaction mixture employing 400 g/L lactose and 1.5 U/mL Biolactase as



**Figure 7.** Kinetics of GOS formation by Biolactase using skim milk: (A) total GOS production; (B) progress of glucose and galactose concentrations. Experimental conditions: 1.5 U/mL, 40 °C.

biocatalyst. The contribution of the main components [glucose, galactose, lactose, and the two major transglycosylation products with  $\beta$ (1 $\rightarrow$ 4) bonds], as well the rest of synthesized



**Figure 8.** HPAEC-PAD analysis of the reaction of skim milk with *Bacillus circulans*  $\beta$ -galactosidase (Biolactase) at the point of maximum GOS concentration (2 h). Peaks: 1, galactose; 2, glucose; 4, lactose; 5, 4-galactobiose; 7, 3-galactosyl-glucose; 10, 4-galactosyl-lactose; 16, Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc.

GOS considered as a pool, is reported. It has been broadly demonstrated that the maximum GOS yield is basically determined by the intrinsic enzyme properties (transgalactosylation to hydrolysis ratio) as well as substrate concentration.<sup>29</sup> In our work, the maximum GOS yield (41.3%, corresponding to a concentration of 165 g/L) was observed at 77.5 h, when lactose conversion accounted for 70%. This value is in the upper range of GOS yields reported (40–45%)<sup>8,30,31</sup> and is similar to that described for the isoform  $\beta$ -galactosidase-2 from *Bacillus circulans*.<sup>21</sup> However, GOS yields are generally lower than those reported for other related prebiotics such as fructooligosaccharides (approximately 65%) using similar strategies.<sup>32,33</sup>

The GOS concentration changes substantially with reaction time because GOS are simultaneously synthesized and hydrolyzed by  $\beta$ -galactosidases.<sup>3,8</sup> As a consequence, the time at which reaction is harvested has a crucial influence on GOS yield.<sup>6</sup> Although the concentration of the two main transgalactosylation products [containing only  $\beta(1\rightarrow4)$  bonds] reached a maximum followed by a progressive decrease (Table 1), the GOS production remained nearly constant after 70 h (Figure 4A) because the synthesis of other GOS increased with time. In particular, two disaccharides with  $\beta(1\rightarrow6)$  and  $\beta(1\rightarrow3)$  bonds [allolactose and Gal- $\beta(1\rightarrow3)$ -Glc, respectively] contributed significantly to the GOS concentration at the latter stages of the reaction. This effect could be explained by the hydrolytic activity of the enzyme, as  $\beta$ -galactosidases are specific for the cleavage of  $\beta(1\rightarrow4)$  bonds compared with  $\beta(1\rightarrow3)$  or  $\beta(1\rightarrow6)$  bonds present in other GOS.

Figure 4A shows the progress of total GOS formation and Figure 4B their distribution based on their polymerization degree. It is worth noting that the remaining lactose at 400 h was still high (23%) compared with similar processes using other glycosidic enzymes.<sup>9,11,13</sup> This effect could be caused by an inactivation of the  $\beta$ -galactosidase during the process. The experiment depicted in Figure 4 was carried out at 1.5 U/mL, which is a lower enzyme concentration than that typically used with  $\beta$ -galactosidases (3–12 U/mL).<sup>12,13,34</sup> In fact, at higher enzyme concentrations, reactions are faster and it is less probable that any effect of enzyme inactivation on reaction progress will be detected. To analyze the possible stability effect, we added fresh enzyme after 440 h (Figure 4A) and confirmed that the lactose diminished up to 31 g/L (7.8%) accompanied by a smooth increase of the GOS formed, especially in the contribution of disaccharides (Figure 4B).

#### Effect of Enzyme Concentration on GOS Formation.

Figure 5 illustrates the thermostability of *Bacillus circulans*  $\beta$ -galactosidase. As shown, the enzyme inactivated very rapidly between 50 and 60 °C, and most of its activity was lost in <2 h. At 40 °C the  $\beta$ -galactosidase lost nearly half of its activity in 10 h, and from that point the activity decay was slower. Taking into account that enzyme stability is enhanced in the presence of sugars,<sup>35</sup> the operational conditions (high concentration of lactose) are favorable for the stability of  $\beta$ -galactosidases.<sup>36</sup> However, the high concentration of residual lactose after 400 h using 1.5 U/mL seems to be related with the inactivation of the enzyme at long reaction times.

To minimize the inactivation effect, we performed the GOS synthesis at 10-fold higher enzyme concentration (15 U/mL). It has been widely reported that, working under kinetic control conditions, enzyme concentration has no effect on the maximum GOS yield as long as no enzyme inactivation takes place,<sup>37</sup> and it exerts a marked influence on only the reaction

time at which the maximum oligosaccharide concentration is achieved.<sup>13,37</sup>

Figure 6A shows that the maximum GOS production at 15 U/mL was achieved in 6.5 h, with a yield of 198 g/L. This value corresponds to 49.4% (w/w) of total sugars (Table 2), which is higher than the value obtained at 1.5 U/mL (165 g/L, 41.3%). Interestingly, the remaining lactose at the end of the reaction (10 g/L, 2.5% of total carbohydrates) is significantly lower than that obtained at 1.5 U/mL. This confirms that the stability of *Bacillus circulans*  $\beta$ -galactosidase is only moderate under typical GOS formation conditions; at 1.5 U/mL, the reaction is stopped before the final composition is reached.

Table 2 summarizes the evolution of the sugar composition of the mixture with time. Again, the two major products were those with only  $\beta(1\rightarrow4)$  bonds, but once formed they were quickly hydrolyzed by the  $\beta$ -galactosidase, enriching the mixture with GOS that contained other bonds. Figure 6B illustrates the distribution of di-, tri-, and tetrasaccharides with time. After 30 h, the contribution of disaccharides was very significant and even surpassed that of trisaccharides. Allolactose, 3-, 4- and 6-galactobioses, and Gal- $\beta(1\rightarrow3)$ -Glc were identified in notable amounts in the HPAEC-PAD chromatograms.

**GOS Production in Skim Milk.** Most reports on the treatment of milk with  $\beta$ -galactosidases focus on the hydrolytic activity to obtain lactose-free products.<sup>38,39</sup> The production of GOS in milk has been scarcely investigated, probably because the lactose concentration in milk (around 5% w/v) is not appropriate for an optimal transglycosylation to hydrolysis ratio.<sup>40</sup> To overcome this limitation, Chen et al. developed a multistep process applying ultrafiltration to separate lactose from milk proteins, followed by concentration of the permeate and further biotransformation with  $\beta$ -galactosidases.<sup>41</sup>

A positive aspect for the direct biotransformation of lactose in milk into GOS is that the pH of milk (approximately 6.7) is not far from the optimum pH of most  $\beta$ -galactosidases. In particular, the  $\beta$ -galactosidase from *Bacillus circulans* is a good choice for dairy products treatment (whole or skim milk, whey, etc.) as it presents a notable activity at pH 6.7 and is not inhibited by calcium cations.<sup>18,42</sup>

Gosling et al. assayed the *Bacillus circulans*  $\beta$ -galactosidase preparation Biolacta for GOS production in milk in the temperature range of 4–60 °C.<sup>43</sup> They observed that GOS yield increased with temperature, as described in other transglycosylation studies,<sup>44</sup> but neither kinetic nor structural analysis of the synthesized GOS was reported. Greenberg et al. reported that GOS accounted for 25% of total sugars in milk (quantified by paper chromatography) using the  $\beta$ -galactosidase from *Streptococcus thermophilus*.<sup>45</sup>

We performed a detailed analysis of GOS formation in skim milk using Biolactase preparation. The initial lactose concentration was 46 g/L, measured by HPAEC-PAD. Figure 7A illustrates the kinetics of GOS synthesis at 40 °C, where the typical pattern with a maximum GOS concentration followed by a progressive disappearance of GOS was observed due to the competition that is established between hydrolysis and transglycosylation (kinetic control).<sup>42,46</sup> This is clearly represented in Figure 7B, which shows how the glucose and galactose concentrations tend to converge at the end of the process. Maximum GOS yield (7.1 g/L, 15.4% of the total carbohydrates present in milk) was obtained at 2 h, when the lactose conversion accounted for 52%. Mozaffar et al. reported a maximum amount of GOS close to 5.5% of total sugars, which

was obtained at 39% conversion of lactose, using a purified  $\beta$ -galactosidase from *B. circulans*.<sup>42</sup>

Figure 8 shows the HPAEC-PAD chromatogram close to the point of maximum GOS concentration. As shown, the main peaks were the same as those described using 400 g/L lactose. The main GOS formed was the trisaccharide Gal- $\beta$ (1 $\rightarrow$ 4)-Gal- $\beta$ (1 $\rightarrow$ 4)-Glc (peak 10). The main difference between both experiments (skim milk vs 400 g/L lactose) was the absence of several minor peaks when milk was the lactose source.

## ■ ASSOCIATED CONTENT

### ● Supporting Information

NMR spectra used to identify the bonds between the galactosyl moieties. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*Postal address: Instituto de Catálisis y Petroleoquímica, CSIC, Cantoblanco, Marie Curie 2, 28049 Madrid, Spain. Fax: +34-91-5854760. E-mail: [fprou@icp.csic.es](mailto:fprou@icp.csic.es).

### Funding

Project BIO2010-20508-C04-01 from the Spanish Ministry of Science and Innovation supported this research. B.R.-C. was supported by a fellowship from the Spanish Ministry of Science and Innovation (FPI program).

### Notes

The authors declare no competing financial interest.

## ■ REFERENCES

- (1) Adam, A. C.; Rubio-Teixeira, M.; Polaina, J. Lactose: the milk sugar from a biotechnological perspective. *Crit. Rev. Food Sci.* **2004**, *44*, 553–557.
- (2) Torres, D. P.; Goncalves, M.; Teixeira, J. A.; Rodrigues, L. R. Galacto-oligosaccharides: production, properties, applications, and significance as prebiotics. *Compr. Rev. Food Sci. F.* **2010**, *9*, 438–454.
- (3) Park, A. R.; Oh, D. K. Galacto-oligosaccharide production using microbial  $\beta$ -galactosidase: current state and perspectives. *Appl. Microbiol. Biotechnol.* **2010**, *85*, 1279–1286.
- (4) Shadid, R.; Haarman, M.; Knol, J.; Theis, W.; Beermann, C.; Rjosk-Dendorfer, D.; Schendel, D. J.; Koletzko, B. V.; Krauss-Etschmann, S. Effects of galactooligosaccharide and long-chain fructooligosaccharide supplementation during pregnancy on maternal and neonatal microbiota and immunity – a randomized, double-blind, placebo-controlled study. *Am. J. Clin. Nutr.* **2007**, *86*, 1426–1437.
- (5) Rastall, R. A.; Gibson, G. R.; Gill, H. S.; Guarner, F.; Klaenhammer, T. R.; Pot, B.; Reid, G.; Rowland, I. R.; Sanders, M. E. Modulation of the microbial ecology of the human colon by probiotics, prebiotics and synbiotics to enhance human health: an overview of enabling science and potential applications. *FEMS Microbiol. Ecol.* **2005**, *52*, 145–152.
- (6) Gosling, A.; Stevens, G. W.; Barber, A. R.; Kentish, S. E.; Gras, S. L. Recent advances refining galactooligosaccharide production from lactose. *Food Chem.* **2010**, *121*, 307–318.
- (7) Cardelle-Cobas, A.; Corzo, N.; Olano, A.; Pelaez, C.; Requena, T.; Avila, M. Galactooligosaccharides derived from lactose and lactulose: influence of structure on *Lactobacillus*, *Streptococcus* and *Bifidobacterium* growth. *Int. J. Food Microbiol.* **2011**, *149*, 81–87.
- (8) Splechtina, B.; Nguyen, T. H.; Steinbock, M.; Kulbe, K. D.; Lorenz, W.; Haltrich, D. Production of prebiotic galacto-oligosaccharides from lactose using  $\beta$ -galactosidases from *Lactobacillus reuteri*. *J. Agric. Food Chem.* **2006**, *54*, 4999–5006.
- (9) Maischberger, T.; Leitner, E.; Nitisinprasert, S.; Juajun, O.; Yamabhai, M.; Nguyen, T. H.; Haltrich, D.  $\beta$ -Galactosidase from *Lactobacillus pentosus*: purification, characterization and formation of galacto-oligosaccharides. *Biotechnol. J.* **2010**, *5*, 838–847.
- (10) Iqbal, S.; Nguyen, T. H.; Nguyen, T. T.; Maischberger, T.; Haltrich, D.  $\beta$ -Galactosidase from *Lactobacillus plantarum* WCFS1: biochemical characterization and formation of prebiotic galacto-oligosaccharides. *Carbohydr. Res.* **2010**, *345*, 1408–1416.
- (11) Rodriguez-Colinas, B.; De Abreu, M. A.; Fernandez-Arrojo, L.; De Beer, R.; Poveda, A.; Jimenez-Barbero, J.; Haltrich, D.; Ballesteros, A. O.; Fernandez-Lobato, M.; Plou, F. J. Production of galacto-oligosaccharides by the  $\beta$ -galactosidase from *Kluyveromyces lactis*: comparative analysis of permeabilized cells versus soluble enzyme. *J. Agric. Food Chem.* **2011**, *59*, 10477–10484.
- (12) Martinez-Villaluenga, C.; Cardelle-Cobas, A.; Corzo, N.; Olano, A.; Villamiel, M. Optimization of conditions for galactooligosaccharide synthesis during lactose hydrolysis by  $\beta$ -galactosidase from *Kluyveromyces lactis* (Lactozym 3000 L HP G). *Food Chem.* **2008**, *107*, 258–264.
- (13) Chockchaisawasdee, S.; Athanasopoulos, V. I.; Niranjana, K.; Rastall, R. A. Synthesis of galacto-oligosaccharide from lactose using  $\beta$ -galactosidase from *Kluyveromyces lactis*: studies on batch and continuous UF membrane-fitted bioreactors. *Biotechnol. Bioeng.* **2005**, *89*, 434–443.
- (14) Maugard, T.; Gaunt, D.; Legoy, M. D.; Besson, T. Microwave-assisted synthesis of galacto-oligosaccharides from lactose with immobilized  $\beta$ -galactosidase from *Kluyveromyces lactis*. *Biotechnol. Lett.* **2003**, *25*, 623–629.
- (15) Guerrero, C.; Vera, C.; Plou, F.; Illanes, A. Influence of reaction conditions on the selectivity of the synthesis of lactulose with microbial  $\beta$ -galactosidases. *J. Mol. Catal. B: Enzym.* **2011**, *72*, 206–212.
- (16) Hsu, C. A.; Lee, S. L.; Chou, C. C. Enzymatic production of galactooligosaccharides by  $\beta$ -galactosidase from *Bifidobacterium longum* BCRC 15708. *J. Agric. Food Chem.* **2007**, *55*, 2225–2230.
- (17) Gosling, A.; Stevens, G. W.; Barber, A. R.; Kentish, S. E.; Gras, S. L. Effect of the substrate concentration and water activity on the yield and rate of the transfer reaction of  $\beta$ -galactosidase from *Bacillus circulans*. *J. Agric. Food Chem.* **2011**, *59*, 3366–3372.
- (18) Mozaffar, Z.; Nakanishi, K.; Matsuno, R. Purification and properties of  $\beta$ -galactosidases from *Bacillus circulans*. *Agric. Biol. Chem. Tokyo* **1984**, *48*, 3053–3061.
- (19) Mozaffar, Z.; Nakanishi, K.; Matsuno, R. Continuous production of galacto-oligosaccharides from lactose using immobilized  $\beta$ -galactosidase from *Bacillus circulans*. *Appl. Microbiol. Biot.* **1986**, *25*, 224–228.
- (20) Fujimoto, H.; Miyasato, M.; Ito, Y.; Sasaki, T.; Ajisaka, K. Purification and properties of recombinant  $\beta$ -galactosidase from *Bacillus circulans*. *Glycoconjugate J.* **1998**, *15*, 155–160.
- (21) Song, J.; Abe, K.; Imanaka, H.; Imamura, K.; Minoda, M.; Yamaguchi, S.; Nakanishi, K. Causes of the production of multiple forms of  $\beta$ -galactosidase by *Bacillus circulans*. *Biosci., Biotechnol., Biochem.* **2011**, *75*, 268–278.
- (22) Wei, L.; Xiaoli, X.; Shufen, T.; Bing, H.; Lin, T.; Yi, S.; Hong, Y.; Xiaoxiong, Z. Effective enzymatic synthesis of lactosucrose and its analogues by  $\beta$ -galactosidase from *Bacillus circulans*. *J. Agric. Food Chem.* **2009**, *57*, 3927–3933.
- (23) Bridiau, N.; Issaoui, N.; Maugard, T. The effects of organic solvents on the efficiency and regioselectivity of *N*-acetyl-lactosamine synthesis, using the  $\beta$ -galactosidase from *Bacillus circulans* in hydro-organic media. *Biotechnol. Prog.* **2010**, *26*, 1278–1289.
- (24) Farkas, E.; Schmidt, U.; Thiem, J.; Kowalczyk, J.; Kunz, M.; Vogel, M. Regioselective synthesis of galactosylated tri- and tetrasaccharides by use of  $\beta$ -galactosidase from *Bacillus circulans*. *Synthesis* **2003**, *5*, 699–706.
- (25) Usui, T.; Kubota, S.; Ohi, H. A convenient synthesis of  $\beta$ -D-galactosyl disaccharide derivatives using the  $\beta$ -D-galactosidase from *Bacillus circulans*. *Carbohydr. Res.* **1993**, *244*, 315–323.
- (26) Torres, P.; Batista-Viera, F. Immobilization of  $\beta$ -galactosidase from *Bacillus circulans* onto epoxy-activated acrylic supports. *J. Mol. Catal. B: Enzym.* **2012**, *74*, 230–235.

- (27) Yanahira, S.; Kobayashi, T.; Suguri, T.; Nakakoshi, M.; Miura, S.; Ishikawa, H.; Nakajima, I. Formation of oligosaccharides from lactose by *Bacillus circulans*  $\beta$ -galactosidase. *Biosci., Biotechnol., Biochem.* **1995**, *59*, 1021–1026.
- (28) Song, J.; Imanaka, H.; Imamura, K.; Minoda, M.; Katase, T.; Hoshi, Y.; Yamaguchi, S.; Nakanishi, K. Cloning and expression of a  $\beta$ -galactosidase gene of *Bacillus circulans*. *Biosci., Biotechnol., Biochem.* **2011**, *75*, 1194–1197.
- (29) Plou, F. J.; Gomez de Segura, A.; Ballesteros, A. Application of glycosidases and transglycosidases for the synthesis of oligosaccharides. In *Industrial enzymes: Structure, Function and Application*; Polaina, J., MacCabe, A. P., Eds.; Springer: New York, 2007; pp 141–157.
- (30) Rabiou, B. A.; Jay, A. J.; Gibson, G. R.; Rastall, R. A. Synthesis and fermentation properties of novel galacto-oligosaccharides by  $\beta$ -galactosidases from *Bifidobacterium* species. *Appl. Environ. Microbiol.* **2001**, *67*, 2526–2530.
- (31) Hansson, T.; Adlercreutz, P. Optimization of galactooligosaccharide production from lactose using  $\beta$ -glycosidases from hyperthermophiles. *Food Biotechnol.* **2001**, *15*, 79–97.
- (32) Ghazi, I.; Fernandez-Arrojo, L.; Gomez de Segura, A. G.; Alcalde, M.; Plou, F. J.; Ballesteros, A. Beet sugar syrup and molasses as low-cost feedstock for the enzymatic production of fructooligosaccharides. *J. Agric. Food Chem.* **2006**, *54*, 2964–2968.
- (33) Alvaro-Benito, M.; De Abreu, M.; Fernandez-Arrojo, L.; Plou, F. J.; Jimenez-Barbero, J.; Ballesteros, A.; Polaina, J.; Fernandez-Lobato, M. Characterization of a  $\beta$ -fructofuranosidase from *Schwanniomyces occidentalis* with transfructosylating activity yielding the prebiotic 6-kestose. *J. Biotechnol.* **2007**, *132*, 75–81.
- (34) Lee, Y. J.; Kim, C. S.; Oh, D. K. Lactulose production by  $\beta$ -galactosidase in permeabilized cells of *Kluyveromyces lactis*. *Appl. Microbiol. Biotechnol.* **2004**, *64*, 787–793.
- (35) Prasad, S.; Roy, I. Effect of disaccharides on the stabilization of bovine trypsin against detergent and autolysis. *Biotechnol. Prog.* **2010**, *26* (3), 627–635.
- (36) Ghazi, I.; Fernandez-Arrojo, L.; Garcia-Arellano, H.; Ferrer, M.; Ballesteros, A.; Plou, F. J. Purification and kinetic characterization of a fructosyltransferase from *Aspergillus aculeatus*. *J. Biotechnol.* **2007**, *128*, 204–211.
- (37) Buchholz, K.; Kasche, V.; Bornscheuer, U. T. *Biocatalysts and Enzyme Technology*; Wiley-VCH Verlag: Weinheim, Germany, 2005.
- (38) Puri, M.; Gupta, S.; Pahuja, P.; Kaur, A.; Kanwar, J. R.; Kennedy, J. F. Cell disruption optimization and covalent immobilization of  $\beta$ -D-galactosidase from *Kluyveromyces marxianus* YW-1 for lactose hydrolysis in milk. *Appl. Biochem. Biotechnol.* **2010**, *160*, 98–108.
- (39) Mlichova, Z.; Rosenberg, M. Current trends of  $\beta$ -galactosidase application in food technology. *J. Food Nutr. Res.* **2006**, *45*, 47–54.
- (40) Illanes, A. Whey upgrading by enzyme biocatalysis. *Electron. J. Biotechnol.* **2011**, *14* (6), <http://dx.doi.org/10.2225/vol14-issue6-fulltext-11>.
- (41) Chen, C. S.; Hsu, C. K.; Chiang, B. H. Optimization of the enzymic process for manufacturing low-lactose milk containing oligosaccharides. *Process Biochem.* **2002**, *38*, 801–808.
- (42) Mozaffar, Z.; Nakanishi, K.; Matsuno, R. Formation of oligosaccharides during hydrolysis of lactose in milk using  $\beta$ -galactosidase from *Bacillus circulans*. *J. Food Sci.* **1985**, *50*, 1602–1606.
- (43) Gosling, A.; Alftren, J.; Stevens, G. W.; Barber, A. R.; Kentish, S. E.; Gras, S. L. Facile pretreatment of *Bacillus circulans*  $\beta$ -galactosidase increases the yield of galactosyl oligosaccharides in milk and lactose reaction systems. *J. Agric. Food Chem.* **2009**, *57*, 11570–11574.
- (44) Linde, D.; Rodriguez-Colinas, B.; Estevez, M.; Poveda, A.; Plou, F. J.; Fernandez-Lobato, M. F. Analysis of neofructooligosaccharides production mediated by the extracellular  $\beta$ -fructofuranosidase from *Xanthophyllomyces dendrorhous*. *Bioresour. Technol.* **2012**, *109*, 123–130.
- (45) Greenberg, N. A.; Mahoney, R. R. Formation of oligosaccharides by  $\beta$ -galactosidase from *Streptococcus thermophilus*. *Food Chem.* **1983**, *10*, 195–204.
- (46) Rodriguez-Alegria, M. E.; Enciso-Rodriguez, A.; Ortiz-Soto, M. E.; Cassani, J.; Olvera, C.; Lopez-Munguia, L. Fructooligosaccharide production by a truncated *Leuconostoc citreum* inulosucrase mutant. *Biocatal. Biotransfor.* **2010**, *28*, 51–59.