

## Erratum to: High-dose remifentanil increases blood pressure and heart rate mediated by sympatho-activation in conscious rats

Tetsuro Shirasaka · Takeshi Yano ·  
Takato Kunitake · Isao Tsuneyoshi

Published online: 13 December 2012  
© Japanese Society of Anesthesiologists 2012

**Erratum to: J Anesth**  
**DOI 10.1007/s00540-012-1515-2**

Unfortunately the original version of this article contained errors. In the third paragraph from the end of the article, the third sentence should read as follows: “Remifentanil at a rate of 2.0 µg/kg/min decreased  $PO_2$ ; however,  $PCO_2$  was preserved within normal range”.

---

The online version of the original article can be found under  
doi:[10.1007/s00540-012-1515-2](https://doi.org/10.1007/s00540-012-1515-2).

---

T. Shirasaka (✉) · T. Yano · I. Tsuneyoshi  
Department of Anesthesiology and Intensive Care,  
Faculty of Medicine, University of Miyazaki,  
5200 Kihara Kiyotake, Miyazaki 889-1692, Japan  
e-mail: shirasak@med.miyazaki-u.ac.jp

T. Kunitake  
Department of Integrative Physiology, Faculty of Medicine,  
University of Miyazaki, Miyazaki, Japan