

# Sevoflurane in combination with propofol, not thiopental, induces a more robust neuroapoptosis than sevoflurane alone in the neonatal mouse brain

Tsuyoshi Tagawa · Shigeki Sakuraba ·  
Kazushi Kimura · Akira Mizoguchi

Received: 27 March 2013 / Accepted: 15 March 2014 / Published online: 6 April 2014  
© Japanese Society of Anesthesiologists 2014

## Abstract

**Purpose** Sevoflurane is the most widely used volatile anesthetic of general anesthesia. In children and neonates, it is commonly used alone or in combination with thiopental or propofol. A few recent studies reported that sevoflurane induced neuronal death in the developing rodent brain. We measured the neurotoxicity of these anesthetics at clinical doses, alone and in combination, in the developing mouse brain.

**Methods** Seven-day-old C57BL/6 mice were randomly assigned to 6 treatment groups. Three groups were exposed to 3 % sevoflurane for 6 h after injection of saline, thiopental (5 mg/kg), or propofol (10 mg/kg), whereas three groups were exposed to room air for 6 h after injection of equal doses of saline, thiopental, or propofol. Apoptosis in the hippocampal CA1 region (CA1) and retrosplenial cortex (RC) was assessed using caspase-3 immunostaining.

**Results** Sevoflurane alone caused significantly higher apoptosis in the CA1 compared with saline plus air ( $P = 0.04$ ). Sevoflurane in combination with propofol resulted in significantly greater numbers of apoptotic neurons than sevoflurane alone in both the CA1 and the RC ( $P = 0.04$ ). However, there was no significant difference in apoptotic neuron density in both the regions between the groups treated with sevoflurane alone and in combination with thiopental ( $P = 0.683$ ).

**Conclusion** Sevoflurane alone can induce neuronal apoptosis, and this effect is enhanced by propofol. Thiopental did not exacerbate the neurotoxicity of sevoflurane. There is the possibility that the combination of sevoflurane and propofol is a more harmful anesthetic technique than sevoflurane alone in pediatric patients.

**Keywords** Sevoflurane · Apoptosis · Children

## Introduction

Anesthetic exposure, even during critical stages of neurodevelopment, has been considered safe and without any adverse long-term consequences. However, many recent animal studies suggest that exposure to certain anesthetics during periods of rapid synaptogenesis can evoke neuronal apoptosis and inhibit neurogenesis, leading to lasting cognitive impairments [1].

Many clinical anesthetics potentiate inhibitory transmission mediated by gamma aminobutyric acid type A ( $GABA_A$ ) receptors and/or inhibit excitatory glutamateric transmission mediated by *N*-methyl-D-aspartate (NMDA) receptors [2]. Simultaneous administration of a  $GABA_A$  agonist and NMDA receptor antagonist in particular can evoke widespread neuronal apoptosis in the developing brain, suggesting that anesthetics targeting these ionotropic receptors may have deleterious effects on the infant brain [3, 4]. Exposure of infant rats to midazolam, nitrous oxide, and isoflurane for 6 h (each at clinically relevant doses) caused widespread neurodegeneration and cognitive deficits that persisted to adulthood [3]. The NMDA antagonist ketamine and a  $GABA_A$  agonist (diazepam, thiopental, or propofol), particularly in combination, induced neuronal apoptosis and long-term motor and cognitive dysfunction

T. Tagawa (✉) · K. Kimura · A. Mizoguchi  
Department of Neural Regeneration and Cell Communication,  
Mie University Graduate School of Medicine, 2-174 Edobashi,  
Tsu, Mie 514-8507, Japan  
e-mail: tsuyoshitagawa@gmail.com

S. Sakuraba  
Department of Anesthesiology, Clinical Care Medicine,  
Kanagawa Dental College, Yokosuka, Kanagawa, Japan

in young mice [4–6]. However, these particular anesthesia cocktails are no longer used for most pediatric anesthetic procedures.

Sevoflurane (2,2,2-trifluoro-1-[trifluoromethyl]ethyl fluoromethyl ether) is currently the most widely used inhalation anesthetic for pediatric surgery and cesarean delivery because of its properties of rapid inhalational induction, tolerable odor, quick emergence and relative cardiovascular stability for infants and children [7]. In a survey conducted by the Society for Pediatric Anesthesia, 95 % of anesthesiologists reported the use of sevoflurane [8]. Sevoflurane enhances GABA<sub>A</sub> receptor currents [9] and likely also blocks NMDA receptors [10]. For induction of routine pediatric anesthesia, sevoflurane is usually used alone or in combination with thiopental or propofol. A few recently published studies reported neurotoxic effects of sevoflurane alone [11–13] but no enhancement when used in combination with other substances.

In light of evidence demonstrating apoptosis in the neonatal brain following administration of sevoflurane, thiopental, and/or propofol, and the frequent use of these agents in current pediatric anesthesia practice, we examined the neurotoxic effects of these agents, alone and in combination, in neonatal mice during the brain development period.

## Materials and methods

### Animals

All experiments were conducted according to the institutional ethical guidelines for animal experiments of Mie University and were approved by the Committee for Animal Research at Mie University School of Medicine, Tsu, Mie, Japan. The C57BL/6 mice were purchased from SLC (SLC Japan Inc., Shizuoka, Japan). The animals were maintained on a 12-h light–dark cycle (lights on from 07:00 to 19:00) in a temperature controlled room ( $21 \pm 1$  °C). The mice had ad libitum access to water and food. Both male and female mice were used in the experimental and control groups. Control and experimental pups were drawn randomly from the same litters. Efforts were made to minimize the number of animals used and their suffering.

### Anesthesia treatment

On postnatal day 7 (P7), mice were randomly assigned to one of six treatment groups ( $n = 8$  per group): (1) a vehicle (Veh) group was administered a single subcutaneous injection of saline followed by 6 h exposure to air, (2) a Thio group given a single subcutaneous injection of

thiopental (5 mg/kg) followed by 6 h exposure to air, (3) a Prop group given a single subcutaneous injection of propofol (10 mg/kg) followed by 6 h exposure to air, (4) a Sevo group administered with a single subcutaneous injection of saline followed by 6 h exposure to 3 % sevoflurane, (5) a Thio + Sevo group was given a single subcutaneous injection of thiopental (5 mg/kg) followed by 6 h exposure to 3 % sevoflurane, and (6) a Prop + Sevo group was given a single subcutaneous injection of propofol (10 mg/kg) followed by 6 h exposure to 3 % sevoflurane. Thiopental and propofol were dissolved in saline and all mice were injected with an equal volume of solution per unit weight (20 ml/kg). For sevoflurane or air exposure, mice were placed in an acrylic box connected to an anesthetic machine. The total gas flow was 2 l/min. Air was used as the carrier for sevoflurane. During anesthetic exposure, the mice were maintained at 38 °C using a heating pad. The concentration of sevoflurane (3 %) was based on previously published work [13], and fell within the range used for pediatric anesthesia [7, 14]. At 3 % sevoflurane, the 6 h exposure had no detrimental effect on cerebral perfusion, arterial blood gas concentrations, or blood glucose concentration in neonatal C57BL/6 mice [13]. The doses of thiopental (5 mg/kg) and propofol (10 mg/kg) administered alone are low and do not increase neonatal neurodegeneration [4]. These low doses were chosen to avoid suppression of respiration when administered with 3 % sevoflurane, because the infant mice are too small to permit adequate airway management.

### SpO<sub>2</sub> analysis

To avoid any potential complications from invasive manipulations and to reduce the number of experimental animals, blood gases and end tidal CO<sub>2</sub> were not measured. However, percutaneous oxygen saturation (SpO<sub>2</sub>) was measured using an SpO<sub>2</sub> monitoring system (Mouse STAT; Kent Scientific Co., Torrington, CT, USA). A monitoring sensor was attached to the leg of each mouse.

### Laser Doppler

Cerebral blood flow (CBF) was measured by a laser-Doppler blood flowmeter (ALF21N; ADVANCE, Tokyo, Japan). Mice were taken out of the chamber before and every hour during anesthetic treatment and were placed face down on the floor while being continuously exposed to sevoflurane via a tube with its opening positioned at the nose of the animals. Veh group mice were exposed to air for a corresponding period to the anesthetized mice. Their head skins were peeled for scanning CBF, and data were captured using appropriate software (LabChart version 8; ADInstruments, Nagoya, Japan). The perfusion response is

presented in arbitrary perfusion units. Arbitrary perfusion unit values were compared among groups at baseline and at 1-h intervals for 6 h.

### Histopathological studies

Immediately following experimental treatments, mice were perfused transcardially with 0.1 M phosphate buffer (PB) containing 4 % paraformaldehyde. Brain sections were prepared and processed for activated caspase-3 immunostaining, a well-established procedure for measuring neonatal apoptosis in the developing brain [12, 15, 16]. In brief, brains were removed and post-fixed overnight in the same fixative at 4 °C. Serial coronal sections of 80- $\mu$ m thickness were cut on a vibratome, blocked for 1 h in blocking solution (PBS with 2 % BSA, 0.2 % skim milk powder, and 0.1 % triton X-100), followed by incubation in rabbit anti-activated caspase-3 polyclonal antibody (1:250; Cell Signaling, Danvers, MA, USA; 9661). Immunolabeled sections were washed and incubated in a goat anti-rabbit Ig conjugated to Alexa Fluor 488 (1:250, A11034; Invitrogen, Carlsbad, CA, USA) or goat anti-mouse antibodies conjugated to Alexa Fluor 594 (1:250, A11032; Invitrogen) for 4 h at room temperature.

Caspase-3-positive cells were counted in the hippocampal CA1 region (CA1) and retrosplenial cortex (RC). These two regions were chosen because they are particularly vulnerable to anesthesia-mediated neurodegeneration [3, 11, 13, 17] and are integral components of the extra-hippocampal circuit, which have been identified as important for mediating learning and memory functions [18]. Images were acquired using an Olympus FV1000 confocal microscope controlled by FV10-ASW 1.7 software (Olympus Corporation, Tokyo, Japan). The images were then printed on paper, and an investigator blind to the treatment history counted the number of caspase-3-positive cells in the CA1 and RC. These regions were defined by plates 48–52 in the Paxinos and Franklin mouse brain atlas [19]. The density of caspase-3-positive cells in a particular brain region was calculated by dividing the number of caspase-3-positive cells by the area of that brain region.

### Statistical analysis

Previous experiments in neonatal rodents and nonhuman primates demonstrated extensive neurotoxicity in response to anesthetics and other toxins, allowing for the use of relatively small numbers of experimental animals (3–8) per treatment group [3, 4, 11, 13, 15–17, 20, 21]. Statistical comparisons between the groups were performed by using a Kruskal–Wallis test. A two-way ANOVA for repeated measures was performed to examine differences in SpO<sub>2</sub> and CBF. All calculations were performed using the SPSS

program (IBM Co., Tokyo, Japan). Statistical significance was accepted at  $P < 0.05$ .

### Results

To test the potential neurotoxicity of sevoflurane (Sevo) alone and in combination with thiopental (Thio) or propofol (Prop), P7 mice were divided into 6 treatment groups and exposed to air or 3 % Sevo for 6 h following injection with saline (Veh and Sevo groups), 5 mg/kg Thio (Thio and Thio + Sevo groups), or 10 mg/kg Prop (Prop and Prop + Sevo groups). All 48 animals survived the anesthetic exposures. Mice subjected to Sevo, Thio + Sevo, and Prop + Sevo reached deep anesthesia as indicated by absent or only minor reactions to pain stimuli. Administration of Thio and Prop alone with air exposure resulted in transient sluggishness but no deep anesthesia. SpO<sub>2</sub> did not differ significantly among groups (Table 1; two-way repeated measures ANOVA, groups:  $F = 2.22$ ,  $P = 0.07$ , time:  $F = 0.72$ ,  $P = 0.64$ , interaction between groups and time:  $F = 0.41$ ,  $P = 0.99$ ), suggesting that all anesthesia treatments did not induce significant disturbances in oxygenation. Moreover, to assess adequacy of cerebral perfusion, we measured CBF during anesthesia using a laser-Doppler blood flowmeter. There were no significant differences in CBF at any point during the 6 h of anesthesia among groups (Table 2; two-way repeated measures ANOVA, groups:  $F = 0.51$ ,  $P = 0.77$ , time:  $F = 1.12$ ,  $P = 0.31$ , interaction between groups and time:  $F = 0.91$ ,  $P = 0.61$ ).

Caspase-3-positive cells were observed sporadically in both the CA1 and the RC of Veh group mice (Fig. 1). Neither Thio nor Prop injection alone induced a significant increase in the number of caspase-3-positive cells (Thio and Prop groups) compared with the Veh group (Fig. 2). The density of apoptotic neurons was significantly higher in the Sevo group compared with the Veh group in the CA1. In both the CA1 and the RC, Prop + Sevo induced a significant increase in the number of caspase-3-positive cells compared with Sevo ( $P = 0.04$ ). However, there were no significant differences in caspase-3-positive cell density between the Sevo and Thio + Sevo groups ( $P = 0.683$ ).

### Discussion

General anesthetics can induce widespread neuronal cell death and persistent behavioral sequelae, including motor and cognitive deficits, in both the neonatal rodent and primate brains [1]. These results have led to a re-evaluation of the safety of many anesthetics used for pediatric patients. Sevoflurane has a more favorable

**Table 1** Percutaneous oxygen saturation in mice

SpO <sub>2</sub> (%)							
Time (h)	0	1	2	3	4	5	6
Veh	97.4 ± 0.5	96.9 ± 0.5	96.9 ± 0.4	97.4 ± 0.5	97.1 ± 0.6	96.9 ± 0.4	97.1 ± 0.3
Thio	95.9 ± 0.4	95.6 ± 0.5	96.5 ± 0.9	97.3 ± 0.6	97.3 ± 0.6	97.0 ± 0.7	97.0 ± 0.6
Prop	96.6 ± 0.6	96.8 ± 0.5	96.3 ± 0.8	96.5 ± 0.5	95.8 ± 0.3	96.5 ± 0.4	97.0 ± 0.6
Sevo	97.1 ± 0.4	95.9 ± 0.9	96.0 ± 0.9	96.0 ± 0.4	95.8 ± 0.7	96.5 ± 1.2	96.8 ± 0.6
Thio + Sevo	96.4 ± 0.6	96.0 ± 0.8	96.3 ± 0.9	96.5 ± 0.5	95.5 ± 1.0	95.8 ± 1.0	96.9 ± 0.6
Prop + Sevo	96.8 ± 0.5	95.6 ± 1.0	96.1 ± 0.8	96.4 ± 0.7	96.4 ± 0.6	96.3 ± 0.6	95.8 ± 1.0

Values are presented as mean ± SEM

Analysis of percutaneous oxygen saturation (SpO<sub>2</sub>) revealed no significant differences at all time points among groups (two-way repeated measures ANOVA, groups:  $P > 0.05$ , time:  $P > 0.05$ , interaction between groups and time:  $P > 0.05$ ,  $n = 8$  per group)

**Table 2** Cerebral blood flow in mice

CBF (arbitrary perfusion unit)							
Time (h)	0	1	2	3	4	5	6
Veh	27.5 ± 2.1	26.6 ± 1.2	22.5 ± 1.7	24.4 ± 1.7	23.8 ± 1.6	25.9 ± 1.9	22.8 ± 1.2
Thio	22.1 ± 1.1	28.6 ± 2.6	26.9 ± 2.5	27.0 ± 2.4	26.5 ± 1.8	24.1 ± 2.3	24.5 ± 3.4
Prop	20.8 ± 3.1	24.6 ± 1.4	25.5 ± 2.3	26.0 ± 2.6	25.9 ± 2.0	22.9 ± 1.4	22.0 ± 1.1
Sevo	25.5 ± 2.4	24.8 ± 1.8	25.3 ± 3.5	26.8 ± 1.9	26.0 ± 2.9	28.9 ± 3.3	25.8 ± 3.3
Thio + Sevo	22.3 ± 2.2	26.0 ± 1.2	28.6 ± 3.1	25.6 ± 1.0	26.1 ± 2.2	26.3 ± 1.3	26.6 ± 2.1
Prop + Sevo	23.3 ± 1.2	23.1 ± 1.5	26.1 ± 1.7	26.5 ± 1.8	25.6 ± 1.0	25.9 ± 0.9	27.6 ± 1.6

Values are presented as mean ± SEM

Analysis of cerebral blood flow (CBF) revealed no significant differences at all time points among groups (two-way repeated measures ANOVA, groups:  $P > 0.05$ , time:  $P > 0.05$ , interaction between groups and time:  $P > 0.05$ ,  $n = 8$  per group)


neurotoxicity profile than the other commonly used inhalational anesthetics isoflurane and desflurane [11, 22] and therefore has become the preferred choice for pediatric anesthesia. To examine the potential neurotoxicity of sevoflurane, both alone and when combined with thiopental or propofol, we measured apoptotic cell numbers in the CA1 and RC following anesthetic treatment.

In the present study, sevoflurane in combination with propofol was more neurotoxic than sevoflurane alone. However, thiopental did not exacerbate sevoflurane-induced neurotoxicity. Propofol, like sevoflurane, acts mainly via GABA<sub>A</sub> receptor agonism, and to a lesser extent via NMDA receptor antagonism [23], while thiopental only potentiates GABA<sub>A</sub> receptor functioning. Neonatal co-administration of an NMDA antagonist and a GABA<sub>A</sub> agonist, even at doses that do not cause apoptosis alone, led to significant deleterious effects compared with when these were administered alone. Furthermore, the effects seem to be synergistic [3, 4].

One limitation of this study is that the blood gases and end tidal CO<sub>2</sub> of individual animals were not measured. To assess oxygenation levels, we examined SpO<sub>2</sub> in mice


every hour during anesthetic treatment. SpO<sub>2</sub> exceeded 90 % in all groups before and every hour during anesthetic treatment. In addition, hypercarbia does not significantly increase brain cell death in the hippocampus and cortex of neonatal rats, compared with control animals exposed to room air [24]. These facts led us to presume the unlikelihood of apoptosis in this study being caused by hypoxia/hypercarbia.

Consistent with previous studies [11–13], we demonstrated that even sevoflurane alone caused significant neuronal apoptosis in the CA1. However, no commonly used anesthetic has a perfect safety record in animals, and withholding anesthesia during surgery or other invasive procedures is unethical. Therefore, there appears to be no current alternative to sevoflurane. Furthermore, painful stimulation in unanesthetized newborn humans and animals can also result in neurological and behavioral abnormalities [25–30]. Most pediatric surgical procedures are not elective and cannot be postponed without incurring some degree of risk. If surgery is required, anesthesia with a small risk of limited neurotoxicity is superior to no anesthesia. Alleviation of pain and stress during treatment should remain the primary goal.



**Fig. 1** Expression pattern of the apoptosis marker activated caspase-3. Images were obtained from coronal brain sections from the hippocampal CA1 region (*left column*) and retrosplenial cortex (*right column*) under a laser confocal microscope immediately after Veh, Sevo, Thio + Sevo, or Prop + Sevo treatment. Cells positive for activated caspase-3 were stained *bright green* using a secondary antibody conjugated to Alexa Fluor 488. *Arrows* mark cells positive for activated caspase-3. The different treatments are symbolized as follows: *Veh* saline + air, *Sevo* saline + 3 % sevoflurane, *Thio + Sevo* thiopental (5 mg/kg) + 3 % sevoflurane and *Prop + Sevo* propofol (10 mg/kg) + 3 % sevoflurane (color figure online)

Based on our results, there is the possibility that the combination of sevoflurane and propofol is more prone to induce apoptosis in the brain than sevoflurane alone in pediatric patients, although there are the difficulties transferring the results with rodents to humans. This possibility



**Fig. 2** The *box plots* showing quantification of caspase-3-positive cells for each treatment condition in the hippocampal CA1 region and retrosplenial cortex. *Horizontal lines* represent respective group medians; *boxes*, 25<sup>th</sup>–75<sup>th</sup> percentile; and *whiskers*, minimum–maximum values. The treatment abbreviations are summarized in Fig. 1. *n* = 8 per group. \**P* = 0.04 versus Sevo; \*\**P* = 0.001 versus Veh or Prop; +*P* = 0.04 versus Veh; ++*P* = 0.006 versus Thio; #*P* = 0.001 versus Veh; ##*P* = 0.04 versus Thio

can occupy an important place in clinical practice because IV propofol followed by inhaled sevoflurane is a commonly used anesthetic technique for induction of pediatric anesthesia. If our current findings can be extrapolated to clinical practice, this study may provide clinicians with neurotoxicity data to make more informed decisions on anesthesia regimens for pediatric patients. Additional research is obviously needed to determine the potential neurotoxicity of various anesthetic drug combinations and possible neuroprotective strategies. In the interim, anesthesiologists should be aware of all current neurotoxicology results in order to select the least harmful anesthetic agents for newborn and infant anesthesia.



## References

- Stratmann G. Neurotoxicity of anesthetic drugs in the developing brain. *Anesth Analg*. 2011;113:1170–9.
- Ikonomidou C, Bittigau P, Koch C, Genz F, Hoerster F, Felderhoff-Mueser U, Tenkova T, Dikranian K, Olney JW. Neurotransmitters and apoptosis in the developing brain. *Biochem Pharmacol*. 2001;62:401–5.
- Jevtovic-Todorovic V, Hartman RE, Izumi Y, Benshoff ND, Dikranian K, Zorumski CF, Olney JW, Wozniak DF. Early exposure to common anesthetic agents causes widespread neurodegeneration in the developing rat brain and persistent learning deficits. *J Neurosci*. 2003;23:876–82.
- Fredriksson A, Pontén E, Gordh T, Eriksson P. Neonatal exposure to a combination of *N*-methyl-D-aspartate and gamma-aminobutyric acid type A receptor anesthetic agents potentiates apoptotic neurodegeneration and persistent behavioral deficits. *Anesthesiology*. 2007;107:427–36.
- Fredriksson A, Archer T, Alm H, Gordh T, Eriksson P. Neurofunctional deficits and potentiated apoptosis by neonatal NMDA antagonist administration. *Behav Brain Res*. 2004;153:367–76.
- Fredriksson A, Archer T. Neurobehavioural deficits associated with apoptotic neurodegeneration and vulnerability for ADHD. *Neurotox Res*. 2004;6:435–56.
- Lerman J, Sikich N, Kleinman S, Yentis S. The pharmacology of sevoflurane in infants and children. *Anesthesiology*. 1994;80:814–24.
- Murray JP, Bhananker SM. Recent findings from the pediatric perioperative cardiac arrest (POCA) registry. *ASA Newslett*. 2005;69:10–2.
- Nishikawa K, Harrison NL. The actions of sevoflurane and desflurane on the gamma-aminobutyric acid receptor type A: effects of TM2 mutations in the alpha and beta subunits. *Anesthesiology*. 2003;99:678–84.
- Hollmann MW, Liu HT, Hoememann CW, Liu WH, Durieux ME. Modulation of NMDA receptor function by ketamine and magnesium, part II: interactions with volatile anesthetics. *Anesth Analg*. 2001;92:1182–91.
- Kodama M, Satoh Y, Otsubo Y, Araki Y, Yonamine R, Masui K, Kazama T. Neonatal desflurane exposure induces more robust neuroapoptosis than do isoflurane and sevoflurane and impairs working memory. *Anesthesiology*. 2011;115:979–91.
- Istaphanous GK, Howard J, Nan X, Hughes EA, McCann JC, McAuliffe JJ, Danzer SC, Loepke AW. Comparison of the neuroapoptotic properties of equipotent anesthetic concentrations of desflurane, isoflurane, or sevoflurane in neonatal mice. *Anesthesiology*. 2011;114:578–87.
- Satomoto M, Satoh Y, Terui K, Miyao H, Takishima K, Ito M, Imaki J. Neonatal exposure to sevoflurane induces abnormal social behaviors and deficits in fear conditioning in mice. *Anesthesiology*. 2009;110:628–37.
- Katoh T, Ikeda K. Minimum alveolar concentration of sevoflurane in children. *Br J Anaesth*. 1992;68:139–41.
- Brambrink AM, Evers AS, Avidan MS, Farber NB, Smith DJ, Martin LD, Dissen GA, Creeley CE, Olney JW. Ketamine-induced neuroapoptosis in the fetal and neonatal rhesus macaque brain. *Anesthesiology*. 2012;116:372–84.
- Brambrink AM, Evers AS, Avidan MS, Farber NB, Smith DJ, Zhang X, Dissen GA, Creeley CE, Olney JW. Isoflurane-induced neuroapoptosis in the neonatal rhesus macaque brain. *Anesthesiology*. 2010;112:834–41.
- Li Y, Liang G, Wang S, Meng Q, Wang Q, Wei H. Effects of fetal exposure to isoflurane on postnatal memory and learning in rats. *Neuropharmacology*. 2007;53:942–50.
- Wozniak DF, Hartman RE, Boyle MP, Vogt SK, Brooks AR, Tenkova T, Young C, Olney JW, Muglia LJ. Apoptotic neurodegeneration induced by ethanol in neonatal mice is associated with profound learning/memory deficits in juveniles followed by progressive functional recovery in adults. *Neurobiol Dis*. 2004;17:403–14.
- Paxinos G, Franklin KBJ. The mouse brain in stereotaxic coordinates. 2nd ed. San Diego: Academic Press; 2001.
- Shu Y, Patel SM, Pac-Soo C, Fidalgo AR, Wan Y, Maze M, Ma D. Xenon pretreatment attenuates anesthetic-induced apoptosis in the developing brain in comparison with nitrous oxide and hypoxia. *Anesthesiology*. 2010;113:360–8.
- Sanders RD, Xu J, Shu Y, Januszewski A, Halder S, Fidalgo A, Sun P, Hossain M, Ma D, Maze M. Dexmedetomidine attenuates isoflurane-induced neurocognitive impairment in neonatal rats. *Anesthesiology*. 2009;110:1077–85.
- Liang G, Ward C, Peng J, Zhao Y, Huang B, Wei H. Isoflurane causes greater neurodegeneration than an equivalent exposure of sevoflurane in the developing brain of neonatal mice. *Anesthesiology*. 2010;112:1325–34.
- Rugolph U, Antkowiak B. Molecular and neuronal substrates for general anaesthetics. *Nat Rev Neurosci*. 2004;5:709–20.
- Stratmann G, May LD, Sall JW, Alvi RS, Bell JS, Ormerod BK, Rau V, Hilton JF, Dai R, Lee MT, Visrodia KH, Ku B, Zusmer EJ, Guggenheim J, Firouzian A. Effect of hypercarbia and isoflurane on brain cell death and neurocognitive dysfunction in 7-day-old rats. *Anesthesiology*. 2009;110:849–61.
- Sanders RD, Ma D, Brooks P, Maze M. Balancing paediatric anaesthesia: preclinical insights into analgesia, hypnosis, neuroprotection, and neurotoxicity. *Br J Anaesth*. 2008;101:597–609.
- Anand KJ, Sippell WG, Aynsley-Green A. Randomised trial of fentanyl anaesthesia in preterm babies undergoing surgery: effects on the stress response. *Lancet*. 1987;1:243–8.
- Fitzgerald M. The development of nociceptive circuits. *Nat Rev Neurosci*. 2005;6:507–20.
- Taddio A, Katz J, Ilersich AL, Koren G. Effect of neonatal circumcision on pain response during subsequent routine vaccination. *Lancet*. 1997;349:599–603.
- Hermann C, Hohmeister J, Demirakça S, Zohsel K, Flor H. Long-term alteration of pain sensitivity in school-aged children with early pain experiences. *Pain*. 2006;125:278–85.
- Page GG, Blakely WP, Kim M. The impact of early repeated pain experiences on stress responsiveness and emotionality at maturity in rats. *Brain Behav Immun*. 2005;19:78–87.