

REACTION OF BLEOMYCIN WITH DNA
STRAND SCISSION OF DNA IN THE ABSENCE OF
SULFHYDRYL OR PEROXIDE COMPOUNDS

ISAO SHIRAKAWA, MASAO AZEGAMI,
SIN-ICHI ISHII and HAMA O UMEZAWA*

Faculty of Pharmaceutical Sciences, Hokkaido University,
Sapporo, Japan

* Institute of Microbial Chemistry, Shinagawa-ku, Tokyo, Japan

(Received for publication July 24, 1971)

The effect of bleomycin on the infectivity and the molecular integrity of bacteriophage ϕ X174 DNA was examined by means of the spheroplast assay and sucrose density gradient centrifugation.

Bleomycins caused degradation of the DNA, both in the single and double stranded forms even in the absence of 2-mercaptoethanol or hydrogen peroxide, although the reaction was accelerated by the addition of either of the compounds. All the bleomycin compounds examined, A₂, A₅ and B₂, were effective for the reaction but copper-chelated A₂ was not. The reaction was inhibited by the addition of EDTA*.

Bleomycins are antibiotics which are produced by *Streptomyces verticillus* and have antitumour and antibacterial activities^{1,2,3}.

It has been reported that bleomycins inhibit DNA synthesis rather than protein or RNA synthesis in *Escherichia coli*, EHRLICH carcinoma and HeLa cells⁴. They cause strand scissions in DNA of growing *E. coli*, and HeLa cells⁵. They bind to DNA and decrease the melting temperature (T_m) and the molecular size in the presence of sulfhydryl compounds or hydrogen peroxide, *in vitro*^{6,7,8,9}. It has been suggested that sulfhydryl compounds or hydrogen peroxide may take part in the *in vivo* action of bleomycin^{5,8}.

In the study of effects of bleomycin on the growth of bacteriophage ϕ X174, which has a single-stranded circular DNA (phage DNA) and also produces double-stranded circular DNA called replicative form (RF)** in the host cell during the development, the degradation of the single-stranded phage DNA and the double-stranded RF DNA was observed *in vitro* in the absence of sulfhydryl compounds or hydrogen peroxide. The results are presented in this paper.

Materials and Methods

The following bleomycins were used: mixture (lot. F-7, A₂; 69.0%, B₂; 28.8%, A₂'; 1.5%, B₁; 1.5%, A₅; 0.9%, A₁; 0.5%). A₂ (lot. F1-8), A₅ (lot. 3, spd), A₂(Cu) (lot. F1-8, copper-chelated), B₂ (lot. 32). All bleomycins were kindly offered by Nippon Kayaku

* EDTA: ethylenediaminetetraacetic acid.

** RF: RF-I (both strands are closed) and RF-II (one or both strands are nicked).

Co., Tokyo. (Methyl-³H) thymidine (specific activity 15.5 c/m mole) was purchased from Schwarz Bioresearch, Inc. Bacterial and phage strains were given to us by Dr. R. L. SINSHEIMER.

Zone Sedimentation and Radioactivity Measurement

DNA or its reaction mixture (in 0.2~0.6 ml) was layered on a 5~20% sucrose density gradient in 0.05 M tris-0.5 M sodium chloride-1 mM EDTA (pH 7.6, Tris-saline-EDTA buffer) and centrifuged at 4°C in an SW 27 rotor of a Beckman L2-65B centrifuge at 25,000 r.p.m. for 16 hours. Fractions were collected from the bottom of tubes. The DNA in each fraction was precipitated with cold 5% (final concentration) trichloroacetic acid (TCA) after adding 100 µg of herring sperm DNA as a carrier. Precipitates containing ³²P or ³H-DNA were collected on glass fibre paper (Whatman GF/C), dried and counted in an Aloka scintillation counter (toluene base scintillator).

Bioassay

The biological activity of the phage DNA was measured by the spheroplast assay as described by GUTHRIE *et al.*¹⁰⁾

Preparation of Phage DNA

³²P-Labeled ϕ X174 *am3* (lysis defective mutant) was prepared essentially according to the procedure of SINSHEIMER *et al.*¹¹⁾ ³²P-Labeled ϕ X174 *am3* DNA was prepared by the method of GUTHRIE *et al.*¹⁰⁾ The phage DNA was purified by sucrose density gradient centrifugation, if necessary. Fractions containing ³²P-phage DNA were pooled and the DNA was precipitated with 2.5 volumes of ethanol after the addition of tRNA (10 µg/ml) or the cold phage DNA (80 µg/ml) as a carrier. Precipitates were dissolved in 0.05 M tris buffer (pH 7.6 or 7.4).

Preparation of ³H-Labeled RF DNA

E. coli HF 4704 (*hcr*⁻, thymine requiring) was grown in TPG-2A medium¹¹⁾ containing 5 µg/ml of thymine to about 1×10^8 cells/ml at 37°C with aeration. The cells were incubated after the addition of 30 µg/ml of mitomycin C for 10 minutes without aeration, collected by centrifugation and resuspended in the fresh medium containing 2 µg/ml of thymine. The cells were infected with ϕ X174 *am3* and at the same time 0.02 mc/ml of ³H-thymidine and 30 µg/ml of chloramphenicol were added. The infected cells were incubated further for 3 hours under vigorous aeration, then collected, washed and resuspended in 0.05 M tris buffer (pH 8.1). The cells were lysed by lysozyme-EDTA treatment¹⁰⁾, followed by freezing and thawing. The lysate was shaken with an equal volume of phenol saturated with 0.05 M tris buffer (pH 8.1) and the aqueous phase was recovered after separation by centrifugation. Remaining phenol in the aqueous phase was removed by the extraction with ether. After adding 0.1 volume of 3 M sodium acetate (pH 5.5), 2.5 volumes of isopropanol and 10 µg/ml of yeast tRNA, the solution was allowed to stand in a freezer overnight. The precipitate was collected by centrifugation.

Results


Effects of Bleomycin A₂ on Infectivity of Single-stranded Phage DNA to Spheroplast

If ϕ X174 phage DNA, single-stranded and closed, is opened by one strand scission or suffers chemical modification in some base residues by the action of hydroxylamine or other reagents, its infectivity to *E. coli* spheroplast is known to be lost. Therefore this spheroplast assay system gives a very sensitive detection method for DNA alteration. The effect of bleomycin A₂ on the infectivity of the single-stranded phage DNA to the spheroplast is shown in Fig. 1. The inactivation of phage DNA by 1 µg/ml of bleomycin A₂ was observed in the presence of neither 2-mercaptoethanol nor hydrogen peroxide.

Fig. 1. Inactivation of phage DNA by bleomycin A₂ as revealed by spheroplast assay.

The phage DNA (about 1×10^{-3} $\mu\text{g}/\text{ml}$) was incubated with bleomycin A₂ in 0.05 M tris buffer (pH 7.6) at 37°C. A portion of reaction mixture at the time indicated was transferred and diluted into ice-cold 0.05 M tris buffer (pH 8.1) containing yeast tRNA (2 $\mu\text{g}/\text{ml}$). Dilution was performed until the bleomycin present caused no effect on the spheroplast with respect to the infectivity of the phage DNA. The biological activity of the treated phage DNA was measured in the spheroplast assay as described by GUTHRIE *et al.*¹⁰⁾

- a. untreated b. bleomycin A₂ (5 $\mu\text{g}/\text{ml}$) c. bleomycin A₂ (1 $\mu\text{g}/\text{ml}$)
To the reaction mixture (c), a half volume of 10 mM EDTA was added after 30 minutes.



Degradation of Single-stranded or Double-stranded DNA by Bleomycin in the Absence of 2-Mercaptoethanol or Hydrogen Peroxide
Effects of three different types of bleomycin, A₂, A₅ and B₂, on the phage DNA were examined by means of sucrose density gradient centrifugation. As shown in

Fig. 2. Effects of bleomycin A₂ and A₂(Cu) and the inhibition of the bleomycin activity by EDTA.

The phage DNA (1×10^{-3} $\mu\text{g}/\text{ml}$) was incubated with bleomycin in 0.05 M tris buffer (pH 7.6) at 37°C for 90 minutes. The reaction mixture was layered on a 5~20% sucrose density gradient in Tris-saline-EDTA buffer and centrifuged as described in materials and methods. Each fraction was collected in a planchet, dried and counted in an Aloka GM counter.

- A. a. untreated b. bleomycin A₂(Cu) (5 $\mu\text{g}/\text{ml}$)
B. a. untreated b. bleomycin A₂ (5 $\mu\text{g}/\text{ml}$)
c. bleomycin A₂ (5 $\mu\text{g}/\text{ml}$)+EDTA (2.2 mM)




Fig. 3. Effects of various bleomycins on the sedimentation behaviour of the phage DNA.

The phage DNA (about $0.7 \mu\text{g/ml}$) was incubated with bleomycin in 0.05 M tris buffer (pH 7.4) at 37°C for 120 minutes. The reaction mixture was centrifuged and counted as described in the legend of Fig. 2.

- a. untreated b. bleomycin B_2 ($50 \mu\text{g/ml}$)
c. bleomycin A_5 ($50 \mu\text{g/ml}$)




Fig. 2B and 3, the phage DNA treated with bleomycins A_2 , A_5 and B_2 showed a decrease in the sedimentation velocity. The double-stranded RF DNA was also degraded by bleomycin as indicated in Fig. 4, where a specimen of bleomycin mixture was used. Fig. 5 presents the result of another experiment to reveal the degradation of phage DNA by bleomycin. The amount (as radioactivity) of TCA-insoluble portion of the DNA was found to decrease during the treatment with bleomycin mixture.

Action of Bleomycin in Cooperation with 2-Mercaptoethanol or Hydrogen Peroxide

The amount of TCA-insoluble portion from the DNA treated with bleomycin decreased more rapidly in the presence of 2-mercaptoethanol than in the absence of that (Fig. 5). The effect of hydrogen peroxide on the DNA degradation by bleomycin is illustrated in Fig. 6. The sedimentation pattern of phage DNA treated with $3.3 \mu\text{g/ml}$ of bleomycin A_2 , at which concentration only a little change was caused in the absence of hydrogen peroxide, showed a remarkable degradation upon the further addition of

Fig. 4. Effect of bleomycin on double-stranded RF DNA.

The RF DNA (about $3 \times 10^{-2} \mu\text{g/ml}$) was incubated with bleomycin in 0.05 M tris buffer (pH 7.6) at 37°C for 90 minutes. The DNA in the reaction mixture was precipitated in the cold with the addition of 0.1 volume of tRNA (1 mg/ml), 0.1 volume of 3 M sodium acetate (pH 5.5) and 2 volumes of ethanol and then redissolved in 0.05 M tris buffer (pH 7.6). The dissolved DNA was centrifuged and counted as described in materials and methods.

- a. untreated
b. bleomycin mixture ($50 \mu\text{g/ml}$)





Fig. 5. Effect of 2-mercaptoethanol on bleomycin activity.

The phage DNA (about $1 \times 10^{-2} \mu\text{g/ml}$) was incubated with bleomycin mixture in 0.05 M tris buffer (pH 7.6) at 37°C . A portion of the sample at the time indicated was added with carrier DNA to make a final concentration of $100 \mu\text{g/ml}$ and with cold TCA at a final concentration of 5%. The precipitates were collected on glass fibre paper, dried and counted in a scintillation counter.

- a. untreated
b. bleomycin mixture (3.4 mg/ml)
c. bleomycin mixture (3.4 mg/ml)
+ 2-mercaptoethanol (1 mM)



0.03 mM of hydrogen peroxide. The pattern of DNA treated with hydrogen peroxide only was the same as that of untreated DNA.

Ineffectiveness of Copper-chelated Bleomycin and Inhibition by EDTA of the Bleomycin Action

The copper-chelated bleomycin has been reported not to bring about the observable strand scission of DNA even in the presence of 2-mercaptoethanol. It is also known that the strand scission of DNA with bleomycin in the presence of 2-mercaptoethanol is inhibited by an addition of EDTA^{4,5,6,7,8}. We examined whether these effects held also for the bleomycin action on the phage DNA without 2-mercaptoethanol. As shown in Fig. 2A, copper-chelated bleomycin A₂ did not cause the observable degradation of DNA in the absence of 2-mercaptoethanol, while bleomycin A₂ freed from copper by the dithizon treatment recovered the activity (not indicated here). The activity of bleomycin A₂ was inhibited rapidly by the addition of EDTA in the absence of 2-mercaptoethanol as shown in the sedimentation (Fig. 2B) or in the spheroplast assay experiment (Fig. 1). These results are consistent with those previously reported.

Discussion


The biological activity of bleomycins has been discussed in connection with DNA degradation^{6,8}. ³H-Labeled bleomycin has been reported to bind to DNA and to cause strand scission and degradation of DNA in the presence of 2-mercaptoethanol or hydrogen peroxide but not in their absence *in vitro* even though it binds to DNA. These conclusions were deduced from the observed change in melting temperature or sedimentation patterns. In our studies it was found that, even in the absence of 2-mercaptoethanol or hydrogen peroxide, bleomycin(s) inactivated the biological activity of phage DNA and moreover degraded both the phage DNA and the double-stranded RF DNA. 2-Mercaptoethanol or hydrogen peroxide only enhanced the degradation of DNA by bleomycin. Consistent with the results of previous authors, the chelation of copper to bleomycin or the addition of EDTA to the reaction mixture inhibited the *in vitro* activity of bleomycin in the present system.

ϕ X174 Phage DNA and RF DNA used in our experiments possess several advantages for the detection of alteration in their molecular integrity. For example, the formation of only a single nick in the DNA strand may be easily revealed, because the circular closed native strand and the resulted linear open strand can be separated from each other by zone sedimentation. Moreover the spheroplast assay can afford a very sensitive detection method for DNA alteration as mentioned above¹². Presumably, the sensitivity of our

Fig. 6. Effect of hydrogen peroxide on bleomycin activity.

The phage DNA (about 1×10^{-3} μ g/ml) was incubated with bleomycin A₂ in 0.05 M tris buffer (pH 7.6) at 37°C for 70 minutes. The reaction mixture was centrifuged and counted as described in the legend of Fig. 2.

- untreated
- bleomycin A₂ (3.3 μ g/ml)
- bleomycin A₂ (3.3 μ g/ml)
+hydrogen peroxide (0.03 mM)
- hydrogen peroxide (0.03 mM)



system detects strand scissions not detected previously.

Although the biological activity of bleomycin *in vivo* might be considered in connection with sulfhydryl or peroxide compounds, their stimulation of DNA degradation by bleomycin seems not to be indispensable, at least *in vitro*.

References

- 1) UMEZAWA, H.; K. MAEDA, T. TAKEUCHI & Y. OKAMI : New antibiotics, bleomycin A and B. J. Antibiotics, Ser. A 19 : 200~209, 1966
- 2) UMEZAWA, H.; Y. SUHARA, T. TAKITA & K. MAEDA : Purification of bleomycins. J. Antibiotics, Ser. A 19 : 210~215, 1966
- 3) ISHIZUKA, M.; H. TAKAYAMA, T. TAKEUCHI & H. UMEZAWA : Activity and toxicity of bleomycin. J. Antibiotics, Ser. A 20 : 15~24, 1967
- 4) SUZUKI, H.; K. NAGAI, H. YAMAKI, N. TANAKA & H. UMEZAWA : Mechanism of action of bleomycins. Studies with the growing culture of bacterial and tumor cells. J. Antibiotics 21 : 379~386, 1968
- 5) SUZUKI, H.; K. NAGAI, H. YAMAKI, N. TANAKA & H. UMEZAWA : On the mechanism of action of bleomycin : Scission of DNA strands *in vitro* and *in vivo*. J. Antibiotics 22 : 446~448, 1969
- 6) NAGAI, K.; H. YAMAKI, H. SUZUKI, N. TANAKA & H. UMEZAWA : The combined effects of bleomycin and sulfhydryl compounds on the thermal denaturation of DNA. Biochim. Biophys. Acta 179 : 165~171, 1969
- 7) NAGAI, K.; H. SUZUKI, N. TANAKA & H. UMEZAWA : Decrease of melting temperature and single strand scission of DNA by bleomycin in the presence of 2-mercaptoethanol. J. Antibiotics 22 : 569~573, 1969
- 8) NAGAI, K.; H. SUZUKI, N. TANAKA & H. UMEZAWA : Decrease of melting temperature and single strand scission of DNA by bleomycin in the presence of hydrogen peroxide. J. Antibiotics 22 : 624~628, 1969
- 9) SUZUKI, H.; K. NAGAI, E. AKUTSU, H. YAMAKI, N. TANAKA & H. UMEZAWA : On the mechanism of action of bleomycin. Strand scission of DNA caused by bleomycin and its binding to DNA *in vitro*. J. Antibiotics 23 : 473~480, 1970
- 10) GUTHRIE, G. D. & R. L. SINSHEIMER : Observations on the infection of bacterial protoplasts with the deoxyribonucleic acid of bacteriophage ϕ X174. Biochim. Biophys. Acta 72 : 290~297, 1963
- 11) LINDQVIST, B. H. & R. L. SINSHEIMER : Process of infection with bacteriophage ϕ X174. XIV. Studies on macromolecular synthesis during infection with a lysis-defective mutant. J. Mol. Biol. 28 : 87~94, 1967
- 12) BURTON, A. & R. L. SINSHEIMER : The process of infection with bacteriophage ϕ X174. VII. Ultracentrifugal analysis of the replicative form. J. Mol. Biol. 14 : 327~347, 1965