

Ushikulides A and B, Immunosuppressants Produced by a Strain of *Streptomyces* sp.

Kosaku Takahashi, Teruhiko Yoshihara, Kazuhiko Kurosawa

Received: April 27, 2005 / Accepted: June 1, 2005

© Japan Antibiotics Research Association

Abstract Novel immunosuppressants, ushikulides A and B, were isolated from the culture broth of *Streptomyces* sp. IUK-102. Ushikulides A and B both have the same molecular formula, determined as $C_{40}H_{68}O_{10}$. The structures of both compounds were elucidated to be novel 22-membered macrolides. Both compounds showed immunosuppressive activity for murine splenocyte proliferation *in vitro*.

Keywords 22-membered macrolides, ushikulide, immunosuppressants

Cyclosporin A [1] and FK506 [2], extremely potent immunosuppressive agents, have revolutionized organ transplantation by blocking the activation of lymphocytes

and are widely used in preventing rejection. Cyclosporin A and FK506 respectively bind to cyclophilin and FK506-binding protein (FKBP), so-called immunophilin [3]. Binding of each of these drug-protein complexes to a catalytic subunit of calcineurin causes inhibition of signal transduction in T cell activation [3]. However, these agents have some adverse side-effects such as nephrotoxicity and chronic rejection [4, 5]. Another potent immunosuppressive agent is desired to develop a safer medical treatment. During our screening for immunosuppressants from microbial products, novel immunosuppressants ushikulides A (1) and B (2) were found in the culture broth of *Streptomyces* sp. IUK-102 (Fig. 1). This paper describes the isolation, structure elucidation, and biological activities of these compounds.

The producing strain, *Streptomyces* sp. IUK-102, was isolated from a soil sample collected in Ibaraki prefecture,




Fig. 1 Structures of ushikulides A (1) and B (2).

K. Takahashi (Corresponding author), **T. Yoshihara**: Graduate School of Agriculture, Hokkaido University, Sapporo, 060-8589, Japan, E-mail: kosaku@chem.agr.hokudai.ac.jp

K. Kurosawa: Department of Biology 68-370, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge MA 02139, USA

Japan. This strain was incubated at 27°C for five days in 500-ml Erlenmeyer flasks, each containing 70 ml of medium composed of (NH₄)₂SO₄ 0.14%, KH₂PO₄ 0.2%, CaCl₂ 0.03%, MgSO₄·7H₂O 0.03%, urea 0.03%, polypepton 0.5%, yeast extracts 0.1%, soybean meal 3%, glucose 1%, soluble starch 0.5%, FeSO₄·7H₂O 0.005%, MnSO₄·5H₂O 0.0016%, ZnSO₄·7H₂O 0.0014%, and CoCl₂ 0.002% (pH 7.0 before sterilization). After fermentation on a rotary shaker (230 rpm), four liters of the culture broth was centrifuged to separate mycelia and supernatant. The mycelia were extracted with acetone and the extract was evaporated to give an aqueous residue. This residue was then extracted with EtOAc and the soluble portion was concentrated *in vacuo* to obtain a brown residue (3.5 g). The residue was subjected to silica gel chromatography and eluted using CHCl₃ followed by a mixture of CHCl₃ and MeOH (90:10). Second elution with activity was evaporated and fractionated on a silica gel column using a mixture of CHCl₃ and MeOH (95:5). The fractions containing active substrates were combined and concentrated *in vacuo*. This active material was then subjected to reverse-phase HPLC (CAPCELL PAK C₁₈ SG 120, Shiseido Co. Ltd., 20×250 mm) using a mixture of CH₃CN and H₂O (70:30) as the isocratic mobile phase at a flow rate of 5.0 ml/minute. Ushikulides A (**1**) and B (**2**) were eluted at 32.1 and 30.0 minutes, respectively. These fractions were collected and concentrated under reduced pressure. Finally, **1** (13 mg) and **2** (5 mg) were obtained as white powders.

Ushikulide A (**1**) was obtained as a colorless powder with a specific rotation of -12.7° (c 0.5, MeOH, 25°C). The molecular formula of **1** was established as C₄₀H₆₈O₁₀ on the basis of HRFAB-MS (found 709.4910, calcd. 709.4883 for M+H). The IR spectrum (KBr) indicated the presence of hydroxyl and carbonyl groups at 3456 and 1719 cm⁻¹, respectively. From ¹H and ¹³C NMR spectral data, compound **1** was shown to possess a ketone, an ester carbonyl, 4 *sp*² methines, 2 *sp*³ quaternary carbons, 13 *sp*³ methines including 7 oxymethines, 11 *sp*³ methylenes, and 8 methyl carbons. Seven degrees of unsaturation were inferred from the molecular formula. The COSY and TOCSY spectra of **1** indicated four ¹H-¹H spin coupling systems (Fig. 2). Spin systems A and B were connected through a series of HMBC correlations from H-34 to C-9, C-10, and C-11, and from H₂-11 (δ_{H} 1.52/1.54) to C-10. Spin systems B and C were linked by HMBC correlations of H-16 and H-18 (δ_{H} 2.73) to C-17 (δ_{C} 212.5). The long range couplings from H-2 (δ_{H} 5.81) and H-21 (δ_{H} 5.33) to C-1 (δ_{C} 166.5) established the ester linkage of spin systems A and C, indicating the presence of a 22-membered lactone ring in the structure. The configurations of the two double




Fig. 2 2D-NMR correlations for **1**. Bold lines show TOCSY correlations, and arrows show HMBC correlations.

bonds in the structure were demonstrated as (*E*) by coupling constant values of H-2 to H-3 (15.7 Hz), and H-14 to H-15 (15.0 Hz). HMBC correlations from H-19, H-22, H-24, and H-27 to C-23 (δ_{C} 98.9) indicated connectivity between spin systems C and D and the presence of 1,1'-dioxaspiro[5.5]undecane, which was fused onto the 22-membered lactone structure. Accordingly, the structure of **1** was determined to be that shown in Fig. 1. All the assigned proton and carbon signals are listed in Table 1.

Ushikulide B (**2**) was obtained as a colorless powder with a specific rotation of -46.7° (c 0.5, MeOH, 25°C). The molecular formula of **2** was determined to be C₄₀H₆₈O₁₀ by HRFAB-MS (found 707.4728, calcd. 707.4734 for M-H), which was the same as that of **1**. The IR spectrum (KBr) indicated the presence of hydroxyl and carbonyl groups at 3437 and 1718 cm⁻¹, respectively. Comparison of the ¹H NMR spectra of **2** and **1** revealed that the double bond proton signals at δ_{H} 5.60 and δ_{H} 5.44 present in **1** were missing in **2**. Instead, a proton signal at δ_{H} 6.97 was identified in **2**. ¹³C NMR spectrum of **2** showed four carbon signals at δ_{C} 152.2, 149.4, 143.5, and 122.7 attributed to olefinic carbons. The significant difference observed between the NMR spectra resulted from a change in the olefin position. Besides this obvious difference, the 1D NMR spectra of these compounds resembled each other. Additionally, TOCSY and HMBC correlations of **2** (Fig. 3a) suggested that the basal structure of this compound was similar to that of **1**, however, more information was still required to establish the structure of **2**. Final structure confirmation was accomplished by a ROESY spectrum (Fig. 3b). The configuration of a double bond composed of C-15 and C-16 was shown to be (*E*) from ROESY correlation between H-15 and H-18 (δ_{H} 2.31). Correlation of H-27 with H-19 revealed a spiroketal

Table 1 NMR spectral data of ushikulides A (**1**) and B (**2**) in CD₃OD

No.	Ushikulide A (1)		Ushikulide B (2)	
	δ_{H} (J Hz)	δ_{C}	δ_{H} (J Hz)	δ_{C}
1		166.5		166.5
2	5.81 d (15.7)	122.7	5.80 d (15.8)	122.7
3	6.76 dd (15.7, 10.0)	152.2	6.75 dd (15.8, 10.0)	152.2
4	2.46 m	42.9	2.45 m	43.0
5	3.62 br.d (9.9)	81.3	3.59 dd (9.9, 1.7)	80.6
6	1.56 m	38.6	1.52 m	39.7
7	4.10 br.s	76.0	4.03 m	76.1
8	1.69 m	36.4	1.52 m	40.0
	1.73 m		1.69 m	
9	3.17 br.d (10.8)	75.0	3.28 (CD ₃ OD overlapped)	75.6
10		75.5		75.4
11	1.52 m	39.8	1.42 m	39.9
	1.54 m		1.57 m	
12	1.43 m	24.1	1.29 m	23.6
	1.52 m		1.47 m	
13	2.04 m	34.6	1.57 m	30.4
	2.23 m		1.63 m	
14	5.60 ddd (15.0, 9.4, 4.2)	136.7	2.37 m	29.8
			2.44 m	
15	5.44 dd (15.0, 9.0)	129.6	6.97 dd (8.7, 5.1)	149.4
16	3.07 ddd (8.6, 8.6, 6.4)	57.8		143.5
17		212.5		203.5
18	2.73 dd (17.4, 8.0)	42.0	2.31 dd (13.3, 4.8)	39.2
	2.78 dd (17.8, 6.4)		3.60 dd (13.3, 10.9)	
19	4.33 br.t (6.3)	67.2	4.19 m	70.2
20	2.05 m	34.7	1.85 m	34.7
21	5.33 ddd (11.1, 5.4, 5.4)	71.3	5.21 ddd (10.6, 4.9, 4.9)	71.4
22	1.68 m	36.4	1.72 m	36.3
	1.72 m			
23		98.9		99.4
24	1.43 m	30.6	1.47 m	30.5
	1.65 m		1.67 m	
25	1.40 m	27.5	1.42 m	27.6
	2.19 m		2.14 m	
26	1.48 m	32.1	1.59 m	32.1
27	4.02 m	69.1	4.09 m	69.3
28	1.28 m	41.7	1.28 m	41.4
	1.63 m		1.62 m	
29	3.73 br.s	70.6	3.78 m	70.4
30	1.47 m	31.9	1.49 m	32.2
	1.53 m		1.54 m	
31	0.97 t (7.4)	10.2	1.00 t (7.4)	10.3
32	1.14 d (6.4)	18.0	1.13 d (6.6)	18.0
33	0.85 d (6.9)	4.2	0.85 d (6.9)	4.9
34	1.06 s	22.7	1.08 s	23.2
35	1.41 m	41.3	2.22 d (7.1)	35.3
	1.53 m			
36	1.53 m	26.7	1.63 m	29.3
37	0.91 d (5.9)	23.2	0.86 d (5.4)	23.0
38	0.87 d (6.2)	22.4	0.82 d (6.6)	22.9
39	0.79 d (6.9)	6.2	0.92 d (6.9)	6.1
40	0.94 d (6.9)	11.5	0.95 d (6.9)	11.5

¹H and ¹³C NMR were observed at 500 and 125 MHz, respectively.




Fig. 3 2D-NMR correlations for **2**: (a) TOCSY (bold lines) and HMBC (arrows) correlations, (b) key ROESY correlations for spiroketal moiety.

subunit connecting spin systems C and D. Moreover, additional ROESY cross peaks were observed as shown in Fig. 3b. The configuration of the spiroketal in **2** was quite similar to that in cytovaricin [6], therefore, the structure of **2** was determined as that in Fig. 1. The complete assignments of proton and carbon signals in **1** and **2** revealed the considerable difference of proton chemical shifts of H₂-18 between these compounds (Table 1). This suggested that the positional change of the olefin near the ketone group caused a conformation of 22-membered lactone ring to be changed. As a result, H₂-18 at δ_{H} 2.31 and 3.60 in **2** were placed in a shielding area of the ketone group and a deshielding area of the ketone group, respectively.

The immunosuppressive activities of these compounds were estimated by concanavalin A (ConA) stimulated lymphocyte blastogenesis [7]. Both compounds inhibited murine splenic lymphocyte proliferation with IC₅₀ values of 70 nM, comparable to that of cyclosporin A. Compounds **1** and **2** did not show significant cytotoxicity for KB cells at a concentration of 10 μM .

Ushikulides A (**1**) and B (**2**) are related to phthoramycin [8], kaimonolide A [9], and the aglycone of cytovaricin [6, 10], which are thought to be members of the oligomycin family from the point of view of biosynthesis. Oligomycin F [11] and 41-demethylhomooligomycin B [12], which have a 26-membered lactone ring in their structures, are known to possess immunosuppressive activities. As far as we know, this is the first report describing a 22-membered macrolide with a spiroketal moiety that shows immunosuppressive activity.

Acknowledgement We thank Dr. E. Fukushi and Mr. K.

Watanabe for measurements of mass spectral data.

References

- Dreyfuss M, Harri E, Hoffmann H, Kobel H, Pache W, Tschertter H. Cyclosporin A and C. New metabolites from *Trichoderma polysporum* (Linx ex Pers.) Rifai. *Eur J Appl Microbiol* 3: 125–133 (1976)
- Kino T, Hatanaka H, Hashimoto M, Nishiyama M, Goto T, Okuhara M, Kohsaka M, Aoki H, Imanaka H. FK-506, a novel immunosuppressant isolated from a *Streptomyces*. I. Fermentation, isolation, and physico-chemical and biological characteristics. *J Antibiot* 40: 1249–1255 (1987)
- Schreiber SL, Crabtree GR. The mechanism of action of cyclosporin A and FK506. *Immunology Today* 13: 136–142 (1992)
- The US Multicenter FK506 Liver Study Group. A comparison of tacrolimus (FK506) and cyclosporin for immunosuppression in liver transplantation. *N Engl J Med* 331: 1110–1115 (1994)
- Atkinson P, Joubert G, Barron A, Grant D, Paradis K, Seidman E, Wall W, Rosenberg H, Howard J, Williams S, Stiller C. Hypertrophic cardiomyopathy associated with tacrolimus in paediatric transplant patients. *Lancet* 345: 894–896 (1995)
- Evans DA, Skalder SW, Jones TK, Clardy J, Stout TJ. Total synthesis of the macrolide antibiotic cytovaricin. *J Am Chem Soc* 112: 7001–7031 (1990)
- Kurosawa K, Takahashi K, Tsuda E. SNF4435C and D, novel immunosuppressants produced by a strain of *Streptomyces spectabilis*. I. Taxonomy, fermentation, isolation and biological activities. *J Antibiot* 54: 541–547 (2001)
- Nakagawa A, Miura S, Imai H, Imamura N, Omura S.

- Structure and biosynthesis of a new antifungal antibiotic, phthoramycin. *J Antibiot* 42: 1324–1327 (1989)
9. Hirota A, Okada H, Kanza T, Nakayama M, Hirota H, Isogai A. Structure of kaimonolide A, a novel macrolide plant growth inhibitor from a *Streptomyces* strain. *Agric Biol Chem* 53: 2831–2833 (1989)
 10. Kihara T, Kusakabe H, Nakamura G, Sakurai T, Isono K. Cytovaricin, a novel antibiotic. *J Antibiot* 34: 1073–1074 (1981)
 11. Laatsch H, Kellner M, Wolf G, Lee YS, Hansske F, Konetschny-Rapp S, Pessara U, Scheuer W, Stockinger H. Oligomycin F, a new immunosuppressive homologue of oligomycin A. *J Antibiot* 46: 1334–1341 (1993)
 12. Kim HS, Han SB, Kim HM, Kim YH, Lee JJ. 41-Demethylhomooligomycin B, a new immunosuppressant antibiotic from *Streptomyces ostreogriseus*. *J Antibiot* 49: 1275–1277 (1996)