

## A New Fungicide Produced by a *Streptomyces* sp. GAAS7310

Guangying Chen, Birun Lin, Yongcheng Lin, Fengchun Xie, Wen Lu, Wang-Fun Fong

Received: April 22, 2005 / Accepted: July 16, 2005

© Japan Antibiotics Research Association

**Abstract** Directed bioassay guided fraction led to a new macrodiolide antimycin A<sub>17</sub> (**1**), isolated from a *Streptomyces* sp. GAAS7310, which showed significant antimicrobial activity against eleven fungal species, including *Curvularia lunata* (Wakker) Boed, *Rhizopus nigricans* Ehrb and *Colletotrichum nigrum* EL. et Halst. The structure was unambiguously established by interpretation of 1D and 2D NMR data and comparison with the known antimycin A<sub>1a</sub>.

**Keywords** *Streptomyces* sp.; antimycin; antimicrobial activity; fungicide; macrodiolide; actinomycete

### Introduction

Actinomycetes are prolific producers of antibiotics and other industrially useful secondary metabolites [1–5]. Because these metabolites are often active against a limited number of specific target species, biodegradable to nontoxic products, and potentially suitable for use in integrated management programs, they can lead to the development of new classes of safer disease control agents. In our course of screening for novel naturally occurring fungicides from actinomycetes, the acetone extract of the mycelium of a *Streptomyces* sp. 7310# collected at Tianshan mountain was found to have strong activity against 12 fungal strains, e.g. *Phomopsis vexans* (Sacc et Syd) Harter etc. By bioassay guided fractionation, we isolated a new macrodiolide, antimycin A<sub>17</sub> (**1**), which showed activity against several fungi. Among these fungi,

*Curvularia lunata* (Wakker) Boed, *Rhizopus nigricans* Ehrb and *Colletotrichum nigrum* EL. et Halst were more sensitive to antimycin A<sub>17</sub> (**1**).

### Materials and Methods

#### General Experimental Procedure

Melting point was determined on a XT-4 apparatus and was uncorrected. NMR spectra were recorded on a Varian Inova-500 NMR spectrometer (500 MHz) using acetone-*d*<sub>6</sub> as solvent and TMS as an internal standard. Mass spectra were measured on a VG-ZAB-HS analytical spectrometer. IR spectra were taken on a Nicolet 5DX-FTIR spectrophotometer. Optical rotation was determined in acetone at room temperature using a Horiba High Sensitivity SEPA-300 Polarimeter, and elemental analysis was recorded on an Elementar Vario EL CHNS-O elemental analysis device. Preparative HPLC were performed on an Waters 2695 system equipped with an autosampler, a quaternary pump system, a photodiode array detector, and a Chemstation data system. A 250 mm × 20 mm i.d., 5 μm particle size Zorbax Eclipse XDB ODS column (Agilent) was selected for HPLC preparation. Analytical thin-layer chromatography (TLC) was carried out on precoated silica gel aluminum plates (Merck, 60 F<sub>254</sub>, 0.2 mm thickness) and HPTLC plates (Merck, RP-18). Silica gel (200–300 mesh, Qingdao Haiyang Chemical Co., Ltd.), RP-18 silica gel (50 μm, Merck), and CAD-45 absorber resin (26–60 mesh, Anhui Sanxing resin technology Co., Ltd.), were used for column

**G. Chen:** Department of Chemistry, Hainan Normal University, Haikou 571158, People's Republic of China

**Y. Lin** (Corresponding author), **W. Lu:** Department of Applied Chemistry, Zhongshan University, Guangzhou 510275, People's Republic of China, E-mail: yclin48@hotmail.com

**B. Lin, F. Xie:** Institute of Plant-protect, Guangdong Academy of Agricultural Science, Guangzhou 510000, People's Republic of China

**G. Chen, W.-F. Fong:** Department of Biology & Chemistry, City University of Hong Kong, Hong Kong

chromatography. The solvents used for high-performance liquid chromatography (HPLC) were of HPLC grade (Sigma); and other solvents and chemicals were of analytical grade.

### Collection and Cultivation

The bacterium *Streptomyces* sp. GAAS7310 was isolated from the soil collected at Tianshan mountain, Xinjiang Province, P.R. China. It has been deposited in the China Center for Type Culture Collection (CCTCC), Wuhan, P.R. China. Deposited number was CCTCC M 203058 *Streptomyces* sp. GAAS7310. Starter cultures were maintained on cornmeal agar. Plugs of agar supporting mycelial growth were cut and transferred aseptically into 250 ml Erlenmeyer flasks containing 100 ml of liquid medium (starch 1%, calcium carbonate 0.2%, ammonium sulfate 0.2%, potassium dihydrogen phosphate 0.1%, magnesium bisulfate 0.1%, sodium chloride 0.1%, yeast extract 1%, pH 7.2). The content were incubated at 28°C on a rotary shaker for 5~7 days, and the mycelium was aseptically transferred to a 500-liter fermentor containing the same culture liquid (300 liters) with the following settings: air, 1.8 liters/minute; impellor speed, 200 rpm for 3 days.

### Extraction and Isolation

The 300-liter culture was filtered through Celite to separate the mycelium and culture liquid. The acetone extract of mycelium exhibited the strong activity against twelve fungal species. The 85% aqueous acetone extract of 900 g mycelium was adsorbed onto a macroporous resin (CAD-45, 50 cm×9 cm i.d.), followed by on elution with water, then ethyl acetate respectively. The ethyl acetate fraction was evaporated in vacuum at 45°C and yielded 15 g pale-yellow oil, then subjected to column chromatography over silica gel (200 g, 200~300 mesh, 80 cm×3.5 cm i.d.) and eluted with petrol (bp 60~90°C)-EtOAc mixtures of increasing 10% polarity. The 30% fraction showed strong fungicidal activities. This fraction (1.5 g) was further purified by means of a silica gel column (80 g, 200~300 mesh, 50 cm×2.5 cm i.d.) using chloroform/methanol (95:5) as the solvent system to yield a bioactive mixture. Further purified by HPLC (a 250 mm×20 mm i.d., 5 μm particle size Zorbax Eclipse XDB ODS column) with methanol/water (70:30) as the mobile phase to yield the active component **1** (retention time 15.1 minutes, 25 mg). The molecular structure of compound **1** was elucidated by means of mass, IR, and 1D and 2D NMR spectroscopic techniques as the new macrodiolide antimycin A<sub>17</sub>.

Compound **1**: colorless crystalline solid (65 mg). mp

140~142°C.  $[\alpha]_D^{20} +114.7$  (*c* 0.002, acetone); IR (KBr)  $\nu_{\max}$ : 3350, 2960, 1797, 1742, 1679, 1638, 1532, 1455, 1367, 1252, 1178, 1075, 875 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR (see Table 1); FAB-MS *m/z*: 535 (M+H)<sup>+</sup>, 521, 492, 451, 372, 265, 164, 107, 102; Anal. (%): found C 60.75, H 7.15, N 5.25, O 26.85; calcd. C 60.66, H 7.16, N 5.24, O 26.9.

### Fungicidal Activity *In Vitro*

Antifungal activities of antimycin A<sub>17</sub> were investigated by disk assay method, using twelve representative fungi. These fungi were grown in potato dextrose agar (PDA) media, at 28°C. The PDA plates were prepared with antimycin A<sub>17</sub> (at a concentration of 100 μg/ml). Then, disks (6.0 mm in diameter) of epiphyte were placed on the plates. Similarly, disks of epiphyte were also placed on the innocuous plates (adding the same dosage sterile water) as blank comparison. All plates were incubated at 28°C for 1~3 days. Each fungus had five repeats. The results are shown in Table 2. Among these fungi, *R. nigricans*, *Cladosporium* sp. and *C. nigrum* were most sensitive.

## Results and Discussion

The acetone extract of the mycelium was adsorbed by CAD-45, followed by washing with water and EtOAc successively. The EtOAc extract was concentrated *in vacuo*, then chromatographed on silica gel column and RP-HPLC as monitored by bioassay against fungi. This afforded the active compound antimycin A<sub>17</sub> (**1**) and no other antimycins were obtained.

The molecular formula of **1** was determined to be C<sub>27</sub>H<sub>38</sub>N<sub>2</sub>O<sub>9</sub> by elemental analysis and FABMS data. UV absorptions at 328 and 212 nm were assigned to a chromophore with extended conjugation. Analysis of the <sup>13</sup>C NMR spectrum (Table 1) showed five carbonyl carbon signals ( $\delta$  175.7, 174.1, 170.9, 170.3, 160.7), six olefinic carbon signals ( $\delta$  151.6, 128.6, 125.2, 122.6, 119.3, 114.5). These accounted for eight of the ten degrees of unsaturation required by the molecular formula, therefore illustrating compound **1** to be bicyclic.

The <sup>1</sup>H NMR signals at  $\delta$  8.47 (1H, dd, *J*=8, 1.5 Hz, H-6'),  $\delta$  7.77 (1H, dd, *J*=8, 1.5 Hz, H-4') and  $\delta$  6.92 (1H, dd, *J*=8, 8 Hz, H-5') indicated the presence of an ortho-trisubstituted benzene ring. In the H-H COSY, the proton correlations of NH-3' and H-8', and in the HMBC the correlations of the proton of NH-3' and C-8' suggested the presence of a NH-CHO moiety. A pentyl side chain was revealed by the contiguous correlations from H-12 to H-16 in <sup>1</sup>H-<sup>1</sup>H COSY and correlations from H-12 to C-6 and C-7 in HMBC. A contiguous sequence from H-18 to H-20 and



**Table 1** NMR data of antimycin A<sub>17</sub> (actone-*d*<sub>6</sub>, TMS)

No.	<sup>13</sup> C	<sup>1</sup> H (mult, J)	COSY	HMBC
2	170.3			
3	54.7	5.41 (dd, 8.0, 7.0 Hz)	NH-3, H-4	2, 4
4	72.2	5.70 (dq, 7.0, 7.0 Hz)	H-3, H-11	2, 6
6	174.1			
7	51.1	2.55 (m)	H-8	6, 12
8	76.3	5.0 (m)	H-7, 9	6, 17
9	74.8	4.9 (m)	H-8, H-10	2, 8, 10
10	18.2	1.30 (d, 7.0 Hz)	H-9	8, 9
11	15.4	1.40 (d, 7.0 Hz)	H-4	3, 4
12	29.7	1.29 (m), 1.63 (m)	H-13, 7	6, 7
13	23.1	0.95 (m), 1.12 (m)	H-12, 14	15
14	36.7	1.28 (m)	H-13, 15	16
15	28.6	1.21 (m)	H-14, 16	
16	13.9	0.86 (t, 7.0 Hz)		
17	175.7			
18	41.8	2.47 (m)	H-19, 21	19, 20
19	27.2	1.51 (m), 1.74 (m)	H-18, 20	17, 18
20	11.9	1.16 (t, 7.0 Hz)	H-19	
21	17.0	1.17 (d, 7.0 Hz)	H-18	17, 18
1'	114.5			
2'	151.6			
3'	128.6			
4'	125.2	8.47 (dd, 8.0, 1.5 Hz)	H-5'	2', 6'
5'	119.3	6.92 (dd, 8.0, 8.0 Hz)	H-4', 6'	1', 3', 4', 6'
6'	122.6	7.77 (dd, 8.0, 1.5 Hz)	H-5'	2', 4', 5', 7'
7'	170.9			
8'	160.7	8.52 (d, 1.5 Hz)	3'-NH	3'
NH-3'	9.09 (s)		H-8'	2', 3', 8'
NH-3	8.31 (d, 8.0 Hz)		H-3	3, 7'
OH-2'	12.83 (s)			

H-21 in <sup>1</sup>H-<sup>1</sup>H COSY, and by combining the HMBC correlations between H-8, H-19 and C-17, and between H-21 and C-17, C-18 respectively, established the presence of a 2-methyl butanoyl group.

The overall structure of **1** was unambiguously established by 2D NMR spectra including COSY, HMQC and HMBC (Table 1 and Fig. 1). This structure was also supported by the mass fragment patterns (Fig. 2).

In 1949, Dunshee *et al.* first isolated antimycins A from *Streptomyces* fermentation products [6]. They thought the products were consisted of at least four closely related compounds, differing only in the substituents at the 7- and 8-positions of the dilactone ring. Subsequently Abide *et al.* isolated the mixture of antimycins A into eight hitherto subcomponents, A1a, A1b, A2a, A2b, A3a, A3b, A4a and A4b by HPLC [7] and identified the structure of A1a by

**Fig. 1** Selected HMBC correlations of **1**.**Fig. 2** Selected mass fragments of **1**.

1D, 2D NMR and X-ray [8, 9]. Tsunoda *et al.* had accomplished the total synthesis of antimycin A<sub>3b</sub> [10]. Compared with known antimycin A series, compound **1** is different at the 7-position. In the antimycin A series, the 7-position substituent is *n*-hexyl or *n*-butyl; but in **1** is *n*-pentyl. Thus we can name **1** as antimycin A<sub>17</sub> [11].

In our present study, a wide range of microorganisms was examined including 12 kinds of representative fungi. It was found that antimycin A<sub>17</sub> has broad inhibitory activity to microorganisms. Especially to three fungi, *Curvularia lunata* (Wakker) Boed, *Rhizopus nigricans* Ehrb and *Colletotrichum nigrum* EL. *et* Halst antimycins A<sub>17</sub> showed more sensitivity (Table 2). The bioassay results suggest that antimycin A<sub>17</sub> may be an ideal candidate as a food preservative, pharmaceutical and natural microorganism-based product.

**Acknowledgments** This work was supported by the National Natural Science Foundation of China (20072058), the 863 Foundation of China (2003AA624010), the Natural Science Foundation of Guangdong Province, China (980317).

**Table 2** The antimicrobial activity study of antimycin A<sub>17</sub> against 12 fungal strains

Fungi	Average diameter of radial growth (cm)		Rate of inhibition* (%)
	CK	Treatments	
<i>Rhizopus nigricans</i> Ehrb	7	3.5	50
<i>Geotrichum candidum</i> Link	6.9	6.55	5.1
<i>Fusarium oxysporum</i> f.sp. <i>cubense</i>	4.4	4.1	6.8
<i>Cladosporium carpophilum</i> Thum.	3.4	3.5	-2.9
<i>Alternaria solani</i> (E. et M.) Jones et Grout	5.3	3.8	28.3
<i>Cladosporium</i> sp.	1.5	1.4	66.7
<i>Sclerotinia sclerotiorum</i> (Lib.) de Bary	2	1.5	25
<i>Peronophthora litchi</i> Chen	2.8	2.5	10.7
<i>Botrytis</i> sp.	1.4	1.2	14.3
<i>Colletotrichum nigrum</i> EL.et Halst	2.8	1.4	50
<i>Curvularia lunata</i> (Wakker) Boed	3	2.2	26.7
<i>Phomopsis vexans</i> (Sacc et Syd) Harter	2.5	2.3	8

\* Rate of inhibition% = [(CK average diameter of radial growth - treatments average diameter of radial growth) / CK average diameter of radial growth] × 100.

## References

- Osada H. Actinomycetes: how fascinating microorganisms. *Actinomycetologica* 12: 85-88 (1998)
- Takahashi Y, Ōmura S. Isolation of new actinomycete strains for the screening of new bioactive compounds. *J. General Applied Microbiology* 49: 141-154 (2003)
- Miyadoh S. Research on antibiotic screening in Japan over the last decade: a producing microorganism approach. *Actinomycetologica* 7: 100-106 (1993)
- Shiomi K, Hatae K, Hatano H, Matsumoto A, Takahashi Y, Jiang CL, Tomoda H, Kobayashi S, Tanaka H, Ōmura S. A new antibiotic, antimycin A9, produced by *Streptomyces* sp. K01-0031. *J Antibiot* 58: 74-78 (2005)
- Barrow CJ, Oleynek JJ, Marinelli V, Sun HH, Kaplita P, Sedlock DM, Gillum AM, Chadwick CC, Cooper R. Antimycins, inhibitors of ATP-citrate lyase, from a *Streptomyces* sp. *J Antibiot* 50: 729-733 (1997)
- Dunshee BR, Leben C, Keitt GW, Strong FM. The isolation and properties of antimycin A. *J Am Chem Soc* 71: 2436 (1949)
- Abidi SL. High-performance liquid chromatographic separation of subcomponents of antimycin A. *J Chromatography* 447: 65-79 (1988)
- Abidi SL, Adams BR. Proton and carbon-13 resonance designation of antimycin A<sub>1</sub> by two-dimensional NMR spectroscopy. *Magnetic Resonance in Chemistry* 12: 1078-1080 (1987)
- Kim H, Esser L, Hossain MB, Xia D, Yu CA, Rizo J, Helm D, Deisenhofer J. Structure of antimycin A<sub>1</sub>, a specific electron transfer inhibitor of ubiquinol-cytochrome *c* oxidoreductase. *J Am Chem Soc* 121: 4902-4903 (1999)
- Tsunoda T, Nishii T, Yoshizuka M, Yamasaki C, Suzuki T, Ito S. A total synthesis of (-)-antimycin A<sub>3b</sub>. *Tetrahedron Letters* 41: 7667-7671 (2000)
- Hosotani N, Kumagai K, Nakagawa H, Shimatani T, Saji I. Antimycins A<sub>10</sub>-A<sub>16</sub>, seven new antimycin antibiotics produced by *Streptomyces* spp. SPA-10191 and SPA-8893. *J Antibiot* 58: 460-467 (2005)