

A New Series of Glycopeptide Antibiotics Incorporating a Squaric Acid Moiety

Synthesis, Structural and Antibacterial Studies[†]

Ferenc Sztaricskai, Gyula Batta, Pál Herczegh, Attila Balázs, József Jeko, Erzsébet Roth, Pál T. Szabó, Szilvia Kardos, Ferenc Rozgonyi, Zoltán Boda

Dedicated to the memory of Professor Kenneth L. Rinehart.

Received: May 8, 2006 / Accepted: September 14, 2006
© Japan Antibiotics Research Association

Abstract The aglycones of the antibiotics eremomycin, vancomycin and ristocetin (**3**, **4** and **6**, respectively) were prepared by deglycosidation of the parent antibiotics with hydrogen fluoride, and complete assignment of their ¹H, ¹³C and ¹⁵N spectra was performed. The squaric acid amide esters (**11**~**14**), were prepared from dimethyl squarate. The corresponding asymmetric diamides (**16**~**19**, **22**, **23**) were also synthesized using 4-phenylbenzylamine and triglycine. The advantage of the method is the high regioselectivity and that no protecting group strategy is required. Electrospray mass spectroscopic method was elaborated for the determination of the site of substitution of the modified antibiotics. The antibacterial activity of the prepared compounds is discussed in detail.

Keywords synthesis, glycopeptide antibiotics, squaric acid amides, antibacterial

Introduction

During the past fifty years the chemistry, biology, and therapeutic application of the glycopeptide antibiotics have been continuously progressing [1, 2]. In each of these antibiotics the specific heptapeptide aglycone is substituted at the alcoholic and/or phenolic hydroxyl groups with aminodeoxy sugars and simple, neutral carbohydrates. Based on the substitution pattern, mono-, bis-, tris-, and tetraglycoside-type glycopeptide antibiotics are classified [3], of which the bis- and tris-glycosides are the most frequently occurring. Until now, no C- or N-glycosidic bond, or a glycosylamine structural unit were detected in the molecules of the glycopeptide antibiotics.

These antibiotics exert biological activity by developing hydrogen bond between the glycopeptide backbone and the L-Lys-D-Ala-D-Ala terminal monomer fragment of the peptidoglycan constituent in the bacterial cell wall. This interaction inhibits the transglycosidation and transpeptidation processes which leads to interruption of

F. Sztaricskai (Corresponding author), **G. Batta**, **P. Herczegh**, **A. Balázs**, **J. Jeko**, **E. Roth**: Research Group for Antibiotics of the Hungarian Academy of Sciences, and Department of Pharmaceutical Chemistry, University of Debrecen, H-4010 Debrecen, P.O.Box 70, Hungary, E-mail: sztari@delphin.unideb.hu
P. T. Szabó: Department of Biochemical Pharmacology, Chemical Research Center, H-1525 Budapest, P.O. Box 17, Hungary

S. Kardos, **F. Rozgonyi**: Institute of Medicinal Microbiology, Semmelweis University, H-1445, Budapest, P.O.Box 370, Hungary
Z. Boda: IInd Department of Medicine, University of Debrecen, H-4012 Debrecen, P.O.Box 58, Hungary

[†]A part of this work was presented at the 9th International Conference on the Chemistry of Antibiotics and other Bioactive Compounds, Bordeaux-Arcachon, France, 25-29 September, 2005. (Abstract No 30)

the cell-wall synthesis and destruction of the bacteria. Interestingly, many glycopeptide antibiotics [vancomycin (2), eremomycin (1), ristocetin A (5), etc.] form dimeric complexes in aqueous solution [4].

Preparation of the aglycones of the glycopeptide antibiotics has been attempted by means of chemical methods, *i.e.* by acid hydrolysis of the parent antibiotics under various conditions. However, these procedures generally led to extremely complex reaction mixtures and low yields. The hydrogen fluoride-method was used by Wanner [5] in glycopeptide chemistry for the deglycosidation of vancomycin, ristocetin and ramoplanin.

In the present work we describe the results obtained by repeating deglycosidation of vancomycin and ristocetin A with hydrogen fluoride, and this procedure was also extended to eremomycin. For comparison of the NMR spectra, the aglycone of teicoplanin was also prepared by acid hydrolysis (Fig. 1). The synthesis, physico-chemical

properties, mass and NMR spectral characteristics and antibacterial activity of the new semisynthetic glycopeptide antibiotics and their aglycones prepared with dimethyl squarate are also presented.

Results and Discussion

Deglycosidation

The cleavage of the *O*-glycosidic bond of the glycopeptide antibiotics with hydrogen fluoride is greatly influenced by several factors, including the type and number of places of attachments of the neutral sugars and aminodeoxy hexoses—either as a monosaccharide unit or an oligosaccharide chain—to the heptapeptide aglycones.

We observed that deglycosidation of the monoglycoside-type vancomycin (2) and of the bis-glycoside-type eremomycin (1) with HF/anisole under the conditions suggested by the American researchers [5] (Scheme 1) furnished the homogeneous aglyco-vancomycin (4) and aglyco-eremomycin (3) with good yields. The purity of the products was proved by TLC, HPLC, and MALDI-TOF (Table 1), and the structures were assigned according to the ¹H and ¹³C NMR data (Table 2a~b). The yields of the reactions were greatly influenced by the temperature and volume of the EtOAc-MeOH mixture (4:1) used for washing in work-up procedure.

In contrast to the data described for 1 and 2, deglycosidation of the tris-glycoside-type ristocetin A (5), containing six carbohydrate units, with anhydrous HF-anisole mixture (Scheme 2) resulted in 3~4 component reaction mixtures in each case. Purification of the




Fig. 1 Structure of aglyco-teicoplanin.

Table 1 Physico-chemical properties of the aglycones


Name of the aglycones	Yield %	HPLC R _t min	TLC R _f	Molecular weight		[α] _D ²⁰	UV λ _{max} nm		
				Formula of the aglycones	Calculated				
3	Eremomycin	57.8	13.70	(1 A) 0.51	C ₅₃ H ₅₃ O ₁₇ N ₈ Cl	1109	1131.5	92.1 (c=0.10 MeOH)	279 (MeOH)
4	Vancomycin	89.87	14.68	(1 A) 0.62	C ₅₃ H ₅₂ O ₁₇ N ₈ Cl ₂	1143	1165.3	55.8 (c=0.26 DMF)	281 (DMF)
6	Ristocetin-A	52.8	14.60	(1 A) 0.63 (2 B) 0.26	C ₆₀ H ₅₁ O ₁₉ N ₇	1173	1196.4	-55.3 (c=0.20 MeOH)	279 (MeOH)

TLC: (1) DC Cellulose
(2) Kieselgel 60F₂₅₄

Solvent system: (A) *n*BuOH-Pyr-AcOH-H₂O (15:10:3:12)


(B) Toluene-MeOH-AcOH (1:1:0.05)

Detection: Pauly-reagent



	X	X ₁	Z	R	R ₁	R ₂	R ₃	R ₄
1 Eremomycin	H	Cl	Me	NH ₂	<i>i</i> -Pr			H
2 Vancomycin	Cl	Cl	Me	NH ₂	<i>i</i> -Pr		H	H

HF/anisole
RT, 2 h



	X	X ₁	Z	R	R ₁
3 Aglyco-eremomycin	H	Cl	Me	NH ₂	
4 Aglyco-vancomycin	Cl	Cl	Me	NH ₂	

Scheme 1 Deglycosidation of eremomycin and vancomycin [5].

Table 2 NMR assignments of the the glycopeptide aglycones(a) $^1\text{H}/^{13}\text{C}$ chemical shift assignment of the aglycones of eremomycin and vancomycin. Solvent reference signals were used for ^{13}C calibration: DMSO=39.51 ppm

Aglycoeremomycin, DMSO, 340 K			Aglycovancomycin, DMSO, 330 K		
Assignment	^1H (ppm)	^{13}C (ppm)	Assignment	^1H (ppm)	^{13}C (ppm)
1a	1.442/1.520	41.51	1a	1.474/1.556	40.62
1b	1.791	24.23	1b	1.741	24.09
1c	0.908	22.13	1c	0.926	22.74
1d	0.937	22.89	1d	0.889	22.31
1e	2.338	34.09	1e	2.378	33.28
2a	—	139.79	2a	—	139.66
2b	7.464	128.35	2b	7.430	128.45
2c	—	129.91	2c	—	126.94
2d	—	150.37	2d	—	150.37
2e	7.243	124.2	2e	7.280	124.31
2f	7.542	126.63	2f	7.548	126.80
3a	2.149/2.644	36.27	3a	2.170/2.542	36.78
4a	—	129.91	4a	—	128.73
4b	5.404	106.31	4b	5.228	104.43
4c	—	147.88	4c	—	147.70
4d	—	134.41	4d	—	133.92
4e	—	149.3	4e	—	149.02
4f	5.617	106.58	4f	5.633	107.02
5a	—	126.93	5a	—	126.19
5b	7.160	135.8	5b	7.172	135.64
5c	—	126.34	5c	—	121.69
5d	—	154.57	5d	—	154.95
5e	6.714	116.24	5e	6.741	116.25
5f	6.789	125.35	5f	6.802	125.37
6a	—	139.69	6a	—	141.99
6b	7.507	128.11	6b	7.854	127.46
6c	7.093	120.63	6c	—	126.10
6d	—	154.39	6d	—	147.49
6e	6.888	121.65	6e	7.231	123.12
6f	7.637	126.73	6f	7.471	127.12
7a	—	136.57	7a	—	136.30
7b	—	118.2	7b	—	117.99
7c	—	156.28	7c	—	156.34
7d	6.390	102.36	7d	6.418	102.27
7e	—	157	7e	—	157.07
7f	6.363	106.19	7f	6.337	106.01
X1	3.024	62.33	X1	3.217	61.69
X2	4.715	58.53	X2	4.825	58.52
X3	4.309	51.08	X3	4.371	51.00
X4	5.675	55.1	X4	5.713	54.64
X5	4.547	53.58	X5	4.465	53.69
X6	4.211	62.15	X6	4.200	61.71
X7	4.491	56.88	X7	4.488	56.84
y1	—	174.3	y1	—	173.22
y2	—	171.8	y2	—	171.37
y3	—	169.87	y3	—	170.19

Table 2 (a) (continued)

Aglycoeremomycin, DMSO, 340 K			Aglycovancomycin, DMSO, 330 K		
Assignment	¹ H (ppm)	¹³ C (ppm)	Assignment	¹ H (ppm)	¹³ C (ppm)
y4	—	169.18	y4	—	169.34
y5	—	169.08	y5	—	169.22
y6	—	167.02	y6	—	167.01
y7	—	167.61	y7	—	167.46
y8 (Asp)	—	172.39	y8 (Asp)	—	172.47
Z2	5.139	71.5	Z2	5.171	71.27
Z6	5.195	71.82	Z6	5.150	71.45

(b) ¹H/¹⁵N chemical shift assignment of the aglycones of eremomycin and vancomycin. ¹⁵N chemical shift scale is derived indirectly from solvent ¹H chemical shifts (2.50 ppm for DMSO) and the known gyromagnetic ratios of ¹H and ¹⁵N nuclei

Aglycoeremomycin, DMSO, 340 K			Aglycovancomycin, DMSO, 330 K		
Assignment	¹ H (ppm)	¹⁵ N (ppm)	Assignment	¹ H (ppm)	¹⁵ N (ppm)
w2	8.560	121.25	w2	(8.335)	(121.48)
w3	6.263	118.60	w3	6.559	117.55
w4	8.105	120.85	w4	8.077	118.50
w5	8.633	123.97	w5	8.494	123.65
w6	6.468	107.13	w6	6.572	106.65
w7	8.556	126.50	w7	8.452	126.99
NH ₂ (Asp)	6.964/7.406	110.84	NH ₂ (Asp)	6.817/7.284	110.14

(c) ¹H/¹³C chemical shift assignment of the aglycones of ristocetin and teicoplanin. Solvent reference signals were used for ¹³C calibration CD₃OD=49.15 DMSO=39.51 ppm

Aglycoristocetin, CD ₃ OD, 300 K			Aglycoteicoplanin, DMSO, 330 K		
Assignment	¹ H (ppm)	¹³ C (ppm)	Assignment	¹ H (ppm)	¹³ C (ppm)
1a	—	125.30	1a	—	131.48
1b	6.774	118.39	1b	6.682	117.45
1c	—	145.04	1c	—	141.47
1d	—	150.81	1d	—	146.83
1e	6.925	119.23	1e	6.936	118.25
1f	7.096	124.31	1f	7.125	124.89
2a	—	137.68	2a	—	135.42
2b	7.908	130.97	2b	7.22	130.90
2c	7.158	129.05	2c	—	126.11
2d	—	157.22	2d	—	150.68
2e	7.118	128.25	2e	7.224	124.55
2f	7.161	124.18	2f	7.646	129.87
3a	—	138.45	3a	—	140.70
3b	6.482	110.03	3b	6.359	110.01
3c	—	157.35	3c	—	158.44
3d	—	115.12	3d	6.340	104.87

Table 2 (c) (continued)

Aglycoristocetin, CD ₃ OD, 300 K			Aglycoteicoplanin, DMSO, 330 K		
Assignment	¹ H (ppm)	¹³ C (ppm)	Assignment	¹ H (ppm)	¹³ C (ppm)
3e	—	158.79	3e	—	157.10
3f	6.630	106.1	3f	6.40	102.66
4a	—	129.2	4a	—	127.61
4b	5.785	107.86	4b	5.568	107.18
4c	—	150.89	4c	—	147.61
4d	—	135.95	4d	—	134.19
4e	—	150.69	4e	—	147.50
4f	5.378	106.87	4f	5.147	104.30
5a	—	127.44	5a	—	125.83
5b	7.06	137.58	5b	7.123	135.53
5c	—	121.92	5c	—	121.13
5d	—	156.94	5d	—	155.40
5e	6.822	121.42	5e	6.670	116.52
5f	6.851	128.21	5f	6.72	125.66
6a	—	139.94	6a	—	142.04
6b	7.427	127.65	6b	7.800	127.16
6c	6.882	122.68	6c	—	127.51
6d	—	156.8	6d	—	148.63
6e	6.796	123.45	6e	7.234	123.08
6f	7.479	129.45	6f	7.466	127.22
7a	—	137.02	7a	—	136.14
7b	—	118.83	7b	—	117.69
7c	—	157.76	7c	—	156.35
7d	6.515	104.36	7d	6.40	102.66
7e	—	159.22	7e	—	157.30
7f	6.277	107.91	7f	6.292	105.94
X1	5.409	57.6	X1	4.587	57.92
X2	5.401	62.87	X2	5.02	54.27
X3	5.459	60.2	X3	5.358	57.94
X4	5.778	56.09	X4	5.631	54.84
X5	4.528	55.49	X5	4.384	53.53
X6	4.147	64.63	X6	4.14	61.63
X7	4.790	58.34	X7	4.459	56.63
y1	—	171.81	y1	—	173.21
y2	—	171.26	y2	—	169.06
y3	—	172.18	y3	—	168.01
y4	—	171.08	y4	—	170.05
y5	—	171.81	y5	—	169.06
y6	—	170.64	y6	—	167.35
y7	—	173.77	y7	—	172.25
Z2	5.076	73.29	Z2	2.869/3.321	36.69
Z6	5.376	73.37	Z6	5.117	71.40
Me3	1.960	8.45			
Me7 (OMe)	3.831	52.99			

Table 2 (continued)

(d) $^1\text{H}/^{15}\text{N}$ chemical shift assignment of the aglycone of ristocetin and teicoplanin. ^{15}N chemical shift scale is derived indirectly from solvent ^1H chemical shifts (3.31 for CD_3OD and 2.50 ppm for DMSO) and the known gyromagnetic ratios of ^1H and ^{15}N nuclei

Aglycoristocetin, CD_3OD , 300 K			Aglycoteicoplanin, DMSO, 330 K		
Assignment	^1H (ppm)	^{15}N (ppm)	Assignment	^1H (ppm)	^{15}N (ppm)
w2	10.297	112.80	w2	7.395	116.06
(w3)	9.305	118.17	w3	7.627	118.75
(w4)	9.894	126.10	w4	7.529	115.98
w5	8.758	124.88	w5	8.301	124.17
w6	7.135	108.47	w6	6.571	106.66
w7	8.948	127.87	w7	8.316	126.40

inhomogeneous crude product on a Silicagel 60 column (see Experimental) furnished aglyco-ristocetin A (**6**) which was homogeneous as shown by TLC, HPLC and MALDI-TOF mass spectrometry (Table 1), as well as by means of ^1H and ^{13}C NMR assignments (Table 2c~d).

Characterization of **6** was very important in the knowledge of the results reported by Ellestad *et al.* [6], who observed epimerization of the methin-group of the *N*-terminal-4-hydroxyphenylglycine unit of avoparcins. In addition, Herrin *et al.* [7] performed inversion of the configuration at the *N*-terminal amino group of aglyco-ristocetin A (ring of ristomycinic acid) by a chemical method to obtain *epi*-aglyco-ristocetin A (**7**). Thus the *R* configuration of the named asymmetric center in **6** changed to *S* in **7**.

Preparation of Semisynthetic Glycopeptide Antibiotics

The goal of the development of new series of the glycopeptide antibiotics is to produce analogues active against methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus faecalis* (VREF) strains.

In a recent work we have successfully applied squaric acid esters, introduced by Tietze *et al.* [8], for the chemical modification of anthracycline glycoside antibiotics [9, 10]. It is known that squaric esters (**7**) selectively react with a small excess of primary or secondary amines in aqueous and/or alcoholic media even in the presence of alcoholic and phenolic hydroxyl, and also carboxyl groups. In neutral media the product is always the squaric acid amide ester (**8**), whereas under more alkaline conditions (pH >7) the corresponding symmetric (**9**) or asymmetric diamide (**10**) are produced (Scheme 3).

The reaction of the amino functions of the glycopeptide antibiotics located at *N*-terminus of the heptapeptide aglycones and at the aminodeoxy sugars with a squaric acid ester


would afford mono and diamide or triamide derivatives. The diamides may possess symmetric or asymmetric structure. The primary and secondary *N*-terminal amino groups of the aglycones are always more reactive than the amino function of the aminodeoxy sugar moieties.

The reaction of the antibiotics **1**, **2** and **5** with the squaric acid ester **7**, furnished (Scheme 4) the squaric acid amide esters **11**, **12** and **13**, which spontaneously separated from the reaction mixtures. Due to the slight water-solubility of **6**, analogous reaction gave the **14** amide ester of aglyco-ristocetin A only in a 1:1 methanol-buffer (pH ~7) mixture. The best yields were obtained with **5** and **6**, carrying an *N*-terminal primary amino group, to give the squaric acid amide esters **13** and **14** of ristomycin A and aglyco-ristomycin A, respectively.

The homogeneity/purity of the above squaric acid amide esters was investigated by TLC and HPLC, and their structures were substantiated by means of MALDI-TOF mass spectrometry, the UV spectral data ($\lambda_{\text{max}}=277\sim 281\text{ nm}$), characteristic of the glycopeptide antibiotics (Table 3), and by the ^1H and ^{13}C NMR spectra (Table 5).

Theoretically, it cannot be excluded that besides hydrogen-bonding, the antibiotic squaric acid amide esters **11**~**14** can bind to the peptidoglycan fragment of the bacterial cell wall also with covalent bonding at an appropriate pH.

Of the numerous, known semisynthetic derivatives [2, 11~16] the most active are those in which the amino group of the heterodisaccharide side-chain is substituted either with a bulky aromatic group, or a longer alkyl chain. We decided to enhance the lipophilic character of the glycopeptide molecules by reacting compounds **11**~**14** with 4-phenylbenzylamine (**15**) to furnish the asymmetric diamides **16**~**19** (Scheme 4). The characteristic physico-chemical and NMR data are summarized in Tables 4 and 5,



Scheme 2 Deglycosidation of ristocetin A.


respectively.

Scheme 5 shows that the reaction of the *N*-terminal amino group of the aglycone of the glycopeptide antibiotics with dimethyl squarate results first in the reactive squaric acid amide esters, whose treatment with the selected amino compound leads to the target amides. However, such asymmetric molecules can also be constructed (Scheme 5) by the preparation of the reactive vinylamide-type squaric acid half-ester **21** from dimethyl squarate (**7**) and triglycine

(**20**), followed by reaction with the *N*-terminal amino group of **5** or its aglycone **6** under basic conditions (pH >8) to obtain the target molecules **22** and **23**. The advantage of both methods is that because of the high regioselectivity no protecting group strategy is needed to perform the experiments.

Mass Spectrometry

In the reaction of the antibiotics with dimethyl squarate we



Scheme 3 The squarate linker method [8].

Table 3 Physico-chemical data of the squaric acid monoamides of glycopeptide antibiotics

Squaric acid monoamides of	Yield %	HPLC R_t	TLC Rf	Molecular weight			$[\alpha]_D^{20}$	UV λ_{max} nm
				Formula of compounds	Calculated	Measured MALDI-TOF (M+Na) ⁺		
11 Eremomycin	53.0	1.72	(1B) 0.31	$\text{C}_{78}\text{H}_{91}\text{N}_{10}\text{O}_{29}\text{Cl}$	1668.04	1689.66	-48.42 ($c=0.28$, DMSO)	273 (DMSO)
12 Vancomycin	35.05	8.05	(1A) 0.26 (2C) 0.64	$\text{C}_{71}\text{H}_{77}\text{N}_9\text{O}_{27}\text{Cl}_2$	1559.29	1580.58	32.08 ($c=0.21$, DMF)	281 (DMF)
13 Ristocetin-A	82.8	4.32	(1A) 0.43	$\text{C}_{100}\text{H}_{112}\text{N}_8\text{O}_{47}$	2177.0	2199.33	-161.94 ($c=0.20$, DMF)	277 (DMF)
14 Ristocetin-aglycone	60.10	36.51	(2D) 0.36	$\text{C}_{65}\text{H}_{53}\text{N}_7\text{O}_{22}$	1284.13	1306.50	-126.99 ($c=0.20$, MeOH)	277 (MeOH)


TLC: (1) DC Cellulose
(2) Kieselgel 60F₂₅₄
Detection: Pauly-reagent

Solvent system: (A) *n*BuOH - Pyr - AcOH - H₂O (15 : 10 : 3 : 12)
(B) *n*BuOH - AcOH - H₂O (4 : 1 : 1)
(C) *n*PrOH - NH₄OH (1 : 1)
(D) Toluene - MeOH - AcOH (1 : 1 : 0.05)

observed the formation of a single product, the site of the attachment was unknown. Theoretically, from eremomycin three types of monosquaramide can be obtained, from vancomycin and ristocetin A two possible regioisomers can be deduced. We solved this problem using electrospray mass spectrometry.

A preliminary experiment was done on the protonated molecular ion of **19** to see the tandem mass spectrometric behavior of the squaric acid-modified aglycone. The loss of the squaric acid moiety was found to be the most characteristic process. In the eremomycin derivative **16**

there are three possible places to which the squaric acid may be bond: the two amino sugar moieties and the *N*-terminus of the peptide core. The molecular weight of **16** (1819.2) is out of the mass range of the instrument so the doubly charged ion at $m/z=910.3$ was selected as precursor in the EPI experiment. Singly and doubly charged ions are also appeared in the spectrum. The most intense peak in the spectrum at $m/z=1676.3$ is formed by loss of 144 Da. The mass of both R₂ and R₃ (for abbreviations see Scheme 1 and 2) is 144 but the consecutive loss of 305 Da resulting a peak at $m/z=1371.1$ clearly shows that the former is due to




Scheme 4 Preparation of the squaric acid amide esters and asymmetric diamides.

the loss of R₃ and the latter is that of the disaccharide chain containing R₂, respectively. Peaks in the lower *m/z* region at 144 and 306 are correspondent to the R₁ and to the disaccharide unit with R₂, respectively. No peaks could be found in the spectrum corresponding to the combination of either of the two aminosugars and the squaric acid (Figure 2). From these data it can be concluded that the squaric acid was bonded to the *N*-terminus of the peptide chain.

There are only two possible positions for the reaction with squaric acid in vancomycin, *i.e.* the R₂ and the *N*-terminus. The EPI experiment with derivative **17** was done on the doubly charged molecule (856.3). The spectrum obtained is quite simple, only several peaks appeared.

The most intense peak at *m/z*=1406.2 is attributed to corresponding to the loss of the disaccharide unit with R₂. This is only possible if the squaric acid is linked at the *N*-terminus in the molecule (Figure 3).

The EPI spectrum of the doubly charged ristocetin derivative **18** seemed the most complicated among the spectra of antibiotics examined, and this is due to the number and complexity of the carbohydrate side chains. There are two possible positions of the squaric acid: at the aminosugar (R₂) and at the X₂. The combination of the consecutive losses of R₁ R₂ R₃ and 167 Da (biphenylmethyl group from the squaric acid part) can be found in the spectrum (Figure 4). There are two peaks in the spectrum at



Scheme 5 Synthesis of the squaric acid asymmetric diamides of triglycine and ristocetin-A or aglyco-ristocetin-A.




Fig. 2 Enhanced product ion (EPI) spectrum of the doubly charged eremomycin derivative (**16**).

SA represents the squaric acid moiety.

$m/z=1598.2$ and 1436.2 , corresponding to the loss of R_1+R_2 and $R_1+R_2+R_3$, respectively, containing the squaric acid moiety but not an aminosugar portion. The latter has the same mass as the protonated molecular ion of the

aglyco-ristocetin A derivative **19**. These data support our assumption that the squaric acid is connected to the antibiotic at position X_2 .




Fig. 3 Enhanced product ion (EPI) spectrum of the doubly charged vancomycin derivative (**17**).




Fig. 4 Enhanced product ion (EPI) spectrum of the doubly charged ristocetin-A derivative (**18**).

SA represents the squaric acid moiety.

Concerted ^1H , ^{13}C and ^{15}N NMR Assignment of the Four Basic Glycopeptide Aglycones and Application in the Structure Elucidation of New Derivatives

Structures of antibiotics are often supported by detailed ^1H -

^{13}C NMR analysis, however ^{15}N data are sparse (missing for aglycones). It has been demonstrated that ^{15}N chemical shifts are extremely useful for locating H-bonds [17, 18] while ^{15}N relaxation gives more insight into dynamics [19].




Fig. 5 Comparison of the $^{15}\text{N}/^1\text{H}$ HSQC spectra of the four aglycones.

Pioneer NMR assignments are difficult to compare, since the experimental conditions and methods are very different. In aqueous solution, many glycopeptide antibiotics form asymmetric dimers, and therefore experiments were often carried out in DMSO solution, at high temperature to get rid of exchange broadened peaks. Certain derivatives, including the aglycones, are poorly soluble in water. Early aglycone assignments were mostly based on chemical shift comparison and homonuclear correlations; *e.g.* for teicoplanin [20] and vancomycin [21, 22], eremomycin [23].

The assignment strategy for the aglycones was similar to our earlier methods, however in this work the starting points were the amides that were identified from the ^{15}N HSQC-spectra (Figure 5). The main chain ^1H assignment of the heptapeptide core was augmented with homonuclear NOESY in addition to COSY and TOCSY spectra. Sequential NH-NH NOE-s can be observed at the *N*-terminus up to residue 4, because of the β -sheet like secondary structure [24]. Due to limited homonuclear connectivities ^{13}C assignments were based on ^{13}C - ^1H HSQC and HMBC spectra for detecting one-bond and long-range heteronuclear connectivities. The latter was especially

useful for the assignments of the aromatic rings. The NMR assignments of the four aglycones are grouped in tables (Table 2a, 2b Table 2c, 2d) containing the vancomycin-eremomycin and ristocetin-teicoplanin pairs. It can be seen that the independent (self-consistent) full assignments of the pairs are well comparable, lending further support to their reliability. Furthermore, comparison with earlier assignments of the intact eremomycin and its aglycone [23] supports that the aglycone does not form dimers in DMSO solution, in contrast to eremomycin in aqueous solution. This is proved by the ^1H chemical shift of the 6e proton ($\delta=6.89$ ppm) in the monomeric aglycone, in contrast to 5.3 ppm in the dimer, where the unusual aromatic shift is due to orthogonal σ - π interaction between H-6e and ring 4. ACD chemical-shift prediction software [25] proved to be useful: though it was insufficient for predicting small chemical shift differences. Because of better solubility and resolution we gave the assignments of the ristocetin aglycone in methanol solution.

Squaric Acid Derivatives of the Glycopeptide Aglycones

In Table 5 we present only the assignments of the

substituents attached the glycopeptide aglycones. The presence of the aglycones was proved by overlaying their 2D-HSQC spectra with the appropriate aglycone, although it is occasionally complicated with signal-doubling due to the presence of the amide group restricted rotation in the squaric acid [8]. Therefore, some assignments are missing (see as ND label) due to low signal—to noise ratio

in their ^{13}C NMR spectra. However, in such cases, mass spectra were convincing about the identity of the semisynthetic derivatives and the site of the squaric acid attachment.

Antibacterial Activity

The aglycones are responsible for the biological activity of


Table 4 Squaric acid asymmetric diamides of glycopeptide antibiotics

Asymmetric diamides of	Yield %	HPLC Rt	TLC Rf	Molecular weight			$[\alpha]_D^{20}$	UV λ_{max} nm
				Formula of compounds	Calculated	Measured MALDI-TOF (M+Na) ⁺		
16 Eremomycin	27.6	26.26	(1A) 0.60	$\text{C}_{90}\text{H}_{100}\text{N}_{11}\text{O}_{28}\text{Cl}$	1819.24	1841.81	-46.7 (<i>c</i> =0.24, DMSO)	287 (DMSO)
17 Vancomycin	29.3	20.22	(1A) 0.58	$\text{C}_{83}\text{H}_{86}\text{N}_{10}\text{O}_{26}\text{Cl}_2$	1710	1732.27	20.5 (<i>c</i> =0.265, DMF)	291 (DMF)
18 Ristocetin-A	27.8	17.59	(1A) 0.60 (3C) 0.40	$\text{C}_{112}\text{H}_{121}\text{N}_9\text{O}_{46}$	2329.16	2351.28	-149 (<i>c</i> =0.2, DMF)	289 (DMF)
19 Aglyco-ristocetin	68.4	18.47	(2B) 0.53	$\text{C}_{77}\text{H}_{62}\text{N}_8\text{O}_{21}$	1435.34	1457.26	-183 (<i>c</i> =0.22, MeOH)	288 (MeOH)
22 Ristocetin-A	73.3	8.75	—	$\text{C}_{105}\text{H}_{119}\text{N}_{11}\text{O}_{50}$	2335.08	2356.80	-127 (<i>c</i> =0.32, DMSO)	289 (DMSO)
23 Aglyco-ristocetin	62.5	10.90	(2A) 0.68	$\text{C}_{70}\text{H}_{60}\text{N}_{10}\text{O}_{25}$	1441.26	1463.51	-157 (<i>c</i> =0.52, DMF)	289 (DMF)


TLC: (1) DC Cellulose
(2) Kieselgel 60F₂₅₄
(3) Reverse phase RP-18F₂₅₄
Detection: Pauly-reagent

Solvent system: (A) *n*BuOH - Pyr - AcOH - H₂O (15 : 10 : 3 : 12)
(B) Toluene - MeOH (1 : 1)
(C) EtOH - AcOH - H₂O (4 : 0.1 : 6)


Table 5 NMR data of squaric acid parts



	11		12		13		14	
	^{13}C (ppm)	^1H (ppm)	^{13}C (ppm)	^1H (ppm)	^{13}C (ppm)	^1H (ppm)	^{13}C (ppm)	^1H (ppm)
a1	175.5	—	(167.69)	—	167.71	—	ND	—
a2	170.83	—	(169.94)	—	169.00	—	ND	—
a3	189.22	—	190.12	—	183.25	—	181.1	—
a4	197.53	—	182.44	—	183.35	—	190.26	—
a5	59.94	4.30	61.59	4.34	60.94	4.29	60.97	4.35

Table 5 (continued)

	17		18		19	
	¹³ C (ppm)	¹ H (ppm)	¹³ C (ppm)	¹ H (ppm)	¹³ C (ppm)	¹ H (ppm)
a1	167.82	—	167.19	—	169.13	—
a2	168.66	—	168.8	—	168.91	—
a3	183.39	—	183.66	—	183.53	—
a4	182.94	—	ND	—	183.27	—
b1	140.05	—	140.4	—	140.26	—
b2. b6	127.70	7.65	127.7	7.65	127.50	7.65
b3. b5	129.13	7.51	129.6	7.45	129.70	7.45
b4	139.22	ND	138.72	—	139.47	—
b7	140.70	ND	140.77	—	140.68	—
b8. b12	127.70	7.65	127.7	7.65	127.40	7.61
b9. b11	129.13	7.41	129.00	7.41	129.00	7.40
b10	128.29	7.37	128.40	7.36	128.20	7.35
b13	47.21	4.83	47.54	4.77	47.37	4.77



	22		23	
	¹³ C (ppm)	¹ H (ppm)	¹³ C (ppm)	¹ H (ppm)
a1	168.94	—	167.35	—
a2	ND	—	168.50	—
a3	183.86	—	184.01	—
a4	ND	—	183.70	—
c1	43.53	3.54	43.83	3.44
c2	43.18	3.77	43.29	3.76
c3	46.82	4.28	46.71	4.29

ND are not detectable signals because of low signal-to noise ratio. The characteristic ¹³C and ¹H NMR signals of the glycopeptide antibiotic parts of the molecules are not listed, though they were unambiguously detected in the overlaid 2D HSQC spectra according to the assigned spectra of the aglycones.

Table 6 Antibacterial activity of the glycopeptide antibiotics and their derivatives

Test microorganism	MIC ($\mu\text{g/ml}$)																Linezolid [30]
	Eremomycin				Vancomycin				Ristocetin								
	1	3	11	16	2	4	12	17	5	6	13	14	18	19	22	23	
<i>Bacillus subtilis</i> ACTC 6633	<0.5	32	1	1	<0.5	1	2	0.5	<0.5	2	8	16	0.5	2	8	8	1.0
<i>Staphylococcus aureus</i> ACTC 29213 (MSSA)	0.5	32	2	8	2	4	4	1	1	2	64	8	4	1	128	4	2.0
<i>S. aureus</i> ACTC 33591 (MRSA)	0.5	32	4	16	2	4	8	2	4	2	128	8	16	8	128	4	1.5
<i>S. epidermidis</i> ACTC 25299 (MSSE)	0.5	8	2	2	2	4	2	2	2	2	64	4	32	0.5	32	8	1.0
<i>S. epidermidis</i> IMMSU 12333/05 (MRSE)	2	128	8	64	4	4	4	8	16	4	128	32	32	8	128	32	1.0
<i>Enterococcus faecalis</i> ACTC 29212 (VSEF)	1	64	8	4	2	4	16	1	2	4	32	16	1	1	1	4	4.0
<i>E. faecalis</i> ACTC 51299 (GREF)	8	>256	64	16	32	8	32	16	8	16	128	64	2	8	4	4	3.0

ACTC: American Collection of Typed Culture, MSSA: Methicillin sensitive *S. aureus*, MRSA: Methicillin resistant *S. aureus*, VSEF: Vancomycin (and teicoplanin) sensitive *E. faecalis*, GREF: Glycopeptide resistant *E. faecalis*, MRSE: Methicillin resistant *S. epidermidis*, MSSE: Methicillin sensitive *S. epidermidis*.

the glycopeptide antibiotics. The antimicrobial activities of the aglycones, obtained after removal of the carbohydrate moieties, in comparison with those of the parent antibiotics are summarized in Table 6.

Deglycosylation of eremomycin (**1**) to **3** results in a surprising decrease in the activity against all of the tested seven microorganisms. The activity of aglyco-vancomycin (**4**) against *Bacillus subtilis* decreased to more than the half of the original value, and *ca.* to the half against methicillin-sensitive and resistant *Staphylococcus aureus* (MSSA, MRSA) and *S. epidermidis* (MRSE, MSSE).

According to Boger [26], the elemental composition and structural features of the aglycones **3** and **4** are very similar, and—besides the atropisomerism (rings 5 and 7) of actinoidinic acid—the chloro substituent of the aromatic rings of **2** and **6** has a complementary role in the affinity and selectivity of binding to the L-Lys-D-Ala-D-Ala tripeptide of the bacterial cell wall. By comparing the antibiotic activity of **1** and **3** and **6**, Russian researchers came to the same conclusion [14, 15].

In the case of ristomycin A (**5**), the aglyco-ristomycin (**6**), which does not carry a covalent chloro atom, a positive change in the activity against both MSSA and MRSA was observed. This aglycone (**6**) was found twice as active against MRSA as the parent antibiotic (**5**). As compared to **5**, the activity of **6** against both *Enterococcus faecalis* strains dropped to half, and to one-fourth against *B. subtilis*. The MIC values for **6** and **2** are approximately the same obtained with the organisms *S. aureus*, *S. epidermidis* and *E. faecalis*. Moreover, these efficacies compare favourably [30] to those of the new synthetic oxazolidinone derivative, linezolid which is in clinical use.

The antibacterial activity of the of the squaric acid amide esters **11** and **13**, derived from eremomycin and ristocetin A, against the test microorganisms is much decreased as compared to both of the parent antibiotics **1** and **5**, and to the corresponding aglycones. At the same time, the squaric acid amide ester of aglyco-ristocetin (**14**) possesses lower decrease in activity than compound **13**.

Of the asymmetric squaric amides **18** and **19**, derived from ristocetin A (**5**) and aglyco-ristocetin A (**6**), respectively, with 4-phenylbenzylamine (**15**), the activity of **18** significantly increased against *B. subtilis* and *E. faecalis* (VSEF, GREF).

Except for MRSA, the effectiveness of **19** is higher than that of **14** towards each of the investigated bacterial strains, and practically identical with that of the starting aglycone **6**.

Experimental

The samples of eremomycin, ristocetin A and aglyco-teicoplanin were received from the Gauze Institute of New Antibiotics (Moscow). Vancomycin hydrochloride was purchased from Aldrich Co.

The solutions were evaporated under diminished pressure at 40°C. Preparative column chromatography was performed on Silica gel 60 (0.063~0.2 mm, Merck) and Silica gel 60 silanized (0.063~0.2 mm, Merck) adsorbents. HPLC was carried out with a Waters 600 instrument using UV detection at 230 nm. Column: Lichrospher RP-8 (4×250 mm 10 μm); injection volume: 20 μl (from 0.1 mg/ml). Eluent A: CF₃COOH–H₂O (pH=2.65); B:

MeCN-water, gradient elution from 10% to 90%. The UV spectra were recorded with a Perkin-Elmer Lambda 11 instrument. The specific optical rotation values were measured with a Perkin-Elmer 341 automatic polarimeter. MALDI-TOF MS spectra were obtained with a Bruker BIFLE X III spectrometer. Positive ions were detected in the refletron mode. Samples were prepared with DBH matrix dissolved in DMSO (20 mg/ml). Concentrations of compounds: 5 mg/ml. An Applied Biosystems 3200 Qtrap hybrid linear ion trap mass spectrometer equipped with TurboV ion source was used for the identification of the position of the squaric acid moiety. The samples were dissolved in acetonitrile : water : formic acid 50 : 50 : 0.1% solution and were infused using a built-in syringe pump with a flow rate of 10 μ l/minute. The ionization potential was 5000 V. The instrument was scanned in linear ion trap (LIT) mode using a scan range of 500~1700 and 50~1700 in enhanced MS (EMS) and enhanced product ion (EPI) experiments, respectively. The Q0 trapping function was turned on. NMR samples were measured in DMSO- d_6 solution at 330 or 340 K, except the aglyco-ristocetin (MeOH- d_4 or MeOH/MeOH- d_4 9:1 mixture for ^{15}N , 300 K). Chemical shifts are referenced to published residual solvent signals (39.51 for DMSO- d_6 and 49.15 MeOH- d_4 for ^{13}C or TMS=0 for ^1H NMR). A Bruker DRX-500 spectrometer equipped with a 5 mm inverse broadband gradient probe head (with 10, 13 and 23 μ s duration of 90° pulses for ^1H , ^{13}C and ^{15}N channels) was used. Simple, gradient-enhanced magnitude representation COSY-60 spectra were run with solvent presaturation when necessary. Gradient and sensitivity enhanced TOCSY spectra were acquired with DIPSI2 mixing sequence of 60 ms. ^{13}C - ^1H and ^{15}N - ^1H HSQC spectra were obtained with gradient and sensitivity enhancement (^{15}N spectra required overnight accumulation at natural abundance). The used NOESY pulse sequence was equipped with the Hahn spin-echo to remove broad solvent lines and baseline bias. The same spin-echo technique was applied in 1D ^1H signal accumulation. In case of derivatives, straight ^{13}C accumulation was sometimes a bottleneck, because of the low ^{13}C sensitivity of the inverse probe. Therefore, in such cases, the basic acquisition scheme of 30° ^{13}C read pulse was applied, instead of the more informative J-modulated spin-echo sequence. The gradient enhanced HMBC sequence was preceded with a filter to suppress one-bond correlations and ^{13}C decoupling was applied during acquisition. SPARKY software package [31] was used for visualization and assignment.

For the deglycosidation experiments a Toho Kasei (Japan) apparatus at the Department of Organic Chemistry, Lóránd Eötvös University of Budapest was used.

Deglycosidation of Eremomycin and Vancomycin

Deglycosidation of eremomycin and vancomycin was carried out as described earlier by Wanner *et al.* [5] (25°C, 2 hours). The physico-chemical data of the prepared aglycones **3** and **4** are collected in Table 1, and the NMR data are presented in Tables 2a~b.

Preparation of Aglyco-ristocetin A (6)

In the teflon tube of the deglycosidation instrument anisole (1.9 ml) was added to ristocetin A sulfate (5200 mg), the mixture was cooled to -70 to -80°C and 50 ml of HF was added. The excess of HF was then removed through a CaO trap in vacuum at -18°C. The crude reaction product was washed three times with an ice-cold 4:1 ethyl acetate-methanol mixture, filtered and dried in a vacuum desiccator. The product contained three UV-detectable components on TLC.

Column Chromatography

The crude aglycone was dissolved in methanol, containing a small amount of acetic acid (6 ml methanol+2 drops of AcOH), and it was applied onto the surface of Silicagel 60 (3.0 g) by evaporation. The residue was applied to the top of a Silicagel 60 (0.063~0.2 mm) column (50×2 cm), and gradient elution was performed with toluene-MeOH-AcOH (7:3:0.1)→toluene-MeOH-AcOH (6:4:0.1). The fractions with Rf: 0.26 was collected, evaporated and dried in a vacuum desiccator to furnish 1.50 g (52.8%) of **6**, whose physical and NMR data are shown in Tables 1 and 2c~d, respectively.

Synthesis of the Squaric Acid Amide Esters **11~14** of the Glycopeptide Antibiotics and Their Aglycones

To a solution of the antibiotic/aglycone **1~6** (0.1 mmol) in Tristisol buffer (5.0 ml, pH=7.3) dimethyl squarate (**7**, 0.1 mmol) was added at room temperature, and the mixture was stirred at the same temperature overnight. The resulting precipitate was filtered off, and washed three times with distilled water (3.0 ml), methanol and ether (3.0~3.0 ml). Column chromatographic purification (6:4:0.025 toluene-MeOH-AcOH) was necessary only in the case of **14**. The yields and the physico-chemical data for the products **11~14** are shown in Table 3, and the NMR characteristics are collected in Table 5.

Preparation of the Asymmetric Squaric Acid Amides (**16~19**, **22**, **23**) of the Glycopeptide Antibiotics and Their Aglycones

Method A

The squaric acid amide esters **11~14** (0.1 mmol) were

dissolved in a mixture (pH >8) of water (4.0 ml), methanol (4.0 ml) and triethyl amine (0.01 ml), and then 4-phenylbenzylamine (**15**, 0.1 mmol) was added. The reaction mixture was stirred overnight, and concentrated under diminished pressure. The residue was washed with methanol and water (3.0~3.0 ml) and dried in a vacuum desiccator over CaCl₂. Compounds **18** and **19** were purified by means of column chromatography. The amide **18** was chromatographed on silanized Silicagel 60 with gradient elution (8:2:0.1→7:3:0.1→6:4:0.15 H₂O - EtOH - AcOH). Pure **19** was eluted from a Silicagel 60 column with 1:1 toluene - methanol. The yields and physico-chemical characteristics and NMR data of the prepared **16**~**19** are presented in Tables 4 and 5, respectively.

Method B. The Reverse Method

Dimethyl squarate (**7**, 0.5 mmol) and triglycine (**20**, 0.5 mmol) were dissolved in a mixture of *N,N*-dimethyl formamide (6.0 ml) and Tristisol buffer (2.0 ml, pH >7), and the mixture was stirred at room temperature for 28 hours. Evaporation of the reaction mixture under diminished pressure afforded the squaric acid triglycine amide **21**, which was purified by column chromatography (eluent: 4:1:1 *n*BuOH - AcOH - H₂O, R_f=0.46). Compound **5** or **6** (0.5 mmol) was dissolved in a mixture of Tristisol buffer (20 ml) and MeOH (30 ml), then **21** (0.5 mmol) was added and the pH was adjusted to 8.0. The reaction mixture was stirred at room temperature for 48 hours, the precipitated product was filtered off, washed with water and 50% aqueous methanol and dried in a vacuum desiccator. The physico-chemical data and NMR spectral characteristics of **22** and **23** are given in Tables 4 and 5, respectively.

Antibacterial Activity Testing

Bacterial Strains

The antibacterial activities of the preparations and their derivatives were measured on *B. subtilis* ACTC 6633, methicillin-sensitive *S. aureus* ACTC 29213 (MSSA), methicillin-resistant *S. aureus* ACTC 33591 (MRSA), methicillin-resistant *S. epidermidis* IMMSU12333/05 (MRSE), vancomycin- and teicoplanin-sensitive *E. faecalis* ACTC 29212 (VSEF), and vancomycin-resistant but teicoplanin-sensitive *E. faecalis* ACTC 51299 (VREF) strains. The confirmation of the presence and absence of the *vanA*~*B* and *mecA* genes was done as previously described [27, 28].

Susceptibility Test

In all cases, except for linezolid, the efficacy of the antimicrobials was determined with the broth micro

dilution method according to the NCCLS guideline [29]. Bacterial strains were grown on 5% bovine blood agar plates at 35°C overnight. Appropriate numbers of colonies were suspended in physiological saline in order to reach the density of 0.5 McFarland for inoculation.

Stock solutions of different concentrations of the antimicrobials were prepared in either distilled water or water - methanol (1:1) or in water - DMSO (1:1) depending on the solubility of the given preparation. These were two-fold serially diluted from 256 to 0.5 µg/ml in cation-adjusted Mueller-Hinton broth then 100 µl of each dilution was transferred into microplate's holes. Inoculation occurred with 10 µl of each of the bacterial suspensions. Incubation was performed at 35°C for 18 hours and reading of minimal inhibitory concentration (MIC) was made with the naked eyes on a mirror.

Solvents were also tested for inhibition of bacterial growth. None of them exerted bacteriostatic effect at the concentration used.

MICs for linezolid were determined with the E-test on cation-adjusted Mueller-Hinton agar plates according to the instructions of the manufacturer (AB Biodisk, Solna, Sweden).

Acknowledgement The authors thank the Hungarian Academy of Sciences and the National Scientific Research Found (Grant No.: OTKA TO46744 and TO46186) for financial support, Nikolett Mihala (Research Group for Peptide Chemistry of the Hungarian Academy of Sciences, Budapest) for her help in the deglycosidation experiments, and Dr. Sándor Kéki (Department of Applied Chemistry, University of Debrecen) for recording the mass spectra. The skillful technical assistance of Ms. Natasa Pesti (Institute of Medical Microbiology, Semmelweis University) is highly appreciated. We are indebted to Olga Miroshnikova (Gauze Institute of New Antibiotics, Moscow) for the preparation of aglyco-teicoplanin.

References

1. Sztaricskai F, Bognár R. The chemistry of the vancomycin group of antibiotics in recent developments in the chemistry of natural carbon compounds. Vol. 10: 91–209, Szántay Cs, ed., Akadémiai Kiadó, Budapest (1984)
2. Nicolaou KC, Boddy CN, Bräse S, Winssinger N. Biology and medicine of the glycopeptide antibiotics. *Angew Chem Int Ed* 38: 2096–2152 (1999)
3. Sztaricskai F, Pelyvás FI. Chemistry of carbohydrate components in glycopeptide antibiotics. *Drugs and Pharm.Sci.*, Nagarajan R, ed., Marcel Dekker Inc. New York, 63: 105–193 (1994)
4. Williams DH, Bardsley B. The vancomycin group of

- antibiotics and the fight against resistant bacteria. *Angew Chem Int Ed* 38: 1172–1193 (1999)
5. Wanner J, Tang O, Melomas CC, Crowley BM, Jiang W, Moss J, Boger DL. A new improved method for deglycosidation of glycopeptide antibiotics exemplified with vancomycin, ristocetin and ramoplanin. *Bioorg Med Chem Lett* 13: 1169–1173 (2003)
 6. Ellestad GA, Leese RA, Morton GO, Barbatschi F, Gorew E, McGahren WJ. Avoparcin and epiavoparcin. *J Am Chem Soc* 103: 5522–5524 (1981)
 7. Herrin TR, Thomas AM, Perun IJ, Mao JC, Fesik SN. Preparation of biologically active ristocetin derivatives replacements of the 1'-amino group. *J Med Chem* 28: 1371–1375 (1985)
 8. Tietze LF, Arlt M, Beller M, Glüsenkamp KH, Jäde E, Rajewsky MF. Squaric acid diethyl ester: a new coupling reagent for the formation of drug biopolymer conjugates. Synthesis of squaric acid ester amides and diamides. *Chem Ber* 124: 1215–1221 (1991)
 9. Tevyashova A, Sztaricskai F, Batta Gy, Herczegh P, Jeney A. Formation of squaric acid amides of anthracycline antibiotics. Synthesis and cytotoxic properties. *Bioorg Med Chem Lett* 14: 4783–4789 (2004)
 10. Sztaricskai F, Sum A, Roth E, Pelyvás FI, Sándor Sz, Batta Gy, Herczegh P, Reményi J, Miklán Zs, Hudecz F. A new class of semisynthetic anthracycline glycoside antibiotics incorporating a squaric acid moiety. *J Antibiot* 58: 704–714 (2005)
 11. Cooper RDG, Snyder NJ, Zweifel MJ, Staszak MA, Wilkie SC, Nicas TI, Mullen DL, Butler ThF, Rodriguez M J, Huff BE, Thompspon RC. Reductive alkylation of glycopeptide antibiotics: synthesis and antibacterial activity. *J Antibiot* 49: 575–581 (1996)
 12. Kahne D, Leimkuhler C, Lu W, Walsch Ch. Glycopeptide and lipoglycopeptide antibiotics. *Chem Rev* 105: 425–448 (2005)
 13. Mu YQ, Nodwell M, Pace JL, Shaw J-J, Judice JK. Vancomycin disulfide derivatives as antibacterial agents. *Bioorg Med Chem Lett* 14: 735–738 (2004)
 14. Kobrin MB, Katrukha HS, Fedorova GB. Preparation and properties of derivatives at the amino group of ristocetin aglycone. *J Antibiot* 42: 1441–1442 (1989)
 15. Pavlov AYu, Olsuf'eva EN, Miroshnikova OM, Reznikova MI, Lazhko EI, Malabarba A, Ciabatti R, Preobrazhenskaya MN. Synthesis and antibacterial activity of nonnatural aglycones of glycopeptide antibiotics of the vancomycin group. *Russian J Bioorg Chem* 23: 383–393 (1997)
 16. Printsevskaya A, Solovieva SE, Olsufjeva EN, Mirchink EP, Isakova EB, DeClerq E, Balzarini J, Preobrazhenskaya MN. Structure-activity relationship studies of a series of antiviral and antibacterial aglycon derivatives of the glycopeptide antibiotics vancomycin, eremomycin, and dechloroeremomycin. *J Med Chem* 48: 3885–3890 (2005)
 17. Hawkes GE, Molinari H, Singh S, Lian LY. A N-15 NMR study of intermolecular interactions between vancomycin and Ac-D-Ala-D-Ala. *J Magnetic Resonance* 74: 188–192 (1987)
 18. Molinari H, Pastore A, Lian LY, Hawkes GE, Sales K. Structure of vancomycin and a vancomycin/D-Ala-D-Ala complex in solution. *Biochemistry* 29: 2271–2277 (1990)
 19. Batta Gy, Sztaricskai F, Makarova MO, Gladkikh EG, Pogozheva VV, Berdnikova TF. Backbone dynamics and amide proton exchange at the two side of eremomycin dimer by ¹⁵NMR. *Chem Comm* 501–502 (2001)
 20. Malabarba A, Ferrari P, Gallo GG, Kettenring J, Cavalleri B. Teicoplanin, antibiotics from *Actinoplanes teichomyceticus* Nov. sp. *J Antibiot* 39: 1430–1442 (1986)
 21. Bongini A, Feeney J, Williamson MP, Williams DH. Assignment of the carbon-13 spectrum of vancomycin and its derivatives. *J Chem Soc Perkin Trans 2* 201–206 (1981)
 22. Williamson MP, Williams DH. A carbon-13 nuclear magnetic resonance study of ristocetin A and B and their derivatives. *J Chem Soc Perkin Trans 1* 1483–1491 (1981)
 23. Batta Gy, Sztaricskai F, Kövér KE, Rüdél C, Berdnikova TF. NMR Study of eremomycin and its derivatives. Full ¹H and ¹³C assignment, motial behaviour, dimerization and complexation with Ac-D-Ala-D-Ala. *J Antibiot* 44: 1208–1221 (1991)
 24. Loll PJ, Axelsen PH. The structural biology of molecular recognition by vancomycin. *Annual Review of Biophysics and Biomolecular Structure* 29: 265–289 (2000)
 25. ACD Software; Toronto, ON, Canada, M5H 3V9; www.acdlabs.com
 26. Boger DL. Vancomycin, teicoplanin and ramoplanin: synthetic and mechanistic studies. *Med Research Reviews* 21: 356–381 (2001)
 27. Knausz M, Ghidán Á, Grossato A, Rozgonyi F. Rapid detection of methicillin resistance in teicoplanin-resistant coagulase-negative *Staphylococci* by a penicillin-binding protein 2' latex agglutination method. *J Microbiol Methods* 60: 413–416 (2005)
 28. Ghidán Á, Jeney Cs, Maródi LCs, Csiszár K, Rozgonyi F. PCR detection of the *vanA* gene in a vancomycin-resistant *Enterococcus faecalis* clinical isolate from Hungary. *J Antimicrob Chemother* 46: 325–327 (2000)
 29. National Committee for Clinical Laboratory Standards Performance Standards for Antimicrobial Susceptibility Testing. 12th Informational supplement M100-S12. NCCLS, Wayne, PA. USA (2002)
 30. Ford ChW. *In: Oxazolidinones: a new class of antibiotics. Ed., Moellering R Jr., 8th International Congress on Infections Diseases Industry-Sponsored Symposium, (Pharmacia and Upjohn), Boston, May 17 (1998)*
 31. Goddard TD, Kneller DG. SPARKY 3, University of California, San Francisco