

Lepidepyrone, a New γ -Pyrone Derivative, from *Neolentinus lepideus*, Inhibits Hyaluronidase

Tomoo Hosoe, Hideo Sakai, Makoto Ichikawa, Takeshi Itabashi, Takayuki Ishizaki, Ken-ichi Kawai

Received: March 7, 2007 / Accepted: May 29, 2007

© Japan Antibiotics Research Association

Abstract In the course of screening for hyaluronidase (HAase) inhibitory agents, a new γ -pyrone derivative, lepedeprone, $C_8H_{10}O_5$, was isolated from the cultured mycelium of the mushroom *Neolentinus lepideus* TMC-1102 as a major HAase inhibitory compound (IC_{50} 3.3 mM). The structure of lepedeprone was established on the basis of spectroscopic investigation.

Keywords *Neolentinus lepideus*, γ -pyrone derivative, lepedeprone, hyaluronidase inhibitor

The glycosaminoglycan, hyaluronan (hyaluronic acid; HA), which consists of repeating disaccharide units of (β ,1-4)-D-glucuronic acid and (β ,1-3)-N-acetyl-D-glucosamine, is a major component of the extracellular matrix of animal tissues. It is present in the skin (dermis and epidermis), brain, and central nervous system in significant quantities [1].

Hyaluronidases (HAases) are enzymes that degrade HA. The mammalian HAases are present in many tissues and organs, e.g. liver, kidney, testes, etc. and are considered to be involved in many physiological processes like fertilization, tumor growth, and metastasis. Therefore HAase inhibitors could be useful as drugs, e.g. in the

treatment of arthritis or degenerative disease in combination with antibiotics, and/or in antibacterial therapy of HA lyase producing bacteria, such as *Streptococcus pneumoniae*, a major human Gram-positive pathogen [2].

In our search for new HAase inhibitors from Japanese basidiomycetes, the extract from the cultured mycelium of *Neolentinus lepideus* (Japanese mushroom name: Matsuouji) TMC 1102 showed a potent inhibitory activity against HAase. The separation of the above extract led us to isolate a new γ -pyrone derivative designated lepedeprone (**1**) along with methyl isoferulate (methyl 3-hydroxy-4-methoxycinnamate) [3] and ergosterol. In this paper we have reported the structure determination and the inhibitory activity against HAase of **1**.

The molecular formula of **1** (Fig. 1) was confirmed as $C_8H_{10}O_5$ by high resolution chemical-ionization mass spectrometry (HRCI-MS). The UV absorption maximum at 252 nm of **1** suggested the presence of γ -pyrone class of compounds [4], whereas the IR absorptions maxima at $3400\sim 3200\text{ cm}^{-1}$ and 1660 cm^{-1} revealed the presence of hydroxyl and conjugated carbonyl functions, respectively. The $^1\text{H-NMR}$ spectrum of **1** showed the presence of two olefinic protons at δ 6.41 (br s) and 8.06 (d), which suggested that the γ -pyrone ring was substituted with two aliphatic side chains. The substituent residues were hydroxymethyl group [δ 4.42 (s)] and 1,2-dihydroxyethyl group [δ 3.52 (dd), 3.78 (dd), and 4.77 (ddd)]. The




Fig. 1 Structure of lepedeprone (**1**).

T. Hosoe (Corresponding author), **H. Sakai**, **M. Ichikawa**, **T. Itabashi**, **K. Kawai**: Faculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41, Shinagawa-ku, Tokyo 142-8501, Japan, E-mail: kawai@hoshi.ac.jp

T. Ishizaki: Tamagawa University Research Institute, Tamagawa-Gakuen 6-1-1, Machida, Tokyo 194-8610, Japan




Fig. 2 HMBC correlation of lepidopyrone (**1**).





Fig. 3 Structures of ascorbic acid (**2**) and its 6-*O*-acylated derivatives.

positions of these side chains on the γ -pyrone ring were finally determined by the analysis of the HMBC spectrum as shown in Fig. 2. Therefore the structure of lepidopyrone was confirmed as shown in **1** (Fig. 1). The stereochemistry of **1** could not yet be determined.

The inhibitory effects on HAase were determined according to the literature [5]. The MeOH extract of *N. lepideus* TMC 1102 showed the inhibition of 25.7% at the concentration of 2.0 mg/ml. The inhibitory activity was raised to 75.7% at 2 mg/ml after the above extract was extracted with BuOH. **1** was the main component isolated from BuOH extract by the repeated purification of LPLC and HPLC. The inhibitory activity of **1** for HAase was determined as IC₅₀ 3.3 mM, whereas IC₅₀ value of catechin as a positive compound was determined to 1.7 mM (Table 1). L-Ascorbic acid (vitamin C; **2**) showed weak inhibitory activity (IC₅₀ 6.1 mM) on HAase (HA lyase) from *Streptococcus pneumoniae* [7], but **2** did not show any inhibitory activity on bovine testicular HAase up to 100 mM [2]. The inhibitory activities of 6-*O*-acylascorbic acid (**3**–**5**) on mammalian HAase showed stronger in accordance with the length of the side chain [6-decanoate (**3**): IC₅₀ 1.4 mM, 6-hexaecaenoate (**4**): IC₅₀ 57 μ M, 6-octadecanoate (**5**): IC₅₀ 39 μ M]. It is interesting that **1** showed still weak inhibitory activity on mammalian HAase and 6-*O*-acylascorbic acid (**4** and **5**) showed the strong inhibitory activity in consideration with the structural similarity between **1** and **2**.

Table 1 Inhibition of hyaluronidase (HAase)

| Compound | IC ₅₀ on HAase (mM) |
|--|--------------------------------|
| Lepidopyrone (1) | 3.3 |
| L-Ascorbic acid (2) | >100 ^[2] |
| L-Ascorbic acid 6-decanoate (3) | 1.4 ^[7] |
| L-Ascorbic acid 6-hexaecaenoate (4) | 0.057 ^[2] |
| L-Ascorbic acid 6-octadecanoate (5) | 0.039 ^[7] |
| Positive Compound (Catechin) | 1.7 |

Experimental

General

Melting points were determined on a Yanagimoto micro-melting point apparatus and are uncorrected. CI-MS were taken with a JEOL JMS-MS600W spectrometer. UV and IR spectra were recorded on a Hitachi U-3210 spectrometer and a JASCO IR-810 spectrometer, respectively. ¹H- and ¹³C-NMR spectra were recorded on a JEOL Lambda-500 (¹H, 500 MHz; ¹³C, 125 MHz) spectrometer, using tetramethylsilane as an internal standard. CD curves were determined on a JASCO J-600 spectropolarimeter. Column chromatography was performed using Kieselgel 60 (Art. 7734, Merck) and Wakogel C-200 (Art. 237-00071, Wako). Low-pressure liquid chromatography (LPLC) was performed with a Chemco Low-Prep 81-M-2 pump and glass column (200×10 mm) packed with Silica gel CQ-3 (30~50 μ m, Wako). HPLC was performed with a Senshu SSC-3160 pump (flow rate, 7 ml/minute) and a YMC-Pack PEGASIL Silica 60-5 (300×10 mm), equipped with a Shimamura YRD-883 RI detector. TLC were detected by UV light on 254 nm and/or by spraying with 5.0% phosphomolybdic acid - ceric acid (trace) in 5.0% H₂SO₄ and then heating.

Origin of Organisms

N. lepideus retained as M-2721 by the Mushroom Research Institute of Japan, was found on a tree stump (*Pinus densiflora*) in hardwood forest in Gunma Prefecture, east Japan. The strain TMC-1102, which was isolated as a pure culture from the tissue of *N. lepideus* M-2721, was used in this experiment.

Fermentation, Extraction and Isolation

N. lepideus TMC 1102 was cultivated on rice (570 g using 4 flasks) at 25°C for four weeks. The cultivated rice was extracted with MeOH and the organic layer was evaporated *in vacuo*. The residue (42 g) was suspended in water and extracted with BuOH. The organic layer was evaporated *in*

vacuo to obtain the BuOH extract (4.8 g). The extract was purified by LPLC on silica gel column using CH₂Cl₂-MeOH (5:1) to give 5 fractions. The second fraction (1.5 g) was purified by LPLC on silica gel using CH₂Cl₂-acetone (60:1) followed by further purification by HPLC on silica gel. [CH₂Cl₂-Me₂CO (60:1) and/or benzene-EtOAc (7:1)] to give methyl isoferulate (21 mg) and ergosterol (95 mg). The third fraction (0.44 g) was further purified by HPLC on silica gel using CH₂Cl₂-MeOH (60:1) to obtain **1** (217 mg).

Physico-chemical Properties of **1**

Colorless amorphous solid. CI-MS *m/z* (%): 187.0611 [(M+H)⁺, 187.0606 for C₈H₁₁O₃]. UV λ_{max}^{MeOH} nm (log ε): 213 (3.77), 252 (3.94). IR ν_{max} (film) cm⁻¹: 3400~3200 (OH), 2890, 1660 (conjugated C=O), 1590, 1440, 1050, 940, 870. CD (MeOH) Δε (nm): -3.71 (251), +3.26 (293). [α]_D²⁵ -9.5° (c 0.1, MeOH). ¹H-NMR (CD₃OD) δ 3.52 (1H, dd, *J*=11.5, 5.8 Hz, 9-H), 3.78 (1H, dd, *J*=11.5, 3.4 Hz, 9-H), 4.42 (2H, s, 7-H), 4.77 (1H, ddd, *J*=5.8, 3.4, 1.2 Hz), 6.41 (1H, br s, 3-H), 8.06 (1H, d, *J*=1.2 Hz, 6-H). ¹³C-NMR (CD₃OD) δ 61.0 (C-7), 65.8 (C-9), 68.4 (C-8), 112.4 (C-3), 129.6 (C-5), 155.7 (C-6), 171.2 (C-2), 180.7 (C-4).

Assay of Inhibitory Effects on HAase [5]

HA sodium salt from rooster comb (Wako Pure Chemical Industries Ltd., Osaka, Japan, 0.25 mg/ml), HAase from bovine testes (Wako Pure Chemical Industries Ltd., Osaka, Japan), and compound 48/80 (Sigma Chemical Co. Ltd., St. Luis, U.S.A., 3.0 mg/ml) containing NaCl (17 mM) and CaCl₂ (27 mM) were dissolved in 0.1 M acetate buffer (pH 4.0), respectively. The test samples were dissolved in dimethylsulfoxide for MeOH extract and BuOH extract (2.0 mg/ml) and in acetate buffer for **1** (0.5~10 mM). *p*-Dimethylaminobenzaldehyde (Wako Pure Chemical Industries Ltd., Osaka, Japan, 20 mg/ml) was dissolved in conc. HCl-acetic acid (1:39). All aqueous solutions were prepared using water filtered through a Milli-Q water system (Millipore, Belford, U.S.A.). All chemicals were of reagent grade.

HAase (460 units/ml, 130 μl), which had been preincubated with the test sample (20 ml) at 37°C for 10 minutes in advance, was incubated with compound 48/80 (100 μl) at 37°C for 10 minutes. After the incubation HA sodium salt (250 μl) was added and the reaction mixture was incubated at 37°C for 40 minutes. Then the reaction was stopped by adding 0.4 M NaOH (100 μl), and then the mixture was allowed to stand for 10 minutes in ice water.

The inhibitory effect was determined by the modified Morgan-Elson method [6]. The reaction mixture was boiled for 3 minutes after adding borate buffer (pH 9.1) (100 μl) and then allowed to stand for 10 minutes in ice water. After centrifuging (10,000 rpm, 3 minutes), the supernatant (140 μl) was incubated with *p*-dimethylaminobenzaldehyde solution (300 μl) at 37°C for 20 minutes. After 20 minutes, the color reaction was detected by the absorbance at 585 nm with a microplate reader Bio-Rad Model 550.

Test samples were replaced with the buffer solution for the control, while the enzyme solution was replaced with the buffer solution for the blank. Percent inhibition was calculated as follows: Inhibition (%)=[(A-B)-(C-D)]/(A-B)×100. A, control OD; B, control blank OD; C, sample OD; D, sample blank OD.

The 50% inhibitory concentration (IC₅₀) was calculated using the mean of 3 observation at each of the 5 concentrations.

Acknowledgements We are grateful to Miss N. Kobayashi, Dr. M. Ikegami and Dr. H. Kasai and of Hoshi University for NMR and mass measurements.

References

1. Laurent TC, Fraser JRE. Hyaluronan. *FASEB J* 6: 2397-2405 (1992)
2. Alexander B, Daniel JR, Stephan B, Masatoshi N, Sumnhild S, Julia H, Günther B, Stefan D, Mark JJ, Armin B. L-Ascorbic acid 6-hexadecanoate, a potent hyaluronidase inhibitor. *J Biol Chem* 44: 45990-45997 (2004)
3. Wat C-K, Neil Towers GH. Production of methylated phenolic acids by species of *Lentinus* (Bashidiomycete). *Phytochem* 16: 290-291 (1977)
4. Chu M, Mierzwa R, Xu L, He L, Terracciano J, Patel M, Zhao W, Black TA, Chan T-M. Structure of Sch 419560, a novel α-pyrone antibiotic produced by *Pseudomonas fluorescens*. *J Antibiot* 55: 215-218 (2002)
5. Kobayashi M, Matsushita H, Yoshida K, Tsukiyama R, Sugiyama T, Yamamoto K. *In vitro* and *in vivo* anti-allergenic activity of soy sauce. *Inter J Molec Medicine* 14: 879-884 (2004)
6. Fujitani N, Sakaki S, Yamaguchi Y, Takenaka H. Inhibitory effects of microalgae on the activation of hyaluronidase. *J Appl Physiol* 13: 489-492 (2000)
7. Spickenreiter M, Braun S, Bernhardt G, Dove S, Buschauer A. Novel 6-*O*-acylated vitamin C derivatives as hyaluronidase inhibitors with selectivity for bacterial lyases. *Bioorg Med Chem Lett* 16: 5313-5316 (2006)