

ERRATUM

Polystyrene/Polybutadiene Blends: An Analysis of the Phase-Inversion Region and Cophase Continuity and a Comparison with Theoretical Predictions

Susan Joseph,¹ Maria Rutkowska,² Mariola Jastrzêbska,² Helena Janik,³ Jozef T. Haponiuk,³ Sabu Thomas⁴

¹*St. Stephen's College, Pathanapuram, Kerala, India, 689695*

²*Gdynia Maritime Academy, 83 Morska Street, 81-225, Gdynia, Poland*

³*Polymer Tech Department, Chemical Faculty, Technical University of Gdansk, 80-952, Gdansk, Poland*

⁴*School of Chemical Sciences, Mahatma Gandhi University, Priyadarshini Hills P.O., Kottayam, Kerala, India, 686560*

[Article in J Appl Polym Sci 2003, 89, 1007]

The affiliations were originally printed incorrectly. They are correct as printed above.

Table IV was originally printed incorrectly. It is correct as shown here:

TABLE IV
Results of Disintegration Tests of the
Different Blend Series

Sample code	Dissolution etchant	
	Butan-2-one for PS phase	n-Heptane for PB phase
S ₁₀	ND	D
S ₂₀	ND	D
S ₃₀	ND	D
S ₄₀	PD	D
S ₅₀	D	D
S ₆₀	D	PD
S ₇₀	D	ND
S ₈₀	D	ND

D = disintegration; ND = no disintegration; PD = partial disintegration.