

Axisymmetric Harmonic Infrapolynomials in R^N

MORRIS MARDEN*

Department of Mathematics, University of Wisconsin, Milwaukee, Wisconsin 53201

Communicated by Oved Shisha

Received July 17, 1970

THIS PAPER IS DEDICATED TO JOSEPH L. WALSH, ON THE OCCASION OF HIS SEVENTY-FIFTH BIRTHDAY ANNIVERSARY.

1. INTRODUCTION

During the last two decades we have witnessed an intensive development of the subject of infrapolynomials on sets $\omega \subset C$. As we recall, an infrapolynomial on ω is a polynomial $p \in P = \{\zeta^n + \sum_1^n a_k \zeta^{n-k}\}$ such that there exists no other $q \in P$ for which $q(\zeta) = 0$ for $\zeta \in \omega' = \{\zeta \in \omega : p(\zeta) = 0\}$ and $|q(\zeta)| < |p(\zeta)|$ for $\zeta \in \omega - \omega'$. A leader in this development was Professor Walsh, the man whom we are honoring and of whom I was privileged to be the first Ph. D. student.

In the present paper we attempt a parallel development for harmonic infrapolynomials on three-dimensional sets. Our results will be expressed in terms of three coordinate systems in R^3 : rectangular (x, y, z) ; cylindrical (x, ρ, ϕ) with

$$\rho^2 = y^2 + z^2, \quad y = \rho \cos \phi, \quad z = \rho \sin \phi;$$

and spherical (r, θ, ϕ) with

$$x = r \cos \theta, \quad \rho = r \sin \theta. \tag{1.1}$$

By an *axisymmetric function* in R^3 we mean one that is independent of ϕ ; that is, a function which assumes the same value at all points of the circle $x = x_0, \rho = \rho_0$ [abbreviated: circle (x_0, ρ_0)]. As the domain of such a function, we take an *axisymmetric set* Ω in R^3 ; that is a set such that, if point $(x_0, \rho_0, \phi_0) \in \Omega$, also point $(x_0, \rho, \phi) \in \Omega$ for all ρ and $\phi, 0 \leq \rho \leq \rho_0$ and $0 \leq \phi \leq 2\pi$. Thus an axisymmetric set Ω may consist of points on the x -axis,

* The work on this paper was done partly under N.S.F. grant No. GP-19615. Abstract appeared in Notices, *Amer. Math. Soc.* 117 (1970), 1072.

circular disks having their centers on the x -axis and their planes perpendicular to the x -axis, and the interiors of surfaces of revolutions which are cut in a single circle by any plane perpendicular to the x -axis. The meridian section $\omega \subset \mathbb{C}$ of Ω is an *axiconvex* region, meaning that $\zeta \in \omega$ implies $\lambda\zeta + (1 - \lambda)\bar{\zeta} \in \omega$ for all real λ , $0 \leq \lambda \leq 1$.

Let us first consider axisymmetric harmonic polynomials $H(x, \rho)$ of degree n . As is well known [5, p. 254], every such polynomial can be written in the form

$$H(x, \rho) = \sum_{j=0}^n a_j r^j P_j(x/r), \tag{1.2}$$

where $P_j(u)$ is the Legendre polynomial of degree j .

Of special importance is the class

$$\mathbf{H} = \{H(x, \rho): a_n = 1\} \tag{1.3}$$

of axisymmetric harmonic polynomials with leading coefficient of one and therefore strictly of degree n . Let us compare two polynomials $P(x, \rho) \in \mathbf{H}$ and $Q(x, \rho) \in \mathbf{H}$ on a given axisymmetric set Ω in \mathbb{R}^3 relative to some suitably defined norm $\|H(x, \rho)\|$. We say that Q is an underpolynomial of P on Ω if

$$\|Q(x, \rho)\| < \|P(x, \rho)\| \tag{1.4}$$

for all circles $(x, \rho) \subset \Omega$. Let us denote by $U(P, \Omega)$ the class of all underpolynomials of P on Ω . If $U(P, \Omega) = \emptyset$ for some $P \in \mathbf{H}$, we say that P is an *axisymmetric harmonic infrapolynomial on Ω* .

In the sequel we shall investigate the properties of the class $\mathbf{I}(\Omega)$ of axisymmetric harmonic infrapolynomials on a given axisymmetric set Ω . We shall determine some conditions on $P \in \mathbf{H}$ in order that $P \in \mathbf{I}(\Omega)$ and also determine the location of the zeros of all $P \in \mathbf{I}(\Omega)$ in relation to the set Ω . In order to do this, we shall bring together the methods of two hitherto disjoint disciplines the theory of infrapolynomials on sets $\omega \subset \mathbb{C}$ and the theory of a certain integral operator, whose development is largely due to Professor Stefan Bergman; see [1 and 2].

2. INTEGRAL REPRESENTATIONS FOR $H(x, \rho)$ AND $\|H(x, \rho)\|$

Let us define as the *associate* of $H(x, \rho)$ the polynomial

$$h(\zeta) = \sum_{k=0}^n a_k \zeta^k, \quad a_n = 1, \quad \zeta \in \mathbb{C}. \tag{2.1}$$

Thus ζ^n is the associate of $r^n P_n(x/r)$.

In view of the Whittaker formula [4, p. 312–315],

$$r^k P_k(x/r) = (1/2\pi) \int_0^{2\pi} (x + i\rho \cos t)^k dt, \quad (2.2)$$

we have the result:

THEOREM 2.1. *Let $H(x, \rho)$ be an axisymmetric harmonic polynomial and let $h(\zeta)$ be its associate.*

Then

$$H(x, \rho) = (1/2\pi) \int_0^{2\pi} h(x + i\rho \cos t) dt. \quad (2.3)$$

More generally, if $f(\zeta)$ is analytic in a region ω which is the meridian section of an axisymmetric region Ω , then

$$F(x, \rho) = (1/2\pi) \int_0^{2\pi} f(x + i\rho \cos t) dt \quad (2.4)$$

satisfies Laplace's equation $\nabla^2 F = 0$ and so is an axisymmetric harmonic function in Ω . In fact, (2.3) and (2.4) are special cases of the operator introduced by Bergman [1, p. 43]:

$$F(x, y, z) = (1/2\pi i) \int_{|\tau|=1} f(\zeta, \tau) \tau^{-1} d\tau, \quad (2.5)$$

acting upon the function $f(\zeta, \tau)$ that is analytic in ζ on some region in \mathbf{C} and continuous in τ for $|\tau| = 1$. On setting

$$\zeta = x + (1/2)(yi + z)\tau + (1/2)(yi - z)\tau^{-1}, \quad (2.6)$$

the operator transforms $f(\zeta, \tau)$ into the function $F(x, y, z)$ which together with $\mathscr{R}F(x, y, z)$ and $\mathscr{I}F(x, y, z)$ is harmonic in a certain region of \mathbf{R}^3 .

In view of the integral representation (2.3) for an axisymmetric harmonic polynomials $H(x, \rho)$, it is natural to define the norm $\|H(x, \rho)\|$ of $H(x, \rho)$ by the formula

$$\|H(x, \rho)\|^2 = (1/2\pi) \int_0^{2\pi} |h(x + i\rho \cos t)|^2 d\sigma(t). \quad (2.7)$$

Here and in the subsequent formulas, $\sigma(t)$ denotes a monotonically increasing function for $0 \leq t \leq 2\pi$. In the special case $\sigma(t) \equiv t$, we denote as norm $\|H(x, \rho)\|_t$. Thus,

$$\|r^n P_n(x/r)\|^2 = (1/2\pi) \int_0^{2\pi} (x^2 + \rho^2 \cos^2 t)^n d\sigma(t).$$

More generally, using (1.2), we may expand (2.7) as a hermitian form in the a_k , the coefficients of which form are homogeneous polynomials in x and ρ .

Let us now consider the harmonic polynomial $H(x, \rho)$ which has for its associate

$$h(\zeta) = p(\zeta) q(\zeta), \tag{2.8}$$

where p and q are, respectively, polynomials of degrees k and $n - k$. Let us denote by $P(x, \rho)$ and $Q(x, \rho)$ the axisymmetric harmonic polynomials which have $p(\zeta)$ and $q(\zeta)$, respectively, as associates. To indicate a kind of factor relation among $H(x, \rho)$, $P(x, \rho)$ and $Q(x, \rho)$, we follow Bergman in defining the operation

$$P(x, \rho) * Q(x, \rho) = (1/2\pi) \int_0^{2\pi} p(x + i\rho \cos t) q(x + i\rho \cos t) d\sigma(t). \tag{2.9}$$

Thus, whereas the product $P(x, \rho) Q(x, \rho)$ is not ordinarily harmonic, the product $P(x, \rho) * Q(x, \rho)$ is harmonic and so the operation converts the family of axisymmetric harmonic polynomials into an algebra.

Obviously we may express the norm of any axisymmetric harmonic polynomial $H(x, \rho)$ in terms of the product in (2.9), as follows.

THEOREM 2.2. *If $H(x, \rho)$ is any axisymmetric harmonic polynomial, its norm $\| H(x, \rho) \|$ as defined by (2.7) satisfies the relation*

$$\| H(x, \rho) \|^2 = H(x, \rho) * \overline{H(x, \rho)}. \tag{2.10}$$

The product $P(x, \rho) * \overline{Q(x, \rho)}$ is in general not a harmonic function but serves the purpose of "inner vector product" in the space of axisymmetric harmonic function.

We now prove the following theorem.

THEOREM 2.3. *Let $P(x, \rho)$ and $Q(x, \rho)$ be any two axisymmetric harmonic polynomials. Then*

$$| P(x, \rho) * Q(x, \rho) | \leq \| P(x, \rho) \| \| Q(x, \rho) \|. \tag{2.11}$$

Proof. Using (2.9) and Schwarz inequality, we infer that

$$\begin{aligned} | P(x, \rho) * Q(x, \rho) | &\leq (1/2\pi) \int_0^{2\pi} | p(x + i\rho \cos t) q(x + i\rho \cos t) | d\sigma \\ &\leq \left\{ (1/2\pi) \int_0^{2\pi} | p(x + i\rho \cos t) |^2 d\sigma \right\}^{1/2} \\ &\quad \times \left\{ (1/2\pi) \int_0^{2\pi} | q(x + i\rho \cos t) |^2 d\sigma \right\}^{1/2}. \end{aligned}$$

That is, (2.11) is valid for all (x, ρ) .

If we choose $Q(x, \rho) = 1$ and $\sigma(t) = t$ in Theorem 2.3, we obtain the following result.

COROLLARY 2.1. *If $H(x, \rho)$ is an axisymmetric harmonic polynomial, then for all (x, ρ)*

$$|H(x, \rho)| \leq \|H(x, \rho)\|_t. \tag{2.12}$$

As may be seen from (2.7), the equality sign holds in both (2.11) and (2.12) when $\rho = 0$ and when P, Q and H are each constants, but does not seem to hold in any other case.

3. STRUCTURE OF AXISYMMETRIC HARMONIC INFRAPOLYNOMIALS

We shall now use well-known theorems about the structure of infrapolynomials on $\omega \subset \mathbb{C}$ in order to get some corresponding results regarding axisymmetric harmonic infrapolynomials on $\Omega \subset \mathbb{R}^3$. It will be helpful first to prove the following.

THEOREM 3.1. *Let $P(x, \rho) \in \mathbf{H}$, the class of axisymmetric harmonic polynomials defined by (1.3). Let $\omega \subset \mathbb{C}$ be a bounded axiconvex region and $\Omega \subset \mathbb{R}^3$ be the axisymmetric region having ω as its meridian section. If $P(x, \rho)$ is an infrapolynomial on the closure $\bar{\Omega}$ of Ω , then its associate $p(\zeta)$ is an infrapolynomial on the closure $\bar{\omega}$ of ω .*

Proof. If the contrary were true, p would have an underpolynomial q on ω ; that is,

$$|q(\zeta)| = |p(\zeta)| \quad \text{for } \zeta \in \omega' = \{\zeta \in \bar{\omega} : p(\zeta) = 0\}, \tag{3.1}$$

$$|q(\zeta)| < |p(\zeta)| \quad \text{for } \zeta \in \bar{\omega} - \omega'. \tag{3.2}$$

Let $q(\zeta)$ be the associate of $Q(x, \rho)$. Clearly, $Q(x, \rho) \in \mathbf{H}$ and

$$\begin{aligned} \|Q(x, \rho)\|^2 &= (1/2\pi) \int_0^{2\pi} |q(x + i\rho \cos t)|^2 d\sigma \\ &< (1/2\pi) \int_0^{2\pi} |p(x + i \cos t)|^2 d\sigma = \|P(x, \rho)\|^2. \end{aligned}$$

Hence, $P(x, \rho)$ would have an underpolynomial $Q(x, \rho)$ on $\bar{\Omega}$, contradicting the hypothesis that $P(x, \rho)$ is an infrapolynomial on $\bar{\Omega}$.

For example, since in \mathbb{C} ζ^n is an infrapolynomial on the unit disk $|\zeta| \leq 1$,

we infer that, in \mathbf{R}^3 , $r^n P_n(x/r)$ is an infrapolynomial on the unit ball $x^2 + \rho^2 \leq 1$.

We now propose to use Theorem 3.1 in conjunction with the following well-known result due to Fekete [3, pp. 15–19].

THEOREM 3.2. *Let E , a closed bounded set in \mathbf{C} containing at least $n + 1$ points, have an infrapolynomial $p(\zeta)$, with $p(\zeta) \neq 0$ for $\zeta \in E$. Then there exist an integer m with $n \leq m \leq 2n$, a set of $m + 1$ constants $\lambda_j > 0$ with $\lambda_0 + \lambda_1 + \dots + \lambda_m = 1$ and a set of $m + 1$ points $\{\zeta_0, \zeta_1, \dots, \zeta_m\} \subset E$ such that $p(\zeta)$ is a factor of the polynomial*

$$f(\zeta) = \sum_{k=0}^m \lambda_k \psi_k(\zeta) \tag{3.3}$$

where

$$\psi_k(\zeta) = (\zeta - \zeta_0) \cdots (\zeta - \zeta_{k-1})(\zeta - \zeta_{k+1}) \cdots (\zeta - \zeta_m). \tag{3.4}$$

In applying Theorem 3.2, we need to choose E to be a bounded axiconvex region ω .

For, the integration in (2.3) requires that, if point $x + i\rho \in \omega$, then also point $x + i\rho \cos t \in \omega$ for $0 \leq t \leq 2\pi$. In view of Theorems (3.1) and (3.2), we are led now to the following theorem.

THEOREM 3.3. *Let Ω be a bounded axisymmetric region in \mathbf{R}^3 and let $P(x, \rho)$ be an n -th degree axisymmetric harmonic infrapolynomial on the closure of Ω . Then there exist an integer m , $n \leq m \leq 2n$, a set of $m + 1$ constants $\lambda_j > 0$ with $\lambda_0 + \lambda_1 + \dots + \lambda_m = 1$, a set of circles*

$$(x_0, \rho_0; x_1, \rho_1; \dots; x_m, \rho_m) \subset \Omega,$$

and an axisymmetric harmonic polynomial $G(x, \rho)$ of degree $m - n$ such that

$$\sum_{k=0}^m \lambda_k \Psi_k(x, \rho) = P(x, \rho)^* G(x, \rho), \tag{3.5}$$

where $\Psi_k(x, \rho)$ is the axisymmetric harmonic polynomial

$$\Psi_k(x, \rho) = -(1/2\pi)(\partial/\partial x_k) \int_0^{2\pi} \prod_{j=0}^m (x + i\rho \cos t - x_j - i\rho_j) dt. \tag{3.6}$$

Proof. Since $P(x, \rho)$ is an infrapolynomial on Ω , its associate $p(\zeta)$ is by Theorem 3.1 an infrapolynomial on ω , the meridian section of Ω . By Theorem 3.2, there is a polynomial $g(\zeta)$ of degree $m - n$ such that

$f(\zeta) = p(\zeta)g(\zeta)$. Hence $f(\zeta)$ is the associate of $P(x, \rho)^*G(x, \rho)$, where $G(x, \rho)$ is the axisymmetric harmonic polynomial having $g(\zeta)$ as associate. On the other hand, $f(\zeta)$ may be written in the form (3.3) and so we are led to (3.5).

4. NULL CIRCLES OF AXISYMMETRIC HARMONIC INFRAPOLYNOMIALS

By a *null circle* (x_0, ρ_0) of an axisymmetric harmonic polynomial $H(x, \rho)$ we mean that $H(x_0, \rho_0) = 0$. A null circle is therefore the intersection of the level surfaces $S_1 : \mathcal{R}H(x, \rho) = 0$ and $S_2 : \mathcal{I}H(x, \rho) = 0$. The set of all null circles of $H(x, \rho)$ is finite unless $\mathcal{R}H(x, \rho) \equiv 0$ or $\mathcal{I}H(x, \rho) \equiv 0$ when it is the entire level surface S_2 or S_1 , respectively.

Let us first recall the following result about the zeros of an infrapolynomial on $\omega \subset \mathbb{C}$, due to Féjer [3, p. 23].

THEOREM 4.1. *Let E be a closed bounded set in \mathbb{C} and $p(\zeta)$ an infrapolynomial on E . Then all the zeros of p lie in the convex hull of E .*

In applying Theorem 4.1, we must again replace E by a bounded axiconvex region ω which is the meridian section of an axisymmetric region Ω . Let us denote by c_1 and c_2 the two points which are on the real axis, left and right of ω , respectively, and from which ω subtends an angle of π/n . Thus, ω lies in the intersection of the two sectors

$$-(\pi/2n) \leq \arg(\zeta - c_1) \leq (\pi/2n), \tag{4.1}$$

$$\pi - (\pi/2n) \leq \arg(\zeta - c_2) \leq \pi + (\pi/2n). \tag{4.2}$$

Let us denote by $K_1(\omega, n)$ and $K_2(\omega, n)$ the cones obtained on revolving about the axis of reals the two sectors

$$\pi - (\pi/2/n) \leq \arg(\zeta - c_1) \leq \pi + (\pi/2n) \tag{4.3}$$

$$-\pi/2n \leq \arg(\zeta - c_2) \leq \pi/2n. \tag{4.4}$$

Alternatively, to obtain, for example, $K_2(\omega, n)$ geometrically, we may take a double nappe cone of vertex angle π/n and slide it as far as possible to the left with its axis along the x -axis and yet have the left nappe contain Ω . The right nappe is then $K_2(\omega, n)$.

We are now in a position to establish

THEOREM 4.2. *Let $\omega \subset \mathbb{C}$ be a bounded axiconvex region, which is the meridian section of Ω , an axisymmetric region in \mathbb{R}^3 . Let $P(x, \rho)$ be an axi-*

symmetric harmonic infrapolynomial on Ω . Then no circle (x, ρ) for which $P(x, \rho) = 0$ may lie in either cone:

$$K_j(\omega, n): \quad 0 < \rho \leq (-1)^j(x - c_j) \tan(\pi/2n), \quad j = 1, 2, \quad (4.5)$$

where the c_j are defined as above.

Proof. By Theorem 3.1, the associate $p(\zeta)$ of $P(x, \rho)$ is an infrapolynomial on ω and by Theorem 4.1 all the zeros $\zeta_j, j = 1, 2, \dots, n$ of $p(\zeta)$ lie in the convex hull κ of ω . The region κ also lies in the intersection of the two sectors (4.1) and (4.1).

Let us write

$$p(\zeta) = (\zeta - \zeta_1)(\zeta - \zeta_2) \cdots (\zeta - \zeta_n)$$

and thus

$$P(x, \rho) = (1/2\pi) \int_0^{2\pi} \prod_{j=1}^n (x + i\rho \cos t - \zeta_j) dt.$$

Let us assume that there is a circle (x_0, ρ_0) in the cone $K_2(\omega, n)$ such that $P(x_0, \rho_0) = 0$. Then

$$\int_0^{2\pi} w(t) dt = 0, \quad (4.5)$$

where

$$w(t) = \prod_{j=1}^n (\zeta_j - x_0 - i\rho_0 \cos t). \quad (4.6)$$

These assumptions require point $x_0 + i\rho_0$ to lie in sector (4.4) and therefore point $x_0 + i\rho_0 \cos t$ also to lie interior to sector (4.4) for all $t, 0 \leq t \leq 2\pi$. Since $\zeta_j \in \kappa$ for $j = 1, 2, \dots, n$ and since κ lies in sector (4.2), it follows that

$$\pi - (\pi/2n) < \arg(\zeta_j - x_0 - i\rho_0 \cos t) < \pi + (\pi/2n)$$

for each j and, because of (4.6),

$$n\pi - (\pi/2) < \arg w(t) < n\pi + (\pi/2).$$

Hence, $\Re[e^{-n\pi i} w(t)] > 0$ for $0 \leq t \leq 2\pi$ and thus $\Re[e^{-\pi i} \int_0^{2\pi} w(t) dt] > 0$. This contradicts (4.5) and thus the assumption that $P(x_0, \rho_0) = 0$ for circle $(x_0, \rho_0) \subset K_2(\omega, n)$, is invalid. Using similar reasoning for circles $(x_0, \rho_0) \subset K_1(\omega, n)$, we complete the proof of Theorem 4.2.

Remark 1. Theorem 4.2 remains valid if $\|H(x, \rho)\|$ as defined by (2.7) is replaced by any other norm for which Theorem 3.1 is true.

5. GENERALIZATION TO \mathbf{R}^N

We shall extend the preceding results to axisymmetric harmonic functions $F(x_1, x_2, \dots, x_N)$ of N real variables; that is, solutions of the Laplace equation

$$\Delta^2 F = \sum_{j=1}^N (\partial^2 F / \partial x_j^2) = 0. \quad (5.1)$$

The axisymmetric case corresponds to the one in which F is a function just of x and ρ where

$$x = x_1, \quad \rho^2 = x_2^2 + x_3^2 + \dots + x_N^2. \quad (5.2)$$

In this case (5.1) reduces to

$$(\partial/\partial x)(\rho^{N-2} \partial F/\partial x) + (\partial/\partial \rho)(\rho^{N-2} \partial F/\partial \rho) = 0. \quad (5.3)$$

On introducing polar coordinates into Eq. (5.3)

$$x = r \cos \theta, \quad \rho = r \sin \theta$$

and using the method of separating variables, we find the basic solutions of (5.3) in the form

$$r^n P_n^{(\mu)}(\cos \theta), \quad 2\mu = N - 2, \quad (5.4)$$

where $P_n^{(\mu)}(\cos \theta) = P_n^{(\alpha, \alpha)}(\cos \theta)$, $2\alpha = N - 3$, and where $P_n^{(\alpha, \beta)}(\cos \theta)$ and $P_n^{(\mu)}(\cos \theta)$ are, respectively, the Jacobi and Gegenbauer polynomials of degree n .

We are thus led to consider the axisymmetric harmonic polynomials in \mathbf{R}^N

$$H(x, \rho) = \sum_{j=0}^n A_j r^j P_j^{(\mu)}(x/r), \quad 2\mu = N - 2, \quad (5.5)$$

with $A_n = 1$. For such a polynomial the following holds.

THEOREM 5.1. *The harmonic function (5.5) may be written in the form*

$$H(x, \rho) = 2^{3-N} \Gamma(\mu)^{-2} \int_0^\pi h(x + i\rho \cos t) \sin^{N-3} t \, dt \quad (5.6)$$

$$h(\zeta) = \sum_{j=0}^n a_j \zeta^j, \quad (5.7)$$

with $a_j = [\Gamma(j + 2\mu)/j!] A_j$.

Proof. The expression (5.5) follows directly from the representation [2, p. 167]

$$r^n P_n^{(\mu)}(\cos \theta) = \frac{2^{1-2\mu} \Gamma(n + 2\mu)}{n! \Gamma(\mu)^2} \int_0^\pi (x + i\rho \cos t)^n \sin^{N-3} t \, dt. \quad (5.8)$$

We refer to the polynomial in (5.7) as the *associate* of the polynomial $H(x, \rho)$ given by (5.5).

By analogy with Section 2, we define the norm $\|H(x, \rho)\|$ by the expression

$$\|H(x, \rho)\|^2 = 2^{3-N} \Gamma(\mu)^{-2} \int_0^\pi |h(x + i\rho \cos t)|^2 \sin^{N-3} t \, d\sigma(t). \quad (5.9)$$

If now we are given two polynomials $P(x, \rho)$ and $Q(x, \rho)$ of type (5.5), we say that $Q(x, \rho)$ is an underpolynomial of $P(x, \rho)$ on an axisymmetric region $\Omega \subset \mathbf{R}^N$ if $\|Q(x, \rho)\| < \|P(x, \rho)\|$ for all $(x, \rho) \in \Omega$ and that $P(x, \rho)$ is an axisymmetric harmonic infrapolynomial on $\bar{\Omega}$ if it has no underpolynomial $Q(x, \rho)$ on $\bar{\Omega}$.

By the same reasoning as for Theorem 3.1, we may establish the following:

THEOREM 5.2. *Let $\omega \subset \mathbf{C}$ be a bounded axiconvex region and let $\Omega \subset \mathbf{R}^N$ be the region comprising the loci $x = x_0, x_2^2 + x_3^2 + \dots + x_N^2 = \rho_0^2$ for all $x_0 + i\rho_0 \in \omega$. If $P(x, \rho)$ is an axisymmetric harmonic infrapolynomial on $\bar{\Omega}$, its associate $p(\zeta)$ is an infrapolynomial on $\bar{\omega}$.*

Again, since (5.6) differs from (2.3) principally because of the nonnegative factors $2^{3-N} \Gamma(\mu)^{-2} \sin^{N-3} t$ in (5.5), we may use the same reasoning as for Theorem 4.2 to show the following theorem to be valid.

THEOREM 5.3. *Let $\omega \subset \mathbf{C}$ be a bounded axiconvex in \mathbf{C} and let $\Omega \subset \mathbf{R}^N$ be the region comprising the loci $x_1 = x_0, x_2^2 + x_3^2 + \dots + x_N^2 = \rho_0^2$ for all $x_0 + i\rho_0 \in \omega$. If $P(x, \rho)$ is an axisymmetric harmonic infrapolynomial on $\bar{\Omega}$, then no locus (x_0, ρ_0) for which $P(x, \rho) = 0$ has points in either of the cones*

$$0 < (x_2^2 + x_3^2 + \dots + x_N^2)^{1/2} \leq \pm(x_1 - c_j) \tan(\pi/2n) \quad (5.10)$$

where the c_j are defined as for Theorem 4.2. for $j = 1, 2$.

Also results analogous to Theorems 2.3 and 3.3 are valid, but their statement and proof are left to the reader.

6. EXTENSION TO CERTAIN OTHER HARMONIC INFRAPOLYNOMIALS IN \mathbf{R}^3

Let us finally consider harmonic polynomials of the form

$$F(x, y, z) = \sum_{j=J}^n A_j r^j P_j^{m(n-j)}(x/r) \cos m(n-j)\phi, \tag{6.1}$$

where $A_n = 1$; m and J are integers with $m > 0$, $J \geq [mn/(m+1)]$, and $P_j^k(\cos \theta)$ is the ‘‘associated Legendre function’’ [4, p. 323]. Clearly, $F(x, y, z)$ is a harmonic polynomial, but not ordinarily axisymmetric. We may show that a representation of $F(x, y, z)$ in the form (2.5) is possible on choosing as associate

$$f(\zeta, \tau) = \tau^{-m} f_0(\tau^m \zeta), \tag{6.2}$$

where

$$f_0(\zeta) = \sum_{j=J}^n a_j \zeta^j = \zeta^J f_1(\zeta) \tag{6.3}$$

with

$$a_j = [j + m(n-j)]! / j! A_j$$

and ζ given by (2.6) or, since $\tau = e^{ti}$, equivalently, by

$$\zeta = x + i(y \cos t + z \sin t) = x + i\rho \cos(t - \phi). \tag{6.4}$$

We may deduce the desired relation directly from the formula [4, p. 326]

$$r^n P_n^k(\cos \theta) = \frac{(j+k)!}{j!(2\pi)} \int_0^{2\pi} (x + i\rho \cos t)^n e^{-kti} dt.$$

That is,

$$F(x, y, z) = (1/2\pi) \int_0^{2\pi} f(x + i\rho \cos(t - \phi), e^{ti}) dt. \tag{6.5}$$

We next define the norm $\|F(x, y, z)\|$ in terms of the variables x, y, z or x, ρ, ϕ in such a way that the norm has the integral representation

$$\|F(x, y, z)\|^2 = (1/2\pi) \int_0^{2\pi} |f(x + i\rho \cos(t - \phi), e^{ti})|^2 d\sigma(t). \tag{6.6}$$

We then say that $F(x, y, z)$ is an infrapolynomial on a given region $\Omega \subset \mathbf{R}^3$ if no polynomial $G(x, y, z)$ of the same type as (6.1) exists such that $\|G(x, y, z)\| < \|F(x, y, z)\|$ for all $(x, y, z) \in \Omega$.

By reasoning similar to that in the proof of Theorem 3.1, we can now establish the following.

THEOREM 6.1. *Let $\omega \subset \mathbf{C}$ be a bounded axiconvex region and let $\Omega \subset \mathbf{R}^3$ be the axisymmetric region whose meridian cross section is ω . If the harmonic polynomial $F(x, y, z)$ given by (6.1) is an infrapolynomial on Ω , then the corresponding polynomial $f(\zeta)$ in (6.3) is an infrapolynomial on ω .*

According to Theorem 4.1, the zeros of the infrapolynomial

$$f_1(\zeta) = \prod_{j=1}^{n-J} (\zeta - \zeta_j)$$

lie in the convex hull κ of ω . Accordingly, since

$$f(\zeta, \tau) = e^{-m\tau i} e^{Jm\tau i} \zeta^J \prod_{j=1}^{n-J} (e^{m\tau i} \zeta - \zeta_j) \tag{6.8}$$

$$f(\zeta, \tau) = \zeta^J \prod_{j=1}^{n-J} (\zeta - \zeta_j e^{-m\tau i})$$

the zeros $\zeta_j e^{-m\tau i}$ therefore lie in the disk $|\zeta| \leq \delta$, where $\delta = \max |\zeta|$ for $\zeta \in \omega$.

If now $F(x, y, z)$ is an infrapolynomial on Ω and if $F(x_0, y_0, z_0) = 0$, then according to (6.5) and (6.8),

$$\int_0^{2\pi} w(t) dt = 0,$$

where

$$w(t) = (0 - x_0 - i\rho_0 \cos(t - \varphi_0))^J \prod_{j=1}^{n-J} [\zeta_j e^{-mti} - x_0 - i\rho_0 \cos(t - \varphi_0)].$$

From here on, the reasoning is similar to that for Theorem 4.2. We thus arrive at the following result.

THEOREM 6.2. *Let $\Omega \subset \mathbf{R}^3$ be an axisymmetric region whose meridian section is a bounded axiconvex region ω .*

Let $\delta = \max |\zeta|$ for $\zeta \in \omega$. If $F(x, y, z)$ given by (6.1) is a harmonic infrapolynomial on Ω , then no point (x, y, z) for which $F(x, y, z) = 0$ lies in either of the cones:

$$0 < \rho \leq \pm x \tan(\pi/2n) - \delta \sec(\pi/2n). \tag{6.9}$$

REFERENCES

1. S. BERGMAN, Integral operators in the theory of linear partial differential equations, in "Ergebnisse der Math. u. Ihrer Grenzgebiete," Heft 23, Springer-Verlag, Berlin 1961.
2. R. P. GILBERT, "Function Theoretic Methods in Partial Differential Equations," Academic Press, New York 1969.
3. M. MARDEN, "Geometry of Polynomials," Math. Surveys No. 3, American Mathematical Society, Providence, RI, 1966.
4. E. T. WHITTAKER AND G. N. WATSON, "A course of Modern Analysis," Amer. ed., Cambridge University Press, New York, 1943.
5. O. KELLOGG, "Foundations of Potential Theory," Springer-Verlag, Berlin, 1929.
6. M. MARDEN, Axisymmetric harmonic vectors, *Amer. J. Math.* 67 (1945), 109-122.