

Remarks on Finite Rank Projections

WILLIAM J. DAVIS

Department of Mathematics, Ohio State University, Columbus, Ohio 43210

Communicated by E. W. Cheney

Received November 25, 1970

If every n -dimensional subspace of X^* is the range of a projection of norm less than C , then every subspace of X with codimension n is the range of a projection having norm less than $1 + C$. Also, projection constants of finite-dimensional spaces are determined by finite-dimensional superspaces. It is further demonstrated that spheres cannot, in general, be nicely embedded into unit balls of finite-dimensional spaces.

This note is primarily concerned with the solution of some problems, stated in the paper of Cheney and Price [1], on projections of finite rank (that is, having finite-dimensional range) in Banach spaces. We see in Section 1 that a sphere cannot always be embedded nicely into the unit ball of a finite-dimensional space: In particular, if f_1, f_2 , and f_3 are in $l_1^{(3)}$ and if for x in $l_1^{(3)}$, $\{f_1(x)^2 + f_2(x)^2 + f_3(x)^2\}^{1/2} \geq \|x\|$, we must have $\|f_i\| > 1$ for some i . This gives a negative solution to part of problem 6 of [1].

The “principle of local reflexivity” of Lindenstrauss and Rosenthal [7] is extended, in the second section, to show that finite rank projections on a conjugate space X^* are, in a certain sense, near adjoints of finite rank projections on X . From this one easily deduces that if every n -dimensional subspace of X^* is complemented with norm $< c_n$, then every subspace of X having deficiency n is complemented with norm $< 1 + c_n$ (this gives an affirmative solution to problem 8 of [1]). From an unpublished result of Kadec to the effect that every n -dimensional subspace of every Banach space is complemented with norm $\leq n^{1/2}$, it follows that if Y has deficiency n in x and if $\epsilon > 0$, there is a projection of norm $< 1 + n^{1/2} + \epsilon$ of X onto Y . This result and the result of Kadec together with its proof, occur in [2].

Finally, the “compactness argument” of Lindenstrauss (see e.g. [6]) is applied directly to show that if Y is a finite-dimensional subspace of X and if P is a “best” (in terms of norm) projection of X onto Y , then $\|P\| = \sup \|R\|$ where the sup is over all “best” projections of Z onto Y , Z is finite-dimensional and $Y \subset Z \subset X$. This answers problem 9 of [1].

1

We show that, in general, spheres cannot be efficiently inscribed in unit balls of finite-dimensional spaces. Suppose that X is an n -dimensional space with norm $\| \cdot \|$. Suppose that there exist functionals f_1, \dots, f_n in the ball of $X^*(B_{X^*})$ such that

$$\left\{ \sum_{j=1}^n f_j(x)^2 \right\}^{1/2} = \|x\|_2 \geq \|x\|$$

for every x in X . There must be a vector x_k in $\bigcap_{i=1, i \neq k}^n \ker(f_i)$ such that $f_k(x_k) = 1 = \|x_k\|_2 \geq \|x_k\|$. Since $\|f_k\| \leq 1$, it follows that $\|x_k\| = \|f_k\| = 1$ for $k = 1, 2, \dots, n$ and that $f_i(x_j) = \delta_{ij}$. The system $(x_i; f_i)$ is called a *normal basis* for X and must satisfy the condition that $\text{sp}\{x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n\}$ is parallel to the supporting hyperplane to B_X at x_k (that is, $\{f_k(x) = 1\}$).

We are now able to show that the ball of $l_1^{(3)}$ has no such inscribed sphere.

THEOREM 1. *If $\{f_1, f_2, f_3\}$ are in*

$$l_\infty^{(3)} (= (l_1^{(3)})^*) \quad \text{and} \quad \{f_1(x)^2 + f_2(x)^2 + f_3(x)^2\}^{1/2} \geq \|x\|$$

for every x in $l_1^{(3)}$, then $\|f_i\| > 1$ for some i .

Proof. Suppose that there is a normal basis (as above) with $\|x\|_2 \geq \|x\|$ always. Then, notice that $x_i = (a_{i1}, a_{i2}, a_{i3})$ must have $|a_{ij}|$ different from zero for each i, j . This is due to the fact that since $\{\|x\|_2 = 1\}$ is tangent to B_X at x_1, x_2 and x_3 and since $\{\|x\|_2 = 1\} \subset B_X$ these are smooth points of the ball of $l_1^{(3)}$. We may as well assume that a_{11}, a_{12} , and a_{13} are all positive. Then $f_1 = (1, 1, 1)$. Since $(x_i; f_i)$ is a normal basis, we can conclude that $a_{21} + a_{22} + a_{23} = a_{31} + a_{32} + a_{33} = 0$. For definiteness, assume that $a_{21} > 0$, $a_{22} > 0$ and $a_{23} < 0$ (the argument will apply to all legitimate choices of sign for the a_{ij} 's). This condition forces $f_2 = (1, 1, -1)$. In turn, $a_{31} + a_{32} - a_{33} = 0$, so that $a_{33} = 0$. This is impossible in our situation, and proves the theorem.

2

Let us recall some elementary facts and notation which will be used here. If R is a finite rank projection on X , then $R: X \rightarrow X$ and R^* is a finite rank projection on X^* . If $\{x_1, \dots, x_n\}$ are in X , then $[x_1, \dots, x_n]$ is to denote the linear span the x_i 's in X . If T is a map from X to Y and W is a subspace of X denote the norm of $T|W$ by $\|T|W\|$.

Theorem 2 below is a modification of the “principle of local reflexivity” of Lindenstrauss and Rosenthal [7]. The author has recently learned that similar versions of this principle occur in [4] and [5]. One change in the proof is the use of the following lemma (suggested to the author by J. Daneman) instead of the separation lemma of Klee [3].

LEMMA. *Let C_1, \dots, C_n be open convex subsets of a Banach space X , and suppose $\bigcap \bar{C}_i^{w*}$ has a nonempty core. Then $\bigcap C_i \neq \emptyset$. (For a set A in X , \bar{A}^{w*} denotes its weak* closure in X^{**}).*

Proof. By induction, first consider $n = 2$ (the case $n = 1$ is trivial and the second case provides the proof of the lemma). Suppose $C_1 \cap C_2 = \emptyset$ so there is an f in X^* and a scalar α such that $f(C_1) < \alpha < f(C_2)$, then, $f(\bar{C}_1^{w*}) \leq \alpha \leq f(\bar{C}_2^{w*})$. Let Ψ in X^{**} be such that $\Psi(f) = 1$. Since there is a core point φ of $\bar{C}_1^{w*} \cap \bar{C}_2^{w*}$, there is $\delta > 0$ such that $|\lambda| < \delta$ implies $\varphi + \lambda\Psi$ is in $\bar{C}_1^{w*} \cap \bar{C}_2^{w*}$. This is incompatible with $(\varphi + \lambda\Psi)(f) = \alpha$ for all such λ , giving the desired contradiction. Now, assuming the conclusion for $n - 1$, let C_1, \dots, C_n satisfy the hypotheses so that $\emptyset \neq D = \bigcap_{j=2}^n C_j$. Let $\varphi \in \text{core } \bar{C}_2^{w*} \cap \dots \cap \bar{C}_n^{w*}$ and $\varphi \notin \bar{D}^{w*}$. Then there is an f in X^* such that $\varphi(f) > 1$ and $f(d) \leq 1$ for all d in D . However, letting $B_i = C_i \cap \{x \mid f(x) > 1\}$ for $i = 2, 3, \dots, n$, we see that the hypotheses for the case $n - 1$ apply to give $\emptyset \neq \bigcap_{j=2}^n B_j \subset D$ which is a contradiction. Thus $\text{core } \bar{C}_2^{w*} \cap \dots \cap \bar{C}_n^{w*} \subset \bar{D}^{w*}$ so that $\bar{C}_1^{w*} \cap \bar{D}^{w*}$ has a core. Now apply the argument for $n = 2$ to the pair C_1, D to see that $\emptyset \neq C_1 \cap D = \bigcap_{j=1}^n C_j$.

THEOREM 2. *Let P be a finite rank projection on X^* and let $\epsilon > 0$. Let V be any finite-dimensional subspace of X^* . Then there is a finite rank projection R on X such that $R^*(X^*) = P(X^*)$, $\|P - R^*\|_V < \epsilon$ and $\|R\| < \|P\| + \epsilon$.*

Proof. Let

$$Pf = \sum_{i=1}^n \varphi_i(f) f_i \quad \text{with} \quad \varphi_i(f_j) = \delta_{ij},$$

where $\{\varphi_1, \dots, \varphi_n\} \subset X^{**}$. Next choose $\{f_{n+1}, \dots, f_m\}$ in $[\varphi_1, \dots, \varphi_n]_{\perp}$ so that $\{f_1, \dots, f_m\}$ is a basis for $\text{sp}\{f_1, \dots, f_n, V\} = [f_1, \dots, f_n, V]$. Now for $\delta > 0$ and $\eta > 0$ (to be determined later), let $\{\Psi_i \mid 1 \leq i \leq p\}$ be a δ -net on the unit sphere of $[\varphi_1, \dots, \varphi_n]$ in X^{**} . Define the following open convex subsets of $X^n (= X \times \dots \times X)$, for $i = 1, 2, \dots, p$:

$$K_i = \left\{ (x_1, \dots, x_n) \mid \left\| \sum_{j=1}^n \Psi_i(f_j) x_j \right\| < 1 + \delta \right\}$$

and

$$D = \left\{ (x_1, \dots, x_n) \mid \sum_{i=1}^m \sum_{j=1}^n |f_i(x_j) - \delta_{ij}| < \eta \right\}.$$

Let K_i^{**} and D^{**} be the similarly defined subsets of $(X^{**})^n$. Then D^{**} is a weak*-open set containing $(\varphi_1, \dots, \varphi_n)$ and K_i^{**} is strongly open, containing $(\varphi_1, \dots, \varphi_n)$. It follows easily (as in [7; proof of Theor. 3.1]) that $\bar{K}_i^{w*} \supset K_i^{**}$, and $\bar{D}^{w*} \supset D^{**}$. The hypotheses of the lemma are now satisfied for the $p + 1$ sets K_1, \dots, K_p, D . Therefore, there is some (x_1, \dots, x_n) in X^n common to all of these sets. Now define $T: [\varphi_1, \dots, \varphi_n] \rightarrow [x_1, \dots, x_n]$ as the linear extension of $T\varphi_i = x_i$; $i = 1, \dots, n$. Let $\psi \in [\varphi_1, \dots, \varphi_n]$ have norm one, let ψ_j satisfy $\|\psi - \psi_j\| < \delta$ and suppose $\|T\psi\| = \|T\|$. Then

$$\|T\| \leq \|T(\psi_j)\| + \|T\| \|\psi - \psi_j\|.$$

Now, $\|T(\psi_j)\| = \|\sum \psi_j(f_i) x_i\| < 1 + \delta$ since (x_1, \dots, x_n) is in K_j . It follows from these inequalities that $\|T\| \leq (1 + \delta)/(1 - \delta)$. (This argument is similar to the same norm estimate in [7].) Since (x_1, \dots, x_n) is in D it follows that the matrix $B = (f_j(x_i) \mid i = 1, \dots, n; j = 1, \dots, n)$ is invertible for $\eta < 1$ (using the Neumann series). Let $A = (a_{ij})$ be its inverse so that $A = \sum (I - B)^k$. Now set $y_i = \sum_{j=1}^n a_{ij} x_j$ and let Δ be the linear extension of $\Delta x_i = y_i$ to all of $[x_1, \dots, x_n]$. It is easy to see that $f_i(y_j) = \delta_{ij}$. Now we estimate the norm of Δ . Let $W = \sum f_i(W) y_i$. Then

$$\|W - \Delta^{-1}(W)\| = \left\| \sum f_i(W)(y_i - x_i) \right\| \leq \|W\| \sum \|f_i\| \|y_i - x_i\|,$$

$$\begin{aligned} & \left(\sum_{i=1}^n \|f_i\| \left\| \left(\sum_{j=1}^n \delta_{ij} - a_{ij} \right) x_j \right\| \right) \|W\| \\ & \leq \left(\sum_{i,j=1}^n |\delta_{ij} - a_{ij}| \|f_i\| \|x_j\| \right) \|W\|, \\ & \leq \left(\left(\frac{\eta}{1 - \eta} \right) \|T\| \sum_{i,j=1}^n \|f_i\| \|\varphi_j\| \right) \|W\|, \\ & \leq \left(\left(\frac{\eta}{1 - \eta} \right) \left(\frac{1 + \delta}{1 - \delta} \right) \sum_{i,j=1}^n \|f_i\| \|\varphi_j\| \right) \|W\|, \\ & \leq \left(2 \left(\frac{\eta}{1 - \eta} \right) \sum_{i,j=1}^n \|f_i\| \|\varphi_j\| \right) \|W\|, \quad \text{for } 0 < \delta \leq \frac{1}{2}. \end{aligned}$$

Therefore, for any $K \in (0, 1)$ there exists $\eta_0 > 0$ such that $0 < \eta < \eta_0$ implies $\|W - \Delta^{-1}(W)\| < K \|W\|$. Thus, $\Delta = \sum (I - \Delta^{-1})^j$ so $\|\Delta\| \leq (1 - K)^{-1}$. Now we define $Ru = \sum f_i(u) y_i$. If J is the canonical map from X to X^{**} , one verifies directly that $R = \Delta TP^*J$, so that

$$\|R\| \leq (1/(1 - K))((1 + \delta)/(1 - \delta)) \|P\|.$$

By choosing η and δ small we get $\|R\| \leq \|P\| + \epsilon$. Further, it is clear that $R^*X^* = PX^*$. Now to check the final assertion,

$$\begin{aligned} \left\| (R^* - P) \sum_{j=1}^m \alpha_j f_j \right\| &= \left\| \sum_{j=n+1}^m \alpha_j R^* f_j \right\| = \left\| \sum_{j=n+1}^m \alpha_j \sum_{i=1}^n f_j(y_i) f_i \right\|, \\ &= \left\| \sum_{j=n+1}^m \sum_{i=1}^n \sum_{k=1}^n \alpha_j a_{ik} f_j(x_k) f_i \right\|, \\ &\leq \left(\sum_{j=n+1}^m \sum_{k=1}^n |f_j(x_k)| \right) \sum |\alpha_j| \sum |a_{ik}| \sum \|f_i\|, \end{aligned}$$

and the first term (being smaller than η) approaches 0 as $\eta \rightarrow 0$ for each $\sum_{j=1}^m \alpha_j f_j$. By choosing η smaller if necessary, the conclusion follows.

COROLLARY. *If every n -dimensional subspace of X^* is complemented with norm $< K_n$, then every subspace of X having deficiency n is complemented with norm $< 1 + K_n$.*

Proof. Let $U = [f_1, \dots, f_n]_{\perp}$, $P: X^* \rightarrow [f_1, \dots, f_n]$ having norm $< K_n$ and $\epsilon < K_n - \|P\|$. If R is the projection of the theorem, then $(I - R)X = (R^*X^*)_{\perp} = U$ and $\|I - R\| \leq 1 + \|R\| \leq 1 + \|P\| + \epsilon < 1 + K_n$.

This gives an affirmative solution to Problem 8 of [1].

Kadec has recently shown the following (see [2]): If Y is an n -dimensional subspace of (any Banach space) X , then there is a projection of X onto Y with norm $\leq n^{1/2}$. This allows the following refinement of Theorem 6 of [1]. (This result also appears in [2]).

COROLLARY. *If Y has deficiency n in X , and if $\epsilon > 0$ there is a projection of norm $< 1 + n^{1/2} + \epsilon$ of X onto Y .*

It is not known whether every Banach space has "nicely" complemented subspaces of arbitrarily large finite dimension. That is, given X , does there exist a constant M such that for every n there is a subspace U of X having dimension $\geq n$ and complemented with norm $\leq M$. The next corollary says that one may as well restrict his attention to conjugate spaces in studying this question.

COROLLARY. *If, for X^* , there is a constant M and subspaces V_n of X^* with $\dim V_n \geq n$ complemented with norm $\leq M$, then X contains subspaces U_n with $\dim U_n \geq n$ and complemented with norm $\leq M$.*

We must remark that if a finite-dimensional subspace is complemented with norm $\leq M + \epsilon$ for every $\epsilon > 0$, then it is complemented with norm $\leq M$.

3

The following is a direct application of the Lindenstrauss "compactness argument" (see e.g. [6]).

THEOREM 3. *Let X be an n -dimensional subspace of a Banach space Z and let P be a projection of least norm of Z onto X . Then $\|P\| = \sup_R \|R\|$ where R ranges over all "minimum norm" projections from W to X , W finite-dimensional, $X \subset W \subset Z$.*

Proof. Let $\mathcal{B} \subset 2^Z$ be the collection of all finite-dimensional superspaces of X partially ordered by inclusion. For each $B \in \mathcal{B}$, let P_B be a best (in terms of norm) projection of B onto X and extend P_B to all of Z by setting $P_{Bz} = 0$ if $z \in Z \setminus B$. By the Kadec result above, it follows that $\|P_{Bz}\| \leq (\dim X)^{1/2} \|z\|$ for every $z \in Z$. Now let

$$W = \prod_{z \in Z} n \|z\| B_x$$

which is compact in the product topology since X is n -dimensional. The net $(P_{Bz})_{z \in Z}$ is in W , and thus has a convergent subnet, say $(P_C(z))$. Thus, $P_C(z)$ converges in X for each z in Z . It is clear that, defining $P: Z \rightarrow X$ by $Pz = \lim P_C z$, P is bounded and $Px = x$ for all x in X . Also, for $z_1, z_2 \in Z$, and all $C \supset [z_1, z_2, X]$, $P_C(\alpha z_1 + \beta z_2) = \alpha P_C(z_1) + \beta P_C(z_2)$, so P is linear. Further $\|Pz\| \leq \{\lim_C \|P_C\|\} \|z\|$ giving the desired result.

Let X be n -dimensional. For any superspace W of X let $P(X, W)$ be the norm of the best projection of W onto X . Define

$$P_m(X) = \sup\{P(X, W) \mid \dim W = m\},$$

$$P(X) = \sup\{P(X, W) \mid W \supset X\}.$$

The affirmative solution to problem 9 of [1] is

COROLLARY. $P(X) = \sup P_m(X)$.

REFERENCES

1. E. W. CHENEY AND K. H. PRICE, "Minimal projections," *Approximation Theory* (A. Talbot, Ed.), pp. 261–289, Academic Press, New York, 1970.
2. D. J. H. GARLING AND Y. GORDON, "Relations between some constants associated with finite dimensional Banach spaces," *Is. J. Math.* **9** (1971), 346–361.
3. V. L. KLEE, "On certain intersection properties of convex sets," *Canad. J. Math.* **3** (1951), 272–275.
4. W. B. JOHNSON, "On the existence of strongly series summable Markushevich bases in Banach spaces," *Trans. A. M. S.* **157** (1971), 481–486.
5. W. B. JOHNSON, H. P. ROSENTHAL, AND M. ZIPPIN, "On bases finite dimensional decompositions and weaker structures in Banach spaces," *Is. J. Math.* **9** (1971), 488–506.
6. J. LINDENSTRAUSS, "On nonlinear projections in Banach spaces," *Michigan Math. J.* (1964), 263–287.
7. J. LINDENSTRAUSS AND H. P. ROSENTHAL, "The \mathcal{L}_p spaces," *Israel J. Math.* **7** (1969), 325–349.